# Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US2005/010912

International filing date: 31 March 2005 (31.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US

Number: 60/559,275


Filing date: 01 April 2004 (01.04.2004)

Date of receipt at the International Bureau: 12 August 2005 (12.08.2005)

Remark: Priority document submitted or transmitted to the International Bureau in

compliance with Rule 17.1(a) or (b)





### 

#### 'and and and vandamentess; presents; searce, comes;

#### UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

August 02, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

**APPLICATION NUMBER: 60/559,275** 

FILING DATE: April 01, 2004

RELATED PCT APPLICATION NUMBER: PCT/US05/10912

Certified by

Under Secretary of Commerce for Intellectual Property and Director of the United States Patent and Trademark Office

#### Abstract of the Disclosure

Provided herein are methods for identifying a risk of osteoarthritis in a subject, reagents and kits for carrying out the methods, methods for identifying candidate therapeutics for treating osteoarthritis, and therapeutic and preventative methods applicable to osteoarthritis. These embodiments are based upon an analysis of polymorphic variations in nucleotide sequences within the human genome.

#### **Application Data Sheet**

#### **Application Information**

Application Type:: Provisional

Subject Matter:: Utility

Suggested Group Art Unit:: Not Yet Assigned

CD-ROM or CD-R?:: None

Sequence submission?:: None

Computer Readable Form (CRF)?:: No

Title:: METHODS FOR IDENTIFYING RISK OF

OSTEOARTHRITIS AND TREATMENTS

**THEREOF** 

Attorney Docket Number:: 524593009000

Request for Early Publication?:: No

Request for Non-Publication?:: No

Total Drawing Sheets?:: 3

Small Entity?:: Yes

Petition included?::

Secrecy Order in Parent Appl.?:: No

#### **Applicant Information**

Applicant Authority Type:: Inventor

Primary Citizenship Country:: US

Status:: Full Capacity

Given Name:: Steven

Family Name:: MAH

City of Residence:: San Diego

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 12820 Via Nieve #74

City of mailing address:: San Diego

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92130

Applicant Authority Type:: Inventor

Primary Citizenship Country:: Germany

Status:: Full Capacity

Given Name:: Andreas

Family Name:: BRAUN

City of Residence:: San Diego

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 3935 Lago Di Grata Circle

City of mailing address:: San Diego

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92130

Applicant Authority Type:: Inventor

Primary Citizenship Country:: Germany

Status:: Full Capacity

Given Name:: Stefan

Middle Name:: M.

Family Name:: KAMMERER

City of Residence:: San Diego

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 3825 Elijah Court, Unit 334

City of mailing address:: San Diego

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92130

Applicant Authority Type:: Inventor

Primary Citizenship Country:: US

Status:: Full Capacity

Page # 2 Initial 04/01/04

Given Name:: Matthew

Middle Name:: Roberts

Family Name:: NELSON

City of Residence:: San Marcos

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 1250 Calle Prospero

City of mailing address:: San Marcos

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92069

Applicant Authority Type:: Inventor

Primary Citizenship Country:: Sweden

Status:: Full Capacity

Given Name:: Rikard

Middle Name:: Henry

Family Name:: RENELAND

City of Residence:: San Diego

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 7555 Charmant Drive, #1114

City of mailing address:: San Diego

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92122

Applicant Authority Type:: Inventor

Primary Citizenship Country:: United Kingdom

Status:: Full Capacity

Given Name:: Maria

Middle Name:: L.

Family Name:: LANGDOWN

City of Residence:: San Diego

Page # 3 Initial 04/01/04

State or Province of Residence:: CA

Country of Residence:: US

Street of mailing address:: 3701 Yosemite Street

City of mailing address:: San Diego

State or Province of mailing address:: CA

Postal or Zip Code of mailing address:: 92109

**Correspondence Information** 

Correspondence Customer Number:: 25225

Representative Information

Representative Customer Number:: 25225

#### What is claimed is:

- 1. A method for identifying a subject at risk of osteoarthritis, which comprises detecting the presence or absence of one or more polymorphic variations associated with osteoarthritis in a nucleic acid sample from a subject, wherein the one or more polymorphic variations are detected in a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
  - (d) a fragment of a nucleotide sequence of (a), (b), or (c);

whereby the presence of the polymorphic variation is indicative of the subject being at risk of osteoarthritis.

- 2. The method of claim 1, which further comprises obtaining the nucleic acid sample from the subject.
- 3. The method of claim 1, wherein the one or more polymorphic variations are detected within a region spanning chromosome positions 102570000 to 102583000 in human genomic DNA.
- 4. The method of claim 1, wherein the one or more polymorphic variations are detected at one or more positions selected from the group consisting of 207, 6019, 6414, 7341, 10984, 12351, 13335, 16584, 16737, 23897, 24057, 25145, 25300, 26262, 26312, 26589, 27302, 27358, 27451, 27552, 30731, 32085, 32139, 33184, 42382, 42569, 44823, 45217, 45548, 45601, 45722, 45967, 47367, 47642, 48126, 49218, 49274, 49433, 49610, 51282, 51466, 53757, 53960, 54031, 54574, 55679, 56100, 56182, 59817, 60533, 60656, 72209, 72778, 74293, 77335, 78029, 78374, 78421, 78434, 79174, 79397, 79562, 79700, 79730, 79904, 79920, 79938, 79972, 80125, 80368, 83484, 85536, 85829, 86425, 88083, 88770, 90622, 90924, 91634, 92029, 95152, 95348, 96145, 96793, 97015, 97064, 97711, 97855 and 98708.
- 5. The method of claim 1, wherein the one or more polymorphic variations are detected at one or more positions in SEQ ID NO: 1 selected from the group consisting of 6414, 51282, 54574, 78374, 92029 and 96793.

- 6. The method of claim 1, wherein the one or more polymorphic variations are detected at one or more positions in linkage disequilibrium with one or more positions in claim 3, 4 or 5.
- 7. The method of claim 1, wherein detecting the presence or absence of the one or more polymorphic variations comprises:

hybridizing an oligonucleotide to the nucleic acid sample, wherein the oligonucleotide is complementary to a nucleotide sequence in the nucleic acid and hybridizes to a region adjacent to the polymorphic variation;

extending the oligonucleotide in the presence of one or more nucleotides, yielding extension products; and

detecting the presence or absence of a polymorphic variation in the extension products.

- 8. The method of claim 1, wherein the subject is a human.
- 9. The method of claim 8, wherein the subject is a human female.
- 10. The method of claim 8, wherein the subject is a human male.
- 11. A method for identifying a polymorphic variation associated with osteoarthritis proximal to an incident polymorphic variation associated with osteoarthritis, which comprises:

identifying a polymorphic variation proximal to the incident polymorphic variation associated with osteoarthritis, wherein the polymorphic variation is detected in a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising a polymorphic variation;

determining the presence or absence of an association of the proximal polymorphic variant with osteoarthritis.

12. The method of claim 11, wherein the incident polymorphic variation is at one or more positions in claim 3, 4 or 5.

- 13. The method of claim 11, wherein the proximal polymorphic variation is within a region between about 5 kb 5' of the incident polymorphic variation and about 5 kb 3' of the incident polymorphic variation.
- 14. The method of claim 11, which further comprises determining whether the proximal polymorphic variation is in linkage disequilibrium with the incident polymorphic variation.
- 15. The method of claim 11, which further comprises identifying a second polymorphic variation proximal to the identified proximal polymorphic variation associated with osteoarthritis and determining if the second proximal polymorphic variation is associated with osteoarthritis.
- 16. The method of claim 15, wherein the second proximal polymorphic variant is within a region between about 5 kb 5' of the incident polymorphic variation and about 5 kb 3' of the proximal polymorphic variation associated with osteoarthritis.
- 17. An isolated nucleic acid comprising a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising a polymorphic variation; and
- (e) a nucleotide sequence complementary to the nucleotide sequences of (a), (b), (c), or (d); wherein the nucleotide sequence comprises a polymorphic variation associated with osteoarthritis selected from the group consisting of an adenine at position 6414, an adenine at position 51282, a cytosine at position 54574, a thymine at position 92029 and an adenine at position 96793.
- 18. An oligonucleotide comprising a nucleotide sequence complementary to a portion of the nucleotide sequence of (a), (b), (c), or (d) in claim 17, wherein the 3' end of the oligonucleotide is adjacent to a polymorphic variation associated with osteoarthritis.
  - 19. A microarray comprising an isolated nucleic acid of claim 17 linked to a solid support.

- 20. An isolated polypeptide encoded by the isolated nucleic acid sequence of claim 17.
- 21. A method for identifying a candidate therapeutic for treating osteoarthritis, which comprises:
- (a) introducing a test molecule to a system which comprises a nucleic acid comprising a nucleotide sequence selected from the group consisting of:
  - (i) a nucleotide sequence in SEQ ID NO: 1-4;
- (ii) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (iii) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
  - (iv) a fragment of a nucleotide sequence of (a), (b), or (c); or introducing a test molecule to a system which comprises a protein encoded by a nucleotide

sequence of (i), (ii), or (iv); and

(b) determining the presence or absence of an interaction between the test molecule and the nucleic acid or protein,

whereby the presence of an interaction between the test molecule and the nucleic acid or protein identifies the test molecule as a candidate therapeutic for treating osteoarthritis.

- 22. The method of claim 21, wherein the system is an animal.
- 23. The method of claim 21, wherein the system is a cell.
- 24. The method of claim 21, wherein the nucleotide sequence comprises one or more polymorphic variations associated with osteoarthritis.
- 25. The method of claim 24, wherein the one or more polymorphic variations associated with osteoarthritis are at one or more positions in claim 3, 4 or 5.
- 26. A method for treating osteoarthritis in a subject, which comprises contacting one or more cells of a subject in need thereof with a nucleic acid, wherein the nucleic acid comprises a nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;

- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
  - (d) a fragment of a nucleotide sequence of (a), (b), or (c); and
- (e) a nucleotide sequence complementary to the nucleotide sequences of (a), (b), (c), or (d); whereby contacting the one or more cells of the subject with the nucleic acid treats the osteoarthritis in the subject.
  - 27. The method of claim 26, wherein the nucleic acid is RNA or PNA.
  - 28. The method of claim 27, wherein the nucleic acid is duplex RNA.
- 29. A method for treating osteoarthritis in a subject, which comprises contacting one or more cells of a subject in need thereof with a protein, wherein the protein is encoded by a nucleotide sequence which comprises a polynucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
  - (d) a fragment of a nucleotide sequence of (a), (b), or (c);

whereby contacting the one or more cells of the subject with the protein treats the osteoarthritis in the subject.

30. A method for treating osteoarthritis in a subject, which comprises:

detecting the presence or absence of one or more polymorphic variations associated with osteoarthritis in a nucleic acid sample from a subject, wherein the one or more polymorphic variation are detected in a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising a polymorphic variation; and

administering an osteoarthritis treatment to a subject in need thereof based upon the presence or absence of the one or more polymorphic variations in the nucleic acid sample.

- 31. The method of claim 30, wherein the one or more polymorphic variations are detected at one or more positions in claim 3, 4 or 5.
- 32. The method of claim 30, wherein the treatment is selected from the group consisting of administering a corticosteroid, a corticosteroid, a nonsteroidal anti-inflammatory drug (NSAID), a cyclooxygenase-2 (COX-2) inhibitor, an antibody, a glucocorticoid, hyaluronic acid, chondrotin sulfate, glucosamine or acetaminophen; prescribing a heat/cold regimen or a joint protection regimen; performing joint surgery; prescribing a weight control regimen; and combinations of the foregoing.
- 33. A method for detecting or preventing osteoarthritis in a subject, which comprises:

  detecting the presence or absence of one or more polymorphic variations associated with
  osteoarthritis in a nucleic acid sample from a subject, wherein the polymorphic variation is detected in a
  nucleotide sequence selected from the group consisting of:
  - (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising a polymorphic variation; and

administering an osteoarthritis prevention or detection procedure to a subject in need thereof based upon the presence or absence of the one or more polymorphic variations in the nucleic acid sample.

- 34. The method of claim 33, wherein the one or more polymorphic variations are detected at one or more positions in claim 3, 4 or 5.
- 35. The method of claim 33, wherein the osteoarthritis prevention is selected from the group consisting of administering a corticosteroid, a nonsteroidal anti-inflammatory drug (NSAID), a cyclooxygenase-2 (COX-2) inhibitor, an antibody, a glucocorticoid, hyaluronic acid, chondrotin sulfate, glucosamine or acetaminophen; prescribing a heat/cold regimen or a joint protection regimen; performing joint surgery; prescribing a weight control regimen; and combinations of the foregoing.

36. A method of targeting information for preventing or treating osteoarthritis to a subject in need thereof, which comprises:

detecting the presence or absence of one or more polymorphic variations associated with osteoarthritis in a nucleic acid sample from a subject, wherein the polymorphic variation is detected in a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence in SEQ ID NO: 1-4;
- (b) a nucleotide sequence which encodes a polypeptide encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to the amino acid sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4;
- (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising a polymorphic variation; and

directing information for preventing or treating osteoarthritis to a subject in need thereof based upon the presence or absence of the one or more polymorphic variations in the nucleic acid sample.

- 37. The method of claim 36, wherein the one or more polymorphic variations are detected at one or more positions in claim 3, 4 or 5.
- 38. A composition comprising a cell from a subject having osteoarthritis or at risk of osteoarthritis and an antibody that specifically binds to a protein, polypeptide or peptide encoded by a nucleotide sequence identical to or 90% or more identical to a nucleotide sequence in SEQ ID NO: 1-4.
- 39. A composition comprising a cell from a subject having osteoarthritis or at risk of osteoarthritis and a RNA, DNA, PNA or ribozyme molecule comprising a nucleotide sequence identical to or 90% or more identical to a portion of a nucleotide sequence in SEQ ID NO: 1-4.
- 40. The composition of claim 39, wherein the RNA molecule is a short inhibitory RNA molecule.

App No.: Not Yet Assigned

Docket No.: 524593009000


Inventor: Steven MAH, et al.

Title: METHODS FOR IDENTIFYING RISK OF


OSTEOARTHRITIS AND TREATMENTS THEREOF

#### FIGURE 1

#### IL1RL1 - DISCOVERY P-VALUES (female only)



#### Chromosome position



App No.: Not Yet Assigned Docket No.: 524593009000

Inventor: Steven MAH, et al.


Title: METHODS FOR IDENTIFYING RISK OF

OSTEOARTHRITIS AND TREATMENTS THEREOF

#### FIGURE 2

IL1RL1 isoform expression (SW1353 monolayers)





MMP8

MMP13

**BVES** 

CHCD1

IL-1RL1 (TM)

IL-1RL1 (Soluble, type 2 or 3)

App No.: Not Yet Assigned


Docket No.: 524593009000

Inventor: Steven MAH, et al.


Title: METHODS FOR IDENTIFYING RISK OF OSTEOARTHRITIS AND TREATMENTS THEREOF

#### FIGURE 3A

#### IL1RL1 isoform expression (3-D alginate)



# FIGURE 3B IL1RL1 isoform expression (3-D alginate)



## METHODS FOR IDENTIFYING RISK OF OSTEOARTHRITIS AND TREATMENTS THEREOF

#### Field of the Invention

[0001] The invention relates to genetic methods for identifying risk of osteoarthritis and treatments that specifically target such diseases.

#### **Background**

[0002] Osteoarthritis (OA) is a chronic disease usually affecting weight-bearing synovial joints. There are approximately 20 million Americans affected by OA and it is the leading cause of disability in the United States. In addition to extensive human suffering, OA also accounts for nearly all knee replacements and more than half of all hip replacements in the United States. Despite its prevalence, OA is poorly understood and there are few treatments available besides anti-inflammatory drugs and joint replacement.

[0003] Most commonly affecting middle-aged and older people, OA can range from very mild to very severe. It affects hands and weight-bearing joints such as knees, hips, feet and the back. Knee OA can be as disabling as any cardiovascular disease except stroke.

[0004] OA is characterized by the breakdown of cartilage in joints. Cartilage in joints cushions the ends of bones, and cartilage breakdown causes bones to rub against each other, causing pain and loss of movement. Type II collagen is the main component of cartilage, comprising 15-25% of the wet weight, approximately half the dry weight, and representing 90-95% of the total collagen content in the tissue. It forms fibrils that endow cartilage with tensile strength (Mayne, R. Arthritis Rhuem. 32:241-246 (1989)).

#### Summary

[0005] It has been discovered that certain polymorphic variations in human genomic DNA are associated with osteoarthritis. In particular, polymorphic variants in a locus containing a *IL1RL1* region in human genomic DNA have been associated with risk of osteoarthritis.

[0006] Thus, featured herein are methods for identifying a subject at risk of osteoarthritis and/or a risk of osteoarthritis in a subject, which comprise detecting the presence or absence of one or more polymorphic variations associated with osteoarthritis in or around the loci described herein in a human nucleic acid sample. In an embodiment, two or more polymorphic variations are detected in two or more regions of which one is the *IL1RL1* region. In certain embodiments, 3 or more, or 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 or more polymorphic variants are detected.

[0007] Also featured are nucleic acids that include one or more polymorphic variations associated with occurrence of osteoarthritis, as well as polypeptides encoded by these nucleic acids. In addition, provided are methods for identifying candidate therapeutic molecules for treating osteoarthritis, as well as methods for treating osteoarthritis in a subject by identifying a subject at risk of osteoarthritis and treating the subject with a suitable prophylactic, treatment or therapeutic molecule.

[0008] Also provided are compositions comprising a cell from a subject having osteoarthritis or at risk of osteoarthritis and/or a *IL1RL1* nucleic acid, with a RNAi, siRNA, antisense DNA or RNA, or ribozyme nucleic acid designed from a *IL1RL1* nucleotide sequence. In an embodiment, the RNAi, siRNA, antisense DNA or RNA, or ribozyme nucleic acid is designed from a *IL1RL1* nucleotide sequence that includes one or more polymorphic variations associated with osteoarthritis, and in some instances, specifically interacts with such a nucleotide sequence. Further, provided are arrays of nucleic acids bound to a solid surface, in which one or more nucleic acid molecules of the array have a *IL1RL1* nucleotide sequence, or a fragment or substantially identical nucleic acid thereof, or a complementary nucleic acid of the foregoing. Featured also are compositions comprising a cell from a subject having osteoarthritis or at risk of osteoarthritis and/or a *IL1RL1* polypeptide, with an antibody that specifically binds to the polypeptide. Thus, featured is an antibody that specifically binds to an epitope in the polypeptide that includes an amino acid encoded by a polymorphic site associated with osteoarthritis. In certain embodiments, the antibody specifically binds to an epitope comprising a glutamate or alanine encoded by rs1041973 (e.g., an antibody that binds to an epitope comprising an alanine at position 78 in an *IL1RL1* polypeptide) An alanine at position 78 is associated with increased risk of osteoarthitis.

#### Brief Description of the Drawings

[0009] Figure 1 shows proximal SNPs in a *IL1RL1* region in genomic DNA. The position of each SNP in the chromosome is shown on the x-axis and the y-axis provides the negative logarithm of the p-value comparing the estimated allele to that of the control group. Also shown in the figure are exons and introns of the gene in the approximate chromosomal positions.

[0010] Figure 2 shows expression profiling results for *IL1RL1*.

[0011] Figures 3A and 3B show *IL1RL1* expression modulation in a human chondrocyte cell line model.

#### **Detailed Description**

[0012] It has been discovered that a polymorphic variant in a locus containing a *IL1RL1* region is associated with occurrence of osteoarthritis in subjects. Thus, detecting genetic determinants associated with an increased risk of osteoarthritis occurrence can lead to early identification of a predisposition to osteoarthritis and early prescription of preventative measures. Also, associating a *IL1RL1* polymorphic

variant with osteoarthritis has provided new targets for screening molecules useful in treatments of osteoarthritis.

#### Osteoarthritis and Sample Selection

[0013] Osteoarthritis (OA), or degenerative joint disease, is one of the oldest and most common types of arthritis. It is characterized by the breakdown of the joint's cartilage. Cartilage is the part of the joint that cushions the ends of bones, and its breakdown causes bones to rub against each other, causing pain and loss of movement. Type II collagen is the main component of cartilage, comprising 15-25% of the wet weight, approximately half the dry weight, and representing 90-95% of the total collagen content in the tissue. It forms fibrils that endow cartilage with tensile strength (Mayne, R. Arthritis Rhuem. 32:241-246 (1989)).

[0014] Most commonly affecting middle-aged and older people, OA can range from very mild to very severe. It affects hands and weight-bearing joints such as knees, hips, feet and the back. Knee OA can be as disabling as any cardiovascular disease except stroke.

[0015] Osteoarthritis affects an estimated 20.7 million Americans, mostly after age 45, with women more commonly affected than men. Physicians make a diagnosis of OA based on a physical exam and history of symptoms. X-rays are used to confirm diagnosis. Most people over 60 reflect the disease on X-ray, and about one-third have actual symptoms.

[0016] There are many factors that can cause OA. Obesity may lead to osteoarthritis of the knees. In addition, people with joint injuries due to sports, work-related activity or accidents may be at increased risk of developing OA.

[0017] Genetics has a role in the development of OA. Some people may be born with defective cartilage or with slight defects in the way that joints fit together. As a person ages, these defects may cause early cartilage breakdown in the joint or the inability to repair damaged or deteriorated cartilage in the joint.

[0018] Inclusion or exclusion of samples for an osteoarthritis pool may be based upon the following criteria: ethnicity (e.g., samples derived from an individual characterized as Caucasian); parental ethnicity (e.g., samples derived from an individual of British paternal and maternal descent); relevant phenotype information for the individual (e.g., case samples derived from individuals diagnosed with specific knee osteoarthritis (OA) and were recruited from an OA knee replacement clinic). Control samples may be selected based on relevant phenotype information for the individual (e.g., derived from individuals free of OA at several sites (knee, hand, hip etc)); and no family history of OA and/or rheumatoid arthritis. Additional phenotype information collected for both cases and controls may include age of the individual, gender, family history of OA, diagnosis with osteoarthritis (joint location of OA, date of primary diagnosis, age of individual as of primary diagnosis), knee history (current symptoms,

any major knee injury, menisectomy, knee replacement surgery, age of surgery), HRT history, osteoporosis diagnosis.

[0019] Based in part upon selection criteria set forth above, individuals having osteoarthritis can be selected for genetic studies. Also, individuals having no history of osteoarthritis often are selected for genetic studies, as described hereafter.

#### Polymorphic Variants Associated with Osteoarthritis

[0020] A genetic analysis provided herein linked osteoarthritis with polymorphic variant nucleic acid sequences in the human genome. As used herein, the term "polymorphic site" refers to a region in a nucleic acid at which two or more alternative nucleotide sequences are observed in a significant number of nucleic acid samples from a population of individuals. A polymorphic site may be a nucleotide sequence of two or more nucleotides, an inserted nucleotide or nucleotide sequence, a deleted nucleotide or nucleotide sequence, or a microsatellite, for example. A polymorphic site that is two or more nucleotides in length may be 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or more, 20 or more, 30 or more, 50 or more, 75 or more, 100 or more, 500 or more, or about 1000 nucleotides in length, where all or some of the nucleotide sequences differ within the region. A polymorphic site is often one nucleotide in length, which is referred to herein as a "single nucleotide polymorphism" or a "SNP."

[0021] Where there are two, three, or four alternative nucleotide sequences at a polymorphic site, each nucleotide sequence is referred to as a "polymorphic variant" or "nucleic acid variant." Where two polymorphic variants exist, for example, the polymorphic variant represented in a minority of samples from a population is sometimes referred to as a "minor allele" and the polymorphic variant that is more prevalently represented is sometimes referred to as a "major allele." Many organisms possess a copy of each chromosome (e.g., humans), and those individuals who possess two major alleles or two minor alleles are often referred to as being "homozygous" with respect to the polymorphism, and those individuals who possess one major allele and one minor allele are normally referred to as being "heterozygous" with respect to the polymorphism. Individuals who are homozygous with respect to one allele are sometimes predisposed to a different phenotype as compared to individuals who are heterozygous or homozygous with respect to another allele.

[0022] In genetic analysis that associate polymorphic variants with osteoarthritis, samples from individuals having osteoarthritis and individuals not having osteoarthritis often are allelotyped and/or genotyped. The term "allelotype" as used herein refers to a process for determining the allele frequency for a polymorphic variant in pooled DNA samples from cases and controls. By pooling DNA from each group, an allele frequency for each SNP in each group is calculated. These allele frequencies are then compared to one another. The term "genotyped" as used herein refers to a process for determining a

genotype of one or more individuals, where a "genotype" is a representation of one or more polymorphic variants in a population.

[0023] A genotype or polymorphic variant may be expressed in terms of a "haplotype," which as used herein refers to two or more polymorphic variants occurring within genomic DNA in a group of individuals within a population. For example, two SNPs may exist within a gene where each SNP position includes a cytosine variation and an adenine variation. Certain individuals in a population may carry one allele (heterozygous) or two alleles (homozygous) having the gene with a cytosine at each SNP position. As the two cytosines corresponding to each SNP in the gene travel together on one or both alleles in these individuals, the individuals can be characterized as having a cytosine/cytosine haplotype with respect to the two SNPs in the gene.

[0024] As used herein, the term "phenotype" refers to a trait which can be compared between individuals, such as presence or absence of a condition, a visually observable difference in appearance between individuals, metabolic variations, physiological variations, variations in the function of biological molecules, and the like. An example of a phenotype is occurrence of osteoarthritis.

[0025] Researchers sometimes report a polymorphic variant in a database without determining whether the variant is represented in a significant fraction of a population. Because a subset of these reported polymorphic variants are not represented in a statistically significant portion of the population, some of them are sequencing errors and/or not biologically relevant. Thus, it is often not known whether a reported polymorphic variant is statistically significant or biologically relevant until the presence of the variant is detected in a population of individuals and the frequency of the variant is determined. Methods for detecting a polymorphic variant in a population are described herein, specifically in Example 2. A polymorphic variant is statistically significant and often biologically relevant if it is represented in 5% or more of a population, sometimes 10% or more, 15% or more, or 20% or more of a population, and often 25% or more, 30% or more, 35% or more, 40% or more, 45% or more, or 50% or more of a population.

[0026] A polymorphic variant may be detected on either or both strands of a double-stranded nucleic acid. Also, a polymorphic variant may be located within an intron or exon of a gene or within a portion of a regulatory region such as a promoter, a 5' untranslated region (UTR), a 3' UTR, and in DNA (e.g., genomic DNA (gDNA) and complementary DNA (cDNA)), RNA (e.g., mRNA, tRNA, and rRNA), or a polypeptide. Polymorphic variations may or may not result in detectable differences in gene expression, polypeptide structure, or polypeptide function.

[0027] It was determined that polymorphic variations associated with an increased risk of osteoarthritis existed in a *IL1RL1* region in SEQ ID NO: 1. In certain embodiments, a polymorphic variant at position rs1041973 in the human genome was associated with an increased risk of osteoarthritis, and in a specific embodiment, a cytosine at position rs1041973 was associated with an increased risk of osteoarthritis.

[0028] Polymorphic variants in and around the *IL1RL1* locus were tested for association with osteoarthritis. These include polymorphic variants at positions in SEQ ID NO: 1 selected from the group consisting of 207, 6019, 6414, 7341, 10984, 12351, 13335, 16584, 16737, 23897, 24057, 25145, 25300, 26262, 26312, 26589, 27302, 27358, 27451, 27552, 30731, 32085, 32139, 33184, 42382, 42569, 44823, 45217, 45548, 45601, 45722, 45967, 47367, 47642, 48126, 49218, 49274, 49433, 49610, 51282, 51466, 53757, 53960, 54031, 54574, 55679, 56100, 56182, 59817, 60533, 60656, 72209, 72778, 74293, 77335, 78029, 78374, 78421, 78434, 79174, 79397, 79562, 79700, 79730, 79904, 79920, 79938, 79972, 80125, 80368, 83484, 85536, 85829, 86425, 88083, 88770, 90622, 90924, 91634, 92029, 95152, 95348, 96145, 96793, 97015, 97064, 97711, 97855 and 98708. Polymorphic variants at the following positions in SEQ ID NO: 1 in particular were associated with an increased risk of osteoarthritis: 6414, 51282, 54574, 78374, 92029 and 96793, where specific embodiments are directed to position 54574. In particular, the following polymorphic variants in SEQ ID NO: 1 were associated with risk of osteoarthritis: an adenine at position 6414, an adenine at position 51282, a cytosine at position 54574, a thymine at position 92029 and an adenine at position 96793.

[0029] Based in part upon analyses summarized in Figure 1, a region with significant association has been identified in a locus associated with osteoarthritis. Any polymorphic variants associated with osteoarthritis in a region of significant association can be utilized for embodiments described herein. For example, polymorphic variants in a region spanning chromosome positions 102570000 to 102583000 (approximately 13,000 nucleotides in length) in a *IL1RL1* locus have significant association (chromosome positions are within NCBI's Genome build 34).

#### Additional Polymorphic Variants Associated with Osteoarthritis

[0030] Also provided is a method for identifying polymorphic variants proximal to an incident, founder polymorphic variant associated with osteoarthritis. Thus, featured herein are methods for identifying a polymorphic variation associated with osteoarthritis that is proximal to an incident polymorphic variation associated with osteoarthritis, which comprises identifying a polymorphic variant proximal to the incident polymorphic variant associated with osteoarthritis, where the incident polymorphic variant is in a *IL1RL1* nucleotide sequence. The nucleotide sequence often comprises a polynucleotide sequence selected from the group consisting of (a) a polynucleotide sequence of SEQ ID NO: 1-4; (b) a polynucleotide sequence that encodes a polypeptide having an amino acid sequence encoded by a polynucleotide sequence of SEQ ID NO: 1-4; and (c) a polynucleotide sequence that encodes a polypeptide having an amino acid sequence that is 90% or more identical to an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4 or a polynucleotide sequence 90% or more identical to the polynucleotide sequence of SEQ ID NO: 1-4. The presence or absence of an association of the proximal polymorphic variant with osteoarthritis then is determined using a known

association method, such as a method described in the Examples hereafter. In an embodiment, the incident polymorphic variant is a polymorphic variant associated with osteoarthritis described herein. In another embodiment, the proximal polymorphic variant identified sometimes is a publicly disclosed polymorphic variant, which for example, sometimes is published in a publicly available database. In other embodiments, the polymorphic variant identified is not publicly disclosed and is discovered using a known method, including, but not limited to, sequencing a region surrounding the incident polymorphic variant in a group of nucleic samples. Thus, multiple polymorphic variants proximal to an incident polymorphic variant are associated with osteoarthritis using this method.

[0031] The proximal polymorphic variant often is identified in a region surrounding the incident polymorphic variant. In certain embodiments, this surrounding region is about 50 kb flanking the first polymorphic variant (*e.g.* about 50 kb 5' of the first polymorphic variant and about 50 kb 3' of the first polymorphic variant), and the region sometimes is composed of shorter flanking sequences, such as flanking sequences of about 40 kb, about 30 kb, about 25 kb, about 20 kb, about 15 kb, about 10 kb, about 7 kb, about 5 kb, or about 2 kb 5' and 3' of the incident polymorphic variant. In other embodiments, the region is composed of longer flanking sequences, such as flanking sequences of about 55 kb, about 60 kb, about 65 kb, about 70 kb, about 75 kb, about 80 kb, about 85 kb, about 90 kb, about 95 kb, or about 100 kb 5' and 3' of the incident polymorphic variant.

[0032] In certain embodiments, polymorphic variants associated with osteoarthritis are identified iteratively. For example, a first proximal polymorphic variant is associated with osteoarthritis using the methods described above and then another polymorphic variant proximal to the first proximal polymorphic variant is identified (e.g., publicly disclosed or discovered) and the presence or absence of an association of one or more other polymorphic variants proximal to the first proximal polymorphic variant with osteoarthritis is determined.

[0033] The methods described herein are useful for identifying or discovering additional polymorphic variants that may be used to further characterize a gene, region or loci associated with a condition, a disease (e.g., osteoarthritis), or a disorder. For example, allelotyping or genotyping data from the additional polymorphic variants may be used to identify a functional mutation or a region of linkage disequilibrium. In certain embodiments, polymorphic variants identified or discovered within a region comprising the first polymorphic variant associated with osteoarthritis are genotyped using the genetic methods and sample selection techniques described herein, and it can be determined whether those polymorphic variants are in linkage disequilibrium with the first polymorphic variant. The size of the region in linkage disequilibrium with the first polymorphic variant also can be assessed using these genotyping methods. Thus, provided herein are methods for determining whether a polymorphic variant is in linkage disequilibrium with a first polymorphic variant associated with osteoarthritis, and such information can be used in prognosis/diagnosis methods described herein.

#### Isolated Nucleic Acids

[0034] Featured herein are isolated *IL1RL1* nucleic acid variants depicted in SEQ ID NO: 1-4, and substantially identical nucleic acids thereof. A nucleic acid variant may be represented on one or both strands in a double-stranded nucleic acid or on one chromosomal complement (heterozygous) or both chromosomal complements (homozygous).

[0035] As used herein, the term "nucleic acid" includes DNA molecules (e.g., a complementary DNA (cDNA) and genomic DNA (gDNA)) and RNA molecules (e.g., mRNA, rRNA, siRNA and tRNA) and analogs of DNA or RNA, for example, by use of nucleotide analogs. The nucleic acid molecule can be single-stranded and it is often double-stranded. The term "isolated or purified nucleic acid" refers to nucleic acids that are separated from other nucleic acids present in the natural source of the nucleic acid. For example, with regard to genomic DNA, the term "isolated" includes nucleic acids which are separated from the chromosome with which the genomic DNA is naturally associated. An "isolated" nucleic acid is often free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and/or 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of 5' and/or 3' nucleotide sequences which flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. As used herein, the term "gene" refers to a nucleotide sequence that encodes a polypeptide.

[0036] Also included herein are nucleic acid fragments. These fragments often have a nucleotide sequence identical to a nucleotide sequence of SEQ ID NO: 1-4, a nucleotide sequence substantially identical to a nucleotide sequence of SEQ ID NO: 1-4, or a nucleotide sequence that is complementary to the foregoing. The nucleic acid fragment may be identical, substantially identical or homologous to a nucleotide sequence in an exon or an intron in a nucleotide sequence of SEQ ID NO: 1-4, and may encode a domain or part of a domain of a polypeptide. Sometimes, the fragment will comprises one or more of the polymorphic variations described herein as being associated with osteoarthritis. The nucleic acid fragment is often 50, 100, or 200 or fewer base pairs in length, and is sometimes about 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 2000, 3000, 4000, 5000, 10000, 15000, or 20000 base pairs in length. A nucleic acid fragment that is complementary to a nucleotide sequence identical or substantially identical to a nucleotide sequence in SEQ ID NO: 1-4 and hybridizes to such a nucleotide sequence under stringent conditions is often referred to as a "probe." Nucleic acid fragments often include one or more polymorphic sites, or sometimes have an end that is adjacent to a polymorphic site as described hereafter.

[0037] An example of a nucleic acid fragment is an oligonucleotide. As used herein, the term "oligonucleotide" refers to a nucleic acid comprising about 8 to about 50 covalently linked nucleotides, often comprising from about 8 to about 35 nucleotides, and more often from about 10 to about 25 nucleotides. The backbone and nucleotides within an oligonucleotide may be the same as those of naturally occurring nucleic acids, or analogs or derivatives of naturally occurring nucleic acids, provided that oligonucleotides having such analogs or derivatives retain the ability to hybridize specifically to a nucleic acid comprising a targeted polymorphism. Oligonucleotides described herein may be used as hybridization probes or as components of prognostic or diagnostic assays, for example, as described herein.

[0038] Oligonucleotides are typically synthesized using standard methods and equipment, such as the ABI™3900 High Throughput DNA Synthesizer and the EXPEDITE™ 8909 Nucleic Acid Synthesizer, both of which are available from Applied Biosystems (Foster City, CA). Analogs and derivatives are exemplified in U.S. Pat. Nos. 4,469,863; 5,536,821; 5,541,306; 5,637,683; 5,637,684; 5,700,922; 5,717,083; 5,719,262; 5,739,308; 5,773,601; 5,886,165; 5,929,226; 5,977,296; 6,140,482; WO 00/56746; WO 01/14398, and related publications. Methods for synthesizing oligonucleotides comprising such analogs or derivatives are disclosed, for example, in the patent publications cited above and in U.S. Pat. Nos. 5,614,622; 5,739,314; 5,955,599; 5,962,674; 6,117,992; in WO 00/75372; and in related publications.

[0039] Oligonucleotides may also be linked to a second moiety. The second moiety may be an additional nucleotide sequence such as a tail sequence (e.g., a polyadenosine tail), an adapter sequence (e.g., phage M13 universal tail sequence), and others. Alternatively, the second moiety may be a non-nucleotide moiety such as a moiety which facilitates linkage to a solid support or a label to facilitate detection of the oligonucleotide. Such labels include, without limitation, a radioactive label, a fluorescent label, a chemiluminescent label, a paramagnetic label, and the like. The second moiety may be attached to any position of the oligonucleotide, provided the oligonucleotide can hybridize to the nucleic acid comprising the polymorphism.

#### Uses for Nucleic Acid Sequence

[0040] Nucleic acid coding sequences may be used for diagnostic purposes for detection and control of polypeptide expression. Also, included herein are oligonucleotide sequences such as antisense RNA, small-interfering RNA (siRNA) and DNA molecules and ribozymes that function to inhibit translation of a polypeptide. Antisense techniques and RNA interference techniques are known in the art and are described herein.

[0041] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme

molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, hammerhead motif ribozyme molecules may be engineered that specifically and efficiently catalyze endonucleolytic cleavage of RNA sequences corresponding to or complementary to *IL1RL1* nucleotide sequences. Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences, GUA, GUU and GUC. Once identified, short RNA sequences of between fifteen (15) and twenty (20) ribonucleotides corresponding to the region of the target gene containing the cleavage site may be evaluated for predicted structural features such as secondary structure that may render the oligonucleotide sequence unsuitable. The suitability of candidate targets may also be evaluated by testing their accessibility to hybridization with complementary oligonucleotides, using ribonuclease protection assays.

[0042] Antisense RNA and DNA molecules, siRNA and ribozymes may be prepared by any method known in the art for the synthesis of RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides well known in the art such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

[0043] DNA encoding a polypeptide also may have a number of uses for the diagnosis of diseases, including osteoarthritis, resulting from aberrant expression of a target gene described herein. For example, the nucleic acid sequence may be used in hybridization assays of biopsies or autopsies to diagnose abnormalities of expression or function (e.g., Southern or Northern blot analysis, in situ hybridization assays).

[0044] In addition, the expression of a polypeptide during embryonic development may also be determined using nucleic acid encoding the polypeptide. As addressed, *infra*, production of functionally impaired polypeptide is the cause of various disease states, such as osteoarthritis. *In situ* hybridizations using polypeptide as a probe may be employed to predict problems related to osteoarthritis. Further, as indicated, *infra*, administration of human active polypeptide, recombinantly produced as described herein, may be used to treat disease states related to functionally impaired polypeptide. Alternatively, gene therapy approaches may be employed to remedy deficiencies of functional polypeptide or to replace or compete with dysfunctional polypeptide.

#### Expression Vectors, Host Cells, and Genetically Engineered Cells

[0045] Provided herein are nucleic acid vectors, often expression vectors, which contain a *IL1RL1* nucleotide sequence, or a substantially identical sequence thereof. As used herein, the term "vector"

refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked and can include a plasmid, cosmid, or viral vector. The vector can be capable of autonomous replication or it can integrate into a host DNA. Viral vectors may include replication defective retroviruses, adenoviruses and adeno-associated viruses for example.

[0046] A vector can include a *IL1RL1* nucleotide sequence in a form suitable for expression of an encoded target polypeptide or target nucleic acid in a host cell. A "target polypeptide" is a polypeptide encoded by a *IL1RL1* nucleotide sequence, or a substantially identical nucleotide sequence thereof. The recombinant expression vector typically includes one or more regulatory sequences operatively linked to the nucleic acid sequence to be expressed. The term "regulatory sequence" includes promoters, enhancers and other expression control elements (*e.g.*, polyadenylation signals). Regulatory sequences include those that direct constitutive expression of a nucleotide sequence, as well as tissue-specific regulatory and/or inducible sequences. The design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of polypeptide desired, and the like. Expression vectors can be introduced into host cells to produce target polypeptides, including fusion polypeptides.

[0047] Recombinant expression vectors can be designed for expression of target polypeptides in prokaryotic or eukaryotic cells. For example, target polypeptides can be expressed in *E. coli*, insect cells (e.g., using baculovirus expression vectors), yeast cells, or mammalian cells. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology 185*, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

[0048] Expression of polypeptides in prokaryotes is most often carried out in *E. coli* with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion polypeptides. Fusion vectors add a number of amino acids to a polypeptide encoded therein, usually to the amino terminus of the recombinant polypeptide. Such fusion vectors typically serve three purposes:

1) to increase expression of recombinant polypeptide; 2) to increase the solubility of the recombinant polypeptide; and 3) to aid in the purification of the recombinant polypeptide by acting as a ligand in affinity purification. Often, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant polypeptide to enable separation of the recombinant polypeptide from the fusion moiety subsequent to purification of the fusion polypeptide. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase. Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith & Johnson, *Gene 67:* 31-40 (1988)), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ) which fuse glutathione S-transferase (GST), maltose E binding polypeptide, or polypeptide A, respectively, to the target recombinant polypeptide.

[0049] Purified fusion polypeptides can be used in screening assays and to generate antibodies specific for target polypeptides. In a therapeutic embodiment, fusion polypeptide expressed in a retroviral expression vector is used to infect bone marrow cells that are subsequently transplanted into irradiated recipients. The pathology of the subject recipient is then examined after sufficient time has passed (e.g., six (6) weeks).

[0050] Expressing the polypeptide in host bacteria with an impaired capacity to proteolytically cleave the recombinant polypeptide is often used to maximize recombinant polypeptide expression (Gottesman, S., Gene Expression Technology: Methods in Enzymology, Academic Press, San Diego, California 185: 119-128 (1990)). Another strategy is to alter the nucleotide sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in E. coli (Wada et al., Nucleic Acids Res. 20: 2111-2118 (1992)). Such alteration of nucleotide sequences can be carried out by standard DNA synthesis techniques.

[0051] When used in mammalian cells, the expression vector's control functions are often provided by viral regulatory elements. For example, commonly used promoters are derived from polyoma, Adenovirus 2, cytomegalovirus and Simian Virus 40. Recombinant mammalian expression vectors are often capable of directing expression of the nucleic acid in a particular cell type (e.g., tissue-specific regulatory elements are used to express the nucleic acid). Non-limiting examples of suitable tissue-specific promoters include an albumin promoter (liver-specific; Pinkert et al., Genes Dev. 1: 268-277 (1987)), lymphoid-specific promoters (Calame & Eaton, Adv. Immunol. 43: 235-275 (1988)), promoters of T cell receptors (Winoto & Baltimore, EMBO J. 8: 729-733 (1989)) promoters of immunoglobulins (Banerji et al., Cell 33: 729-740 (1983); Queen & Baltimore, Cell 33: 741-748 (1983)), neuron-specific promoters (e.g., the neurofilament promoter; Byrne & Ruddle, Proc. Natl. Acad. Sci. USA 86: 5473-5477 (1989)), pancreas-specific promoters (Edlund et al., Science 230: 912-916 (1985)), and mammary gland-specific promoters (e.g., milk whey promoter; U.S. Patent No. 4,873,316 and European Application Publication No. 264,166). Developmentally-regulated promoters are sometimes utilized, for example, the murine hox promoters (Kessel & Gruss, Science 249: 374-379 (1990)) and the α-fetopolypeptide promoter (Campes & Tilghman, Genes Dev. 3: 537-546 (1989)).

[0052] A *IL1RL1* nucleic acid also may be cloned into an expression vector in an antisense orientation. Regulatory sequences (e.g., viral promoters and/or enhancers) operatively linked to a *IL1RL1* nucleic acid cloned in the antisense orientation can be chosen for directing constitutive, tissue specific or cell type specific expression of antisense RNA in a variety of cell types. Antisense expression vectors can be in the form of a recombinant plasmid, phagemid or attenuated virus. For a discussion of the regulation of gene expression using antisense genes *see*, *e.g.*, Weintraub *et al.*, Antisense RNA as a molecular tool for genetic analysis, *Reviews - Trends in Genetics*, Vol. 1(1) (1986).

[0053] Also provided herein are host cells that include a *IL1RL1* nucleotide sequence within a recombinant expression vector or a fragment of such a nucleotide sequence which facilitate homologous recombination into a specific site of the host cell genome. The terms "host cell" and "recombinant host cell" are used interchangeably herein. Such terms refer not only to the particular subject cell but rather also to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein. A host cell can be any prokaryotic or eukaryotic cell. For example, a target polypeptide can be expressed in bacterial cells such as *E. coli*, insect cells, yeast or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells). Other suitable host cells are known to those skilled in the art.

[0054] Vectors can be introduced into host cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection" are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, transduction/infection, DEAE-dextranmediated transfection, lipofection, or electroporation.

[0055] A host cell provided herein can be used to produce (*i.e.*, express) a target polypeptide or a substantially identical polypeptide thereof. Accordingly, further provided are methods for producing a target polypeptide using host cells described herein. In one embodiment, the method includes culturing host cells into which a recombinant expression vector encoding a target polypeptide has been introduced in a suitable medium such that a target polypeptide is produced. In another embodiment, the method further includes isolating a target polypeptide from the medium or the host cell.

[0056] Also provided are cells or purified preparations of cells which include a *IL1RL1* transgene, or which otherwise misexpress target polypeptide. Cell preparations can consist of human or non-human cells, *e.g.*, rodent cells, *e.g.*, mouse or rat cells, rabbit cells, or pig cells. In preferred embodiments, the cell or cells include a *IL1RL1* transgene (*e.g.*, a heterologous form of a *IL1RL1* gene, such as a human gene expressed in non-human cells). The transgene can be misexpressed, *e.g.*, overexpressed or underexpressed. In other preferred embodiments, the cell or cells include a gene which misexpress an endogenous target polypeptide (*e.g.*, expression of a gene is disrupted, also known as a knockout). Such cells can serve as a model for studying disorders which are related to mutated or mis-expressed alleles or for use in drug screening. Also provided are human cells (*e.g.*, a hematopoietic stem cells) transfected with a *IL1RL1* nucleic acid.

[0057] Also provided are cells or a purified preparation thereof (e.g., human cells) in which an endogenous *IL1RL1* nucleic acid is under the control of a regulatory sequence that does not normally control the expression of the endogenous gene. The expression characteristics of an endogenous gene within a cell (e.g., a cell line or microorganism) can be modified by inserting a heterologous DNA

regulatory element into the genome of the cell such that the inserted regulatory element is operably linked to the corresponding endogenous gene. For example, an endogenous corresponding gene (e.g., a gene which is "transcriptionally silent," not normally expressed, or expressed only at very low levels) may be activated by inserting a regulatory element which is capable of promoting the expression of a normally expressed gene product in that cell. Techniques such as targeted homologous recombinations, can be used to insert the heterologous DNA as described in, e.g., Chappel, US 5,272,071; WO 91/06667, published on May 16, 1991.

#### Transgenic Animals

[0058] Non-human transgenic animals that express a heterologous target polypeptide (e.g., expressed from a IL1RL1 nucleic acid or substantially identical sequence thereof) can be generated. Such animals are useful for studying the function and/or activity of a target polypeptide and for identifying and/or evaluating modulators of the activity of IL1RL1 nucleic acids and encoded polypeptides. As used herein, a "transgenic animal" is a non-human animal such as a mammal (e.g., a non-human primate such as chimpanzee, baboon, or macaque; an ungulate such as an equine, bovine, or caprine; or a rodent such as a rat, a mouse, or an Israeli sand rat), a bird (e.g., a chicken or a turkey), an amphibian (e.g., a frog, salamander, or newt), or an insect (e.g., Drosophila melanogaster), in which one or more of the cells of the animal includes a transgene. A transgene is exogenous DNA or a rearrangement (e.g., a deletion of endogenous chromosomal DNA) that is often integrated into or occurs in the genome of cells in a transgenic animal. A transgene can direct expression of an encoded gene product in one or more cell types or tissues of the transgenic animal, and other transgenes can reduce expression (e.g., a knockout). Thus, a transgenic animal can be one in which an endogenous nucleic acid homologous to a IL1RL1 nucleic acid has been altered by homologous recombination between the endogenous gene and an exogenous DNA molecule introduced into a cell of the animal (e.g., an embryonic cell of the animal) prior to development of the animal.

[0059] Intronic sequences and polyadenylation signals can also be included in the transgene to increase expression efficiency of the transgene. One or more tissue-specific regulatory sequences can be operably linked to a *IL1RL1* nucleotide sequence to direct expression of an encoded polypeptide to particular cells. A transgenic founder animal can be identified based upon the presence of a *IL1RL1* nucleotide sequence in its genome and/or expression of encoded mRNA in tissues or cells of the animals. A transgenic founder animal can then be used to breed additional animals carrying the transgene. Moreover, transgenic animals carrying a *IL1RL1* nucleotide sequence can further be bred to other transgenic animals carrying other transgenes.

[0060] Target polypeptides can be expressed in transgenic animals or plants by introducing, for example, a *IL1RL1* nucleic acid into the genome of an animal that encodes the target polypeptide. In

preferred embodiments the nucleic acid is placed under the control of a tissue specific promoter, e.g., a milk or egg specific promoter, and recovered from the milk or eggs produced by the animal. Also included is a population of cells from a transgenic animal.

#### Target Polypeptides

[0061] Also featured herein are isolated target polypeptides, which are encoded by a *IL1RL1* nucleotide sequence (e.g., SEQ ID NO: 1-4), or a substantially identical nucleotide sequence thereof. Examples of *IL1RL1* polypeptides are set forth in SEQ ID NO: 5-7. The term "polypeptide" as used herein includes proteins and peptides. An "isolated" or "purified" polypeptide or protein is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. In one embodiment, the language "substantially free" means preparation of a target polypeptide having less than about 30%, 20%, 10% and more preferably 5% (by dry weight), of non-target polypeptide (also referred to herein as a "contaminating protein"), or of chemical precursors or non-target chemicals. When the target polypeptide or a biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, specifically, where culture medium represents less than about 20%, sometimes less than about 10%, and often less than about 5% of the volume of the polypeptide preparation. Isolated or purified target polypeptide preparations are sometimes 0.01 milligrams or more or 0.1 milligrams or more, and often 1.0 milligrams or more and 10 milligrams or more in dry weight.

[0062] Further included herein are target polypeptide fragments. The polypeptide fragment may be a domain or part of a domain of a target polypeptide. The polypeptide fragment may have increased, decreased or unexpected biological activity. The polypeptide fragment is often 50 or fewer, 100 or fewer, or 200 or fewer amino acids in length, and is sometimes 300, 400, 500, 600, 700, or 900 or fewer amino acids in length.

[0063] Interleukin 1 receptor-like 1 isoform 1 (SEQ ID NO: 5) is a member of the interleukin 1 receptor family with no known ligand (orphan receptor). *IL1RL1* exists in soluble (SEQ ID NO: 6-7) and transmembrane forms, suggesting that it may have ligand or ligand scavenging activity. In an embodiment, *IL1RL1* protein agents may be administered to treat or prevent the occurrence of OA. *IL1RL1* protein agents include *IL1RL1* polypeptides or fragments thereof that have *IL1RL1* ligand activity (e.g., recombinant polypeptides of SEQ ID NO: 6-7). In a related embodiment, *IL1RL1* protein agents include *IL1RL1* polypeptides or fragments thereof that have *IL1RL1* ligand scavenging activity (e.g., recombinant polypeptide of SEQ ID NO: 5). Isolated *IL1RL1* polypeptides featured herein include the full-length polypeptide, the mature polypeptide (i.e., the polypeptide without the signal sequence

MGFWILAILTILMYSTAA) or a polypeptide fragment containing a domain or part of a *IL1RL1* domain. The polypeptide fragment may have increased, decreased or unexpected biological activity.

[0064] Substantially identical target polypeptides may depart from the amino acid sequences of target polypeptides in different manners. For example, conservative amino acid modifications may be introduced at one or more positions in the amino acid sequences of target polypeptides. A "conservative amino acid substitution" is one in which the amino acid is replaced by another amino acid having a similar structure and/or chemical function. Families of amino acid residues having similar structures and functions are well known. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Also, essential and non-essential amino acids may be replaced. A "non-essential" amino acid is one that can be altered without abolishing or substantially altering the biological function of a target polypeptide, whereas altering an "essential" amino acid abolishes or substantially alters the biological function of a target polypeptide. Amino acids that are conserved among target polypeptides are typically essential amino acids. In certain embodiments, the polypeptide includes one or more nonsynonymous polymorphic variants associated with osteoarthritis.

[0065] Also, target polypeptides may exist as chimeric or fusion polypeptides. As used herein, a target "chimeric polypeptide" or target "fusion polypeptide" includes a target polypeptide linked to a non-target polypeptide. A "non-target polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a polypeptide which is not substantially identical to the target polypeptide, which includes, for example, a polypeptide that is different from the target polypeptide and derived from the same or a different organism. The target polypeptide in the fusion polypeptide can correspond to an entire or nearly entire target polypeptide or a fragment thereof. The non-target polypeptide can be fused to the N-terminus or C-terminus of the target polypeptide.

[0066] Fusion polypeptides can include a moiety having high affinity for a ligand. For example, the fusion polypeptide can be a GST-target fusion polypeptide in which the target sequences are fused to the C-terminus of the GST sequences, or a polyhistidine-target fusion polypeptide in which the target polypeptide is fused at the N- or C-terminus to a string of histidine residues. Such fusion polypeptides can facilitate purification of recombinant target polypeptide. Expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide), and a nucleotide sequence in SEQ ID NO: 1-4, or a substantially identical nucleotide sequence thereof, can be cloned into an expression vector such that the fusion moiety is linked in-frame to the target polypeptide. Further, the fusion polypeptide can be a target polypeptide containing a heterologous signal sequence at its N-

terminus. In certain host cells (e.g., mammalian host cells), expression, secretion, cellular internalization, and cellular localization of a target polypeptide can be increased through use of a heterologous signal sequence. Fusion polypeptides can also include all or a part of a serum polypeptide (e.g., an IgG constant region or human serum albumin).

[0067] Target polypeptides can be incorporated into pharmaceutical compositions and administered to a subject *in vivo*. Administration of these target polypeptides can be used to affect the bioavailability of a substrate of the target polypeptide and may effectively increase target polypeptide biological activity in a cell. Target fusion polypeptides may be useful therapeutically for the treatment of disorders caused by, for example, (i) aberrant modification or mutation of a gene encoding a target polypeptide; (ii) misregulation of the gene encoding the target polypeptide; and (iii) aberrant post-translational modification of a target polypeptide. Also, target polypeptides can be used as immunogens to produce anti-target antibodies in a subject, to purify target polypeptide ligands or binding partners, and in screening assays to identify molecules which inhibit or enhance the interaction of a target polypeptide with a substrate.

[0068] In addition, polypeptides can be chemically synthesized using techniques known in the art (See, e.g., Creighton, 1983 Proteins. New York, N.Y.: W. H. Freeman and Company; and Hunkapiller et al., (1984) Nature July 12 -18;310(5973):105-11). For example, a relative short fragment can be synthesized by use of a peptide synthesizer. Furthermore, if desired, non-classical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the fragment sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, a-amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, g-Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoroamino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

[0069] Polypeptides and polypeptide fragments sometimes are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH4; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; and the like. Additional post-translational modifications include, for example, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell

expression. The polypeptide fragments may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the polypeptide.

[0070] Also provided are chemically modified derivatives of polypeptides that can provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see e.g., U.S. Pat. No: 4,179,337. The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

[0071] The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog).

[0072] The polymers should be attached to the polypeptide with consideration of effects on functional or antigenic domains of the polypeptide. There are a number of attachment methods available to those skilled in the art (e.g., EP 0 401 384 (coupling PEG to G-CSF) and Malik et al. (1992) Exp Hematol. September;20(8):1028-35 (pegylation of GM-CSF using tresyl chloride)). For example, polyethylene glycol may be covalently bound through amino acid residues via a reactive group, such as a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues, glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. For therapeutic purposes, the attachment sometimes is at an amino group, such as attachment at the N-terminus or lysine group.

[0073] Proteins can be chemically modified at the N-terminus. Using polyethylene glycol as an illustration of such a composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, and the like), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus may be accomplished by

reductive alkylation, which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

#### Substantially Identical Nucleic Acids and Polypeptides

[0074] Nucleotide sequences and polypeptide sequences that are substantially identical to a *IL1RL1* nucleotide sequence and the target polypeptide sequences encoded by those nucleotide sequences, respectively, are included herein. The term "substantially identical" as used herein refers to two or more nucleic acids or polypeptides sharing one or more identical nucleotide sequences or polypeptide sequences, respectively. Included are nucleotide sequences or polypeptide sequences that are 55% or more, 60% or more, 65% or more, 70% or more, 75% or more, 80% or more, 85% or more, 90% or more, 95% or more (each often within a 1%, 2%, 3% or 4% variability) identical to a *IL1RL1* nucleotide sequence or the encoded target polypeptide amino acid sequences. One test for determining whether two nucleic acids are substantially identical is to determine the percent of identical nucleotide sequences or polypeptide sequences shared between the nucleic acids or polypeptides.

[0075] Calculations of sequence identity are often performed as follows. Sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). The length of a reference sequence aligned for comparison purposes is sometimes 30% or more, 40% or more, 50% or more, often 60% or more, and more often 70% or more, 80% or more, 90% or more, or 100% of the length of the reference sequence. The nucleotides or amino acids at corresponding nucleotide or polypeptide positions, respectively, are then compared among the two sequences. When a position in the first sequence is occupied by the same nucleotide or amino acid as the corresponding position in the second sequence, the nucleotides or amino acids are deemed to be identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, introduced for optimal alignment of the two sequences.

[0076] Comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. Percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of Meyers & Miller, *CABIOS 4*: 11-17 (1989), which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. Also, percent identity between two amino acid sequences can be determined using the Needleman & Wunsch, *J. Mol. Biol. 48*: 444-453 (1970) algorithm which has been incorporated into the GAP program in the GCG software package (available at

the http address www.gcg.com), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. Percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software package (available at http address www.gcg.com), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A set of parameters often used is a Blossum 62 scoring matrix with a gap open penalty of 12, a gap extend penalty of 4, and a frameshift gap penalty of 5.

[0077] Another manner for determining if two nucleic acids are substantially identical is to assess whether a polynucleotide homologous to one nucleic acid will hybridize to the other nucleic acid under stringent conditions. As use herein, the term "stringent conditions" refers to conditions for hybridization and washing. Stringent conditions are known to those skilled in the art and can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y., 6.3.1-6.3.6 (1989). Aqueous and non-aqueous methods are described in that reference and either can be used. An example of stringent hybridization conditions is hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 50°C. Another example of stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 55°C. A further example of stringent hybridization conditions is hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 60°C. Often, stringent hybridization conditions are hybridization in 6X sodium chloride/sodium citrate (SSC) at about 45°C, followed by one or more washes in 0.2X SSC, 0.1% SDS at 65°C. More often, stringency conditions are 0.5M sodium phosphate, 7% SDS at 65°C, followed by one or more washes at 0.2X SSC, 1% SDS at 65°C.

[0078] An example of a substantially identical nucleotide sequence to a nucleotide sequence in SEQ ID NO: 1-4 is one that has a different nucleotide sequence but still encodes the same polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO: 1-4. Another example is a nucleotide sequence that encodes a polypeptide having a polypeptide sequence that is more than 70% or more identical to, sometimes more than 75% or more, 80% or more, or 85% or more identical to, and often more than 90% or more and 95% or more identical to a polypeptide sequence encoded by a nucleotide sequence in SEQ ID NO: 1-4.

[0079] Nucleotide sequences in SEQ ID NO: 1-4 and amino acid sequences of encoded polypeptides can be used as "query sequences" to perform a search against public databases to identify other family members or related sequences, for example. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul *et al.*, *J. Mol. Biol. 215*: 403-10 (1990). BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to nucleotide sequences in SEQ ID NO: 1-4. BLAST polypeptide

searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to polypeptides encoded by the nucleotide sequences of SEQ ID NO: 1-4. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, *Nucleic Acids Res. 25(17):* 3389-3402 (1997). When utilizing BLAST and Gapped BLAST programs, default parameters of the respective programs (*e.g.*, XBLAST and NBLAST) can be used (*see* the http address www.ncbi.nlm.nih.gov).

[0080] A nucleic acid that is substantially identical to a nucleotide sequence in SEQ ID NO: 1-4 may include polymorphic sites at positions equivalent to those described herein when the sequences are aligned. For example, using the alignment procedures described herein, SNPs in a sequence substantially identical to a sequence in SEQ ID NO: 1-4 can be identified at nucleotide positions that match (*i.e.*, align) with nucleotides at SNP positions in each nucleotide sequence in SEQ ID NO: 1-4. Also, where a polymorphic variation results in an insertion or deletion, insertion or deletion of a nucleotide sequence from a reference sequence can change the relative positions of other polymorphic sites in the nucleotide sequence.

[0081] Substantially identical nucleotide and polypeptide sequences include those that are naturally occurring, such as allelic variants (same locus), splice variants, homologs (different locus), and orthologs (different organism) or can be non-naturally occurring. Non-naturally occurring variants can be generated by mutagenesis techniques, including those applied to polynucleotides, cells, or organisms. The variants can contain nucleotide substitutions, deletions, inversions and insertions. Variation can occur in either or both the coding and non-coding regions. The variations can produce both conservative and non-conservative amino acid substitutions (as compared in the encoded product). Orthologs, homologs, allelic variants, and splice variants can be identified using methods known in the art. These variants normally comprise a nucleotide sequence encoding a polypeptide that is 50% or more, about 55% or more, often about 70-75% or more or about 80-85% or more, and sometimes about 90-95% or more identical to the amino acid sequences of target polypeptides or a fragment thereof. Such nucleic acid molecules can readily be identified as being able to hybridize under stringent conditions to a nucleotide sequence in SEQ ID NO: 1-4 or a fragment of this sequence. Nucleic acid molecules corresponding to orthologs, homologs, and allelic variants of a nucleotide sequence in SEQ ID NO: 1-4 can further be identified by mapping the sequence to the same chromosome or locus as the nucleotide sequence in SEQ ID NO: 1-4.

[0082] Also, substantially identical nucleotide sequences may include codons that are altered with respect to the naturally occurring sequence for enhancing expression of a target polypeptide in a particular expression system. For example, the nucleic acid can be one in which one or more codons are altered, and often 10% or more or 20% or more of the codons are altered for optimized expression in

bacteria (e.g., E. coli.), yeast (e.g., S. cervesiae), human (e.g., 293 cells), insect, or rodent (e.g., hamster) cells.

## Methods for Identifying Risk of Osteoarthritis

[0083] Methods for prognosing and diagnosing osteoarthritis are included herein. These methods include detecting the presence or absence of one or more polymorphic variations in a nucleotide sequence associated with osteoarthritis, such as variants in or around the loci set forth herein, or a substantially identical sequence thereof, in a sample from a subject, where the presence of a polymorphic variant described herein is indicative of a risk of osteoarthritis. Determining a risk of osteoarthritis sometimes refers to determining whether an individual is at an increased risk of osteoarthritis (e.g., intermediate risk or higher risk).

[0084] Thus, featured herein is a method for identifying a subject who is at risk of osteoarthritis, which comprises detecting an aberration associated with osteoarthritis in a nucleic acid sample from the subject. An embodiment is a method for detecting a risk of osteoarthritis in a subject, which comprises detecting the presence or absence of a polymorphic variation associated with osteoarthritis at a polymorphic site in a nucleotide sequence in a nucleic acid sample from a subject, where the nucleotide sequence comprises a polynucleotide sequence selected from the group consisting of: (a) a nucleotide sequence of SEQ ID NO: 1-4; (b) a nucleotide sequence which encodes a polypeptide consisting of an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4; (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4, or a nucleotide sequence about 90% or more identical to a nucleotide sequence of SEQ ID NO: 1-4; and (d) a fragment of a nucleotide sequence of (a), (b), or (c) comprising the polymorphic site; whereby the presence of the polymorphic variation is indicative of a predisposition to osteoarthritis in the subject. In certain embodiments, polymorphic variants at the positions described herein are detected for determining a risk of osteoarthritis, and polymorphic variants at positions in linkage disequilibrium with these positions are detected for determining a risk of osteoarthritis. As used herein, "SEQ ID NO: 1-4" refers to individual sequences in SEQ ID NO: 1, 2, 3 or 4, each sequence being separately applicable to embodiments described herein.

[0085] Risk of osteoarthritis sometimes is expressed as a probability, such as an odds ratio, percentage, or risk factor. Risk often is based upon the presence or absence of one or more polymorphic variants described herein, and also may be based in part upon phenotypic traits of the individual being tested. Methods for calculating risk based upon patient data are well known (see, e.g., Agresti, Categorical Data Analysis, 2nd Ed. 2002. Wiley). Allelotyping and genotyping analyses may be carried out in populations other than those exemplified herein to enhance the predictive power of the prognostic

method. These further analyses are executed in view of the exemplified procedures described herein, and may be based upon the same polymorphic variations or additional polymorphic variations.

[0086] In certain embodiments, determining the presence of a combination of two or more polymorphic variants associated with osteoarthritis in one or more genetic loci (e.g., one or more genes) of the sample is determined to identify, quantify and/or estimate, risk of osteoarthritis. The risk often is the probability of having or developing osteoarthritis. The risk sometimes is expressed as a relative risk with respect to a population average risk of osteoarthritis, and sometimes is expressed as a relative risk with respect to the lowest risk group. Such relative risk assessments often are based upon penetrance values determined by statistical methods, and are particularly useful to clinicians and insurance companies for assessing risk of osteoarthritis (e.g., a clinician can target appropriate detection, prevention and therapeutic regimens to a patient after determining the patient's risk of osteoarthritis, and an insurance company can fine tune actuarial tables based upon population genotype assessments of osteoarthritis risk). Risk of osteoarthritis sometimes is expressed as an odds ratio, which is the odds of a particular person having a genotype has or will develop osteoarthritis with respect to another genotype group (e.g., the most disease protective genotype or population average). In related embodiments, the determination is utilized to identify a subject at risk of osteoarthritis. In an embodiment, two or more polymorphic variations are detected in two or more regions in human genomic DNA associated with increased risk of osteoarthritis, such as a locus containing a IL1RL1, for example. In certain embodiments, 3 or more, or 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20 or more polymorphic variants are detected in the sample. In specific embodiments, polymorphic variants are detected in a IL1RL1 region, for example. In certain embodiments, polymorphic variants are detected at other genetic loci (e.g., the polymorphic variants can be detected in IL1RL1 in addition to other loci or only in other loci), where the other loci include but are not limited to those described in concurrentlyfiled patent applications having attorney docket number 524593008700, 524593008800, 524593008900, 524593009000 or 524593009200, which is incorporated herein by reference in its entirety.

[0087] Results from prognostic tests may be combined with other test results to diagnose osteoarthritis. For example, prognostic results may be gathered, a patient sample may be ordered based on a determined predisposition to osteoarthritis, the patient sample is analyzed, and the results of the analysis may be utilized to diagnose osteoarthritis. Also osteoarthritis diagnostic method can be developed from studies used to generate prognostic methods in which populations are stratified into subpopulations having different progressions of osteoarthritis. In another embodiment, prognostic results may be gathered, a patient's risk factors for developing osteoarthritis (e.g., age, weight, race, diet) analyzed, and a patient sample may be ordered based on a determined predisposition to osteoarthritis.

[0088] The nucleic acid sample typically is isolated from a biological sample obtained from a subject. For example, nucleic acid can be isolated from blood, saliva, sputum, urine, cell scrapings, and

biopsy tissue. The nucleic acid sample can be isolated from a biological sample using standard techniques, such as the technique described in Example 2. As used herein, the term "subject" refers primarily to humans but also refers to other mammals such as dogs, cats, and ungulates (e.g., cattle, sheep, and swine). Subjects also include avians (e.g., chickens and turkeys), reptiles, and fish (e.g., salmon), as embodiments described herein can be adapted to nucleic acid samples isolated from any of these organisms. The nucleic acid sample may be isolated from the subject and then directly utilized in a method for determining the presence of a polymorphic variant, or alternatively, the sample may be isolated and then stored (e.g., frozen) for a period of time before being subjected to analysis.

[0089] The presence or absence of a polymorphic variant is determined using one or both chromosomal complements represented in the nucleic acid sample. Determining the presence or absence of a polymorphic variant in both chromosomal complements represented in a nucleic acid sample from a subject having a copy of each chromosome is useful for determining the zygosity of an individual for the polymorphic variant (*i.e.*, whether the individual is homozygous or heterozygous for the polymorphic variant). Any oligonucleotide-based diagnostic may be utilized to determine whether a sample includes the presence or absence of a polymorphic variant in a sample. For example, primer extension methods, ligase sequence determination methods (*e.g.*, U.S. Pat. Nos. 5,679,524 and 5,952,174, and WO 01/27326), mismatch sequence determination methods (*e.g.*, U.S. Pat. Nos. 5,851,770; 5,958,692; 6,110,684; and 6,183,958), microarray sequence determination methods, restriction fragment length polymorphism (RFLP), single strand conformation polymorphism detection (SSCP) (*e.g.*, U.S. Pat. Nos. 5,891,625 and 6,013,499), PCR-based assays (*e.g.*, TAQMAN® PCR System (Applied Biosystems)), and nucleotide sequencing methods may be used.

[0090] Oligonucleotide extension methods typically involve providing a pair of oligonucleotide primers in a polymerase chain reaction (PCR) or in other nucleic acid amplification methods for the purpose of amplifying a region from the nucleic acid sample that comprises the polymorphic variation. One oligonucleotide primer is complementary to a region 3' of the polymorphism and the other is complementary to a region 5' of the polymorphism. A PCR primer pair may be used in methods disclosed in U.S. Pat. Nos. 4,683,195; 4,683,202, 4,965,188; 5,656,493; 5,998,143; 6,140,054; WO 01/27327; and WO 01/27329 for example. PCR primer pairs may also be used in any commercially available machines that perform PCR, such as any of the GENEAMP® Systems available from Applied Biosystems. Also, those of ordinary skill in the art will be able to design oligonucleotide primers based upon a *IL1RL1* nucleotide sequence using knowledge available in the art.

[0091] Also provided is an extension oligonucleotide that hybridizes to the amplified fragment adjacent to the polymorphic variation. As used herein, the term "adjacent" refers to the 3' end of the extension oligonucleotide being often 1 nucleotide from the 5' end of the polymorphic site, and sometimes 2, 3, 4, 5, 6, 7, 8, 9, or 10 nucleotides from the 5' end of the polymorphic site, in the nucleic

acid when the extension oligonucleotide is hybridized to the nucleic acid. The extension oligonucleotide then is extended by one or more nucleotides, and the number and/or type of nucleotides that are added to the extension oligonucleotide determine whether the polymorphic variant is present. Oligonucleotide extension methods are disclosed, for example, in U.S. Pat. Nos. 4,656,127; 4,851,331; 5,679,524; 5,834,189; 5,876,934; 5,908,755; 5,912,118; 5,976,802; 5,981,186; 6,004,744; 6,013,431; 6,017,702; 6,046,005; 6,087,095; 6,210,891; and WO 01/20039. Oligonucleotide extension methods using mass spectrometry are described, for example, in U.S. Pat. Nos. 5,547,835; 5,605,798; 5,691,141; 5,849,542; 5,869,242; 5,928,906; 6,043,031; and 6,194,144, and a method often utilized is described herein in Example 2.

[0092] A microarray can be utilized for determining whether a polymorphic variant is present or absent in a nucleic acid sample. A microarray may include any oligonucleotides described herein, and methods for making and using oligonucleotide microarrays suitable for diagnostic use are disclosed in U.S. Pat. Nos. 5,492,806; 5,525,464; 5,589,330; 5,695,940; 5,849,483; 6,018,041; 6,045,996; 6,136,541; 6,142,681; 6,156,501; 6,197,506; 6,223,127; 6,225,625; 6,229,911; 6,239,273; WO 00/52625; WO 01/25485; and WO 01/29259. The microarray typically comprises a solid support and the oligonucleotides may be linked to this solid support by covalent bonds or by non-covalent interactions. The oligonucleotides may also be linked to the solid support directly or by a spacer molecule. A microarray may comprise one or more oligonucleotides complementary to a polymorphic site set forth herein.

[0093] A kit also may be utilized for determining whether a polymorphic variant is present or absent in a nucleic acid sample. A kit often comprises one or more pairs of oligonucleotide primers useful for amplifying a fragment of a nucleotide sequence of SEQ ID NO: 1-4 or a substantially identical sequence thereof, where the fragment includes a polymorphic site. The kit sometimes comprises a polymerizing agent, for example, a thermostable nucleic acid polymerase such as one disclosed in U.S. Pat. Nos. 4,889,818 or 6,077,664. Also, the kit often comprises an elongation oligonucleotide that hybridizes to a IL1RL1 nucleotide sequence in a nucleic acid sample adjacent to the polymorphic site. Where the kit includes an elongation oligonucleotide, it also often comprises chain elongating nucleotides, such as dATP, dTTP, dGTP, dCTP, and dITP, including analogs of dATP, dTTP, dGTP, dCTP and dITP, provided that such analogs are substrates for a thermostable nucleic acid polymerase and can be incorporated into a nucleic acid chain elongated from the extension oligonucleotide. Along with chain elongating nucleotides would be one or more chain terminating nucleotides such as ddATP, ddTTP, ddGTP, ddCTP, and the like. In an embodiment, the kit comprises one or more oligonucleotide primer pairs, a polymerizing agent, chain elongating nucleotides, at least one elongation oligonucleotide, and one or more chain terminating nucleotides. Kits optionally include buffers, vials, microtiter plates, and instructions for use.

[0094] An individual identified as being at risk of osteoarthritis may be heterozygous or homozygous with respect to the allele associated with a higher risk of osteoarthritis. A subject homozygous for an allele associated with an increased risk of osteoarthritis is at a comparatively high risk of osteoarthritis, a subject heterozygous for an allele associated with an increased risk of osteoarthritis is at a comparatively intermediate risk of osteoarthritis, and a subject homozygous for an allele associated with a decreased risk of osteoarthritis is at a comparatively low risk of osteoarthritis. A genotype may be assessed for a complementary strand, such that the complementary nucleotide at a particular position is detected.

[0095] Also featured are methods for determining risk of osteoarthritis and/or identifying a subject at risk of osteoarthritis by contacting a polypeptide or protein encoded by a *IL1RL1* nucleotide sequence from a subject with an antibody that specifically binds to an epitope associated with increased risk of osteoarthritis in the polypeptide (e.g., an epitope comprising an alanine at position 78 in an *IR1LR1* polypeptide).

# Applications of Prognostic and Diagnostic Results to Pharmacogenomic Methods

[0096] Pharmacogenomics is a discipline that involves tailoring a treatment for a subject according to the subject's genotype as a particular treatment regimen may exert a differential effect depending upon the subject's genotype. For example, based upon the outcome of a prognostic test described herein, a clinician or physician may target pertinent information and preventative or therapeutic treatments to a subject who would be benefited by the information or treatment and avoid directing such information and treatments to a subject who would not be benefited (e.g., the treatment has no therapeutic effect and/or the subject experiences adverse side effects).

[0097] The following is an example of a pharmacogenomic embodiment. A particular treatment regimen can exert a differential effect depending upon the subject's genotype. Where a candidate therapeutic exhibits a significant interaction with a major allele and a comparatively weak interaction with a minor allele (e.g., an order of magnitude or greater difference in the interaction), such a therapeutic typically would not be administered to a subject genotyped as being homozygous for the minor allele, and sometimes not administered to a subject genotyped as being heterozygous for the minor allele. In another example, where a candidate therapeutic is not significantly toxic when administered to subjects who are homozygous for a major allele but is comparatively toxic when administered to subjects heterozygous or homozygous for a minor allele, the candidate therapeutic is not typically administered to subjects who are genotyped as being heterozygous or homozygous with respect to the minor allele.

[0098] The methods described herein are applicable to pharmacogenomic methods for preventing, alleviating or treating osteoarthritis. For example, a nucleic acid sample from an individual may be subjected to a prognostic test described herein. Where one or more polymorphic variations associated

with increased risk of osteoarthritis are identified in a subject, information for preventing or treating osteoarthritis and/or one or more osteoarthritis treatment regimens then may be prescribed to that subject.

[0099] In certain embodiments, a treatment or preventative regimen is specifically prescribed and/or administered to individuals who will most benefit from it based upon their risk of developing osteoarthritis assessed by the methods described herein. Thus, provided are methods for identifying a subject predisposed to osteoarthritis and then prescribing a therapeutic or preventative regimen to individuals identified as having a predisposition. Thus, certain embodiments are directed to a method for reducing osteoarthritis in a subject, which comprises: detecting the presence or absence of a polymorphic variant associated with osteoarthritis in a nucleotide sequence in a nucleic acid sample from a subject, where the nucleotide sequence comprises a polynucleotide sequence selected from the group consisting of: (a) a nucleotide sequence of SEQ ID NO: 1-4; (b) a nucleotide sequence which encodes a polypeptide consisting of an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4; (c) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4, or a nucleotide sequence about 90% or more identical to a nucleotide sequence of SEQ ID NO: 1-4; and (d) a fragment of a polynucleotide sequence of (a), (b), or (c); and prescribing or administering a treatment regimen to a subject from whom the sample originated where the presence of a polymorphic variation associated with osteoarthritis is detected in the nucleotide sequence. In these methods, predisposition results may be utilized in combination with other test results to diagnose osteoarthritis.

[0100] Certain preventative treatments often are prescribed to subjects having a predisposition to osteoarthritis and where the subject is diagnosed with osteoarthritis or is diagnosed as having symptoms indicative of an early stage of osteoarthritis. The treatment sometimes is preventative (e.g., is prescribed or administered to reduce the probability that osteoarthritis arises or progresses), sometimes is therapeutic, and sometimes delays, alleviates or halts the progression of osteoarthritis. Any known preventative or therapeutic treatment for alleviating or preventing the occurrence of osteoarthritis is prescribed and/or administered. For example, the treatment often is directed to decreasing pain and improving joint movement. Examples of OA treatments include exercises to keep joints flexible and improve muscle strength. Different medications to control pain, including corticosteroids and nonsteroidal anti-inflammatory drugs (NSAIDs, e.g., Voltaren); cyclooxygenase-2 (COX-2) inhibitors (e.g., Celebrex, Vioxx, Mobic, and Bextra); monoclonal antibodies (e.g., Remicade); tumor necrosis factor inhibitors (e.g., Enbrel); or injections of glucocorticoids, hyaluronic acid or chondrotin sulfate into joints that are inflamed and not responsive to NSAIDS. Orally administered chondroitin sulfate also may be used as a therapeutic, as it may increase hyaluronic acid levels and viscosity of synovial fluid, and decrease collagenase levels in synovial fluid. Also, glucosamine can serve as an OA therapeutic as delivering it into joints may inhibit enzymes involved in cartilage degradation and enhance the

production of hyaluronic acid. For mild pain without inflammation, acetaminophen may be used. Other treatments include: heat/cold therapy for temporary pain relief; joint protection to prevent strain or stress on painful joints; surgery to relieve chronic pain in damaged joints; and weight control to prevent extra stress on weight-bearing joints.

[0101] As therapeutic approaches for treating osteoarthritis continue to evolve and improve, the goal of treatments for osteoarthritis related disorders is to intervene even before clinical signs first manifest. Thus, genetic markers associated with susceptibility to osteoarthritis prove useful for early diagnosis, prevention and treatment of osteoarthritis.

[0102] As osteoarthritis preventative and treatment information can be specifically targeted to subjects in need thereof (e.g., those at risk of developing osteoarthritis or those in an early stage of osteoarthritis), provided herein is a method for preventing or reducing the risk of developing osteoarthritis in a subject, which comprises: (a) detecting the presence or absence of a polymorphic variation associated with osteoarthritis at a polymorphic site in a nucleotide sequence in a nucleic acid sample from a subject; (b) identifying a subject with a predisposition to osteoarthritis, whereby the presence of the polymorphic variation is indicative of a predisposition to osteoarthritis in the subject; and (c) if such a predisposition is identified, providing the subject with information about methods or products to prevent or reduce osteoarthritis or to delay the onset of osteoarthritis. Also provided is a method of targeting information or advertising to a subpopulation of a human population based on the subpopulation being genetically predisposed to a disease or condition, which comprises: (a) detecting the presence or absence of a polymorphic variation associated with osteoarthritis at a polymorphic site in a nucleotide sequence in a nucleic acid sample from a subject; (b) identifying the subpopulation of subjects in which the polymorphic variation is associated with osteoarthritis; and (c) providing information only to the subpopulation of subjects about a particular product which may be obtained and consumed or applied by the subject to help prevent or delay onset of the disease or condition.

[0103] Pharmacogenomics methods also may be used to analyze and predict a response to osteoarthritis treatment or a drug. For example, if pharmacogenomics analysis indicates a likelihood that an individual will respond positively to osteoarthritis treatment with a particular drug, the drug may be administered to the individual. Conversely, if the analysis indicates that an individual is likely to respond negatively to treatment with a particular drug, an alternative course of treatment may be prescribed. A negative response may be defined as either the absence of an efficacious response or the presence of toxic side effects. The response to a therapeutic treatment can be predicted in a background study in which subjects in any of the following populations are genotyped: a population that responds favorably to a treatment regimen, a population that does not respond significantly to a treatment regimen, and a population that responds adversely to a treatment regimen (e.g., exhibits one or more side effects). These populations are provided as examples and other populations and subpopulations may be analyzed. Based

upon the results of these analyses, a subject is genotyped to predict whether he or she will respond favorably to a treatment regimen, not respond significantly to a treatment regimen, or respond adversely to a treatment regimen.

[0104] The tests described herein also are applicable to clinical drug trials. One or more polymorphic variants indicative of response to an agent for treating osteoarthritis or to side effects to an agent for treating osteoarthritis may be identified using the methods described herein. Thereafter, potential participants in clinical trials of such an agent may be screened to identify those individuals most likely to respond favorably to the drug and exclude those likely to experience side effects. In that way, the effectiveness of drug treatment may be measured in individuals who respond positively to the drug, without lowering the measurement as a result of the inclusion of individuals who are unlikely to respond positively in the study and without risking undesirable safety problems.

[0105] Thus, another embodiment is a method of selecting an individual for inclusion in a clinical trial of a treatment or drug comprising the steps of: (a) obtaining a nucleic acid sample from an individual; (b) determining the identity of a polymorphic variation which is associated with a positive response to the treatment or the drug, or at least one polymorphic variation which is associated with a negative response to the treatment or the drug in the nucleic acid sample, and (c) including the individual in the clinical trial if the nucleic acid sample contains said polymorphic variation associated with a positive response to the treatment or the drug or if the nucleic acid sample lacks said polymorphic variation associated with a negative response to the treatment or the drug. In addition, the methods described herein for selecting an individual for inclusion in a clinical trial of a treatment or drug encompass methods with any further limitation described in this disclosure, or those following, specified alone or in any combination. The polymorphic variation may be in a sequence selected individually or in any combination from the group consisting of (i) a nucleotide sequence of SEQ ID NO: 1-4; (ii) a nucleotide sequence which encodes a polypeptide consisting of an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4; (iii) a nucleotide sequence which encodes a polypeptide that is 90% or more identical to an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4, or a nucleotide sequence about 90% or more identical to a nucleotide sequence of SEQ ID NO: 1-4; and (iv) a fragment of a polynucleotide sequence of (i), (ii), or (iii) comprising the polymorphic site. The including step (c) optionally comprises administering the drug or the treatment to the individual if the nucleic acid sample contains the polymorphic variation associated with a positive response to the treatment or the drug and the nucleic acid sample lacks said biallelic marker associated with a negative response to the treatment or the drug.

[0106] Also provided herein is a method of partnering between a diagnostic/prognostic testing provider and a provider of a consumable product, which comprises: (a) the diagnostic/prognostic testing provider detects the presence or absence of a polymorphic variation associated with osteoarthritis at a

polymorphic site in a nucleotide sequence in a nucleic acid sample from a subject; (b) the diagnostic/prognostic testing provider identifies the subpopulation of subjects in which the polymorphic variation is associated with osteoarthritis; (c) the diagnostic/prognostic testing provider forwards information to the subpopulation of subjects about a particular product which may be obtained and consumed or applied by the subject to help prevent or delay onset of the disease or condition; and (d) the provider of a consumable product forwards to the diagnostic test provider a fee every time the diagnostic/prognostic test provider forwards information to the subject as set forth in step (c) above.

## Compositions Comprising Osteoarthritis-Directed Molecules

[0107] Featured herein is a composition comprising a cell from a subject having osteoarthritis or at risk of osteoarthritis and one or more molecules specifically directed and targeted to a nucleic acid comprising a IL1RL1 nucleotide sequence or amino acid sequence. Such directed molecules include, but are not limited to, a compound that binds to a ILIRLI nucleotide sequence or amino acid sequence referenced herein; a RNAi or siRNA molecule having a strand complementary or substantially complementary to a IL1RL1 nucleotide sequence (e.g., hybridizes to a IL1RL1 nucleotide sequence under conditions of high stringency); an antisense nucleic acid complementary or substantially complementary to an RNA encoded by a IL1RL1 nucleotide sequence (e.g., hybridizes to a IL1RL1 nucleotide sequence under conditions of high stringency); a ribozyme that hybridizes to a IL1RL1 nucleotide sequence (e.g., hybridizes to a IL1RL1 nucleotide sequence under conditions of high stringency); a nucleic acid aptamer that specifically binds a polypeptide encoded by IL1RL1 nucleotide sequence; and an antibody that specifically binds to a polypeptide encoded by IL1RL1 nucleotide sequence or binds to a nucleic acid having such a nucleotide sequence. In specific embodiments, the osteoarthritis directed molecule interacts with a nucleic acid or polypeptide variant associated with osteoarthritis, such as variants referenced herein. In other embodiments, the osteoarthritis directed molecule interacts with a polypeptide involved in a signal pathway of a polypeptide encoded by a IL1RL1 nucleotide sequence, or a nucleic acid comprising such a nucleotide sequence.

[0108] Compositions sometimes include an adjuvant known to stimulate an immune response, and in certain embodiments, an adjuvant that stimulates a T-cell lymphocyte response. Adjuvants are known, including but not limited to an aluminum adjuvant (e.g., aluminum hydroxide); a cytokine adjuvant or adjuvant that stimulates a cytokine response (e.g., interleukin (IL)-12 and/or gamma-interferon cytokines); a Freund-type mineral oil adjuvant emulsion (e.g., Freund's complete or incomplete adjuvant); a synthetic lipoid compound; a copolymer adjuvant (e.g., TitreMax); a saponin; Quil A; a liposome; an oil-in-water emulsion (e.g., an emulsion stabilized by Tween 80 and pluronic polyoxyethlene/polyoxypropylene block copolymer (Syntex Adjuvant Formulation); TitreMax; detoxified endotoxin (MPL) and mycobacterial cell wall components (TDW, CWS) in 2% squalene (Ribi

Adjuvant System)); a muramyl dipeptide; an immune-stimulating complex (ISCOM, e.g., an Agmodified saponin/cholesterol micelle that forms stable cage-like structure); an aqueous phase adjuvant that does not have a depot effect (e.g., Gerbu adjuvant); a carbohydrate polymer (e.g., AdjuPrime); L-tyrosine; a manide-oleate compound (e.g., Montanide); an ethylene-vinyl acetate copolymer (e.g., Elvax 40W1,2); or lipid A, for example. Such compositions are useful for generating an immune response against osteoarthritis directed molecule (e.g., an HLA-binding subsequence within a polypeptide encoded by a *IL1RL1* nucleotide sequence). In such methods, a peptide having an amino acid subsequence of a polypeptide encoded by a *IL1RL1* nucleotide sequence is delivered to a subject, where the subsequence binds to an HLA molecule and induces a CTL lymphocyte response. The peptide sometimes is delivered to the subject as an isolated peptide or as a minigene in a plasmid that encodes the peptide. Methods for identifying HLA-binding subsequences in such polypeptides are known (see e.g., publication WO02/20616 and PCT application US98/01373 for methods of identifying such sequences).

[0109] The cell may be in a group of cells cultured *in vitro* or in a tissue maintained *in vitro* or present in an animal *in vivo* (e.g., a rat, mouse, ape or human). In certain embodiments, a composition comprises a component from a cell such as a nucleic acid molecule (e.g., genomic DNA), a protein mixture or isolated protein, for example. The aforementioned compositions have utility in diagnostic, prognostic and pharmacogenomic methods described previously and in therapeutics described hereafter. Certain osteoarthritis directed molecules are described in greater detail below.

#### Compounds

[0110] Compounds can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: biological libraries; peptoid libraries (libraries of molecules having the functionalities of peptides, but with a novel, non-peptide backbone which are resistant to enzymatic degradation but which nevertheless remain bioactive (see, e.g., Zuckermann et al., J. Med. Chem.37: 2678-85 (1994)); spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; "one-bead one-compound" library methods; and synthetic library methods using affinity chromatography selection. Biological library and peptoid library approaches are typically limited to peptide libraries, while the other approaches are applicable to peptide, non-peptide oligomer or small molecule libraries of compounds (Lam, Anticancer Drug Des. 12: 145, (1997)). Examples of methods for synthesizing molecular libraries are described, for example, in DeWitt et al., Proc. Natl. Acad. Sci. U.S.A. 90: 6909 (1993); Erb et al., Proc. Natl. Acad. Sci. USA 91: 11422 (1994); Zuckermann et al., J. Med. Chem. 37: 2678 (1994); Cho et al., Science 261: 1303 (1993); Carrell et al., Angew. Chem. Int. Ed. Engl. 33: 2061 (1994); and in Gallop et al., J. Med. Chem. 37: 1233 (1994).

[0111] Libraries of compounds may be presented in solution (e.g., Houghten, Biotechniques 13: 412-421 (1992)), or on beads (Lam, Nature 354: 82-84 (1991)), chips (Fodor, Nature 364: 555-556 (1993)), bacteria or spores (Ladner, United States Patent No. 5,223,409), plasmids (Cull et al., Proc. Natl. Acad. Sci. USA 89: 1865-1869 (1992)) or on phage (Scott and Smith, Science 249: 386-390 (1990); Devlin, Science 249: 404-406 (1990); Cwirla et al., Proc. Natl. Acad. Sci. 87: 6378-6382 (1990); Felici, J. Mol. Biol. 222: 301-310 (1991); Ladner supra.).

[0112] A compound sometimes alters expression and sometimes alters activity of a polypeptide target and may be a small molecule. Small molecules include, but are not limited to, peptides, peptidomimetics (e.g., peptoids), amino acids, amino acid analogs, polynucleotides, polynucleotide analogs, nucleotides, nucleotide analogs, organic or inorganic compounds (i.e., including heteroorganic and organometallic compounds) having a molecular weight less than about 10,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 5,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 1,000 grams per mole, organic or inorganic compounds having a molecular weight less than about 500 grams per mole, and salts, esters, and other pharmaceutically acceptable forms of such compounds.

# Antisense Nucleic Acid Molecules, Ribozymes, RNAi, siRNA and Modified Nucleic Acid Molecules

[0113] An "antisense" nucleic acid refers to a nucleotide sequence complementary to a "sense" nucleic acid encoding a polypeptide, e.g., complementary to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. The antisense nucleic acid can be complementary to an entire coding strand, or to a portion thereof or a substantially identical sequence thereof. In another embodiment, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence (e.g., 5' and 3' untranslated regions in SEQ ID NO: 1).

[0114] An antisense nucleic acid can be designed such that it is complementary to the entire coding region of an mRNA encoded by a nucleotide sequence (e.g., SEQ ID NO: 1), and often the antisense nucleic acid is an oligonucleotide antisense to only a portion of a coding or noncoding region of the mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of the mRNA, e.g., between the -10 and +10 regions of the target gene nucleotide sequence of interest. An antisense oligonucleotide can be, for example, about 7, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, or more nucleotides in length. The antisense nucleic acids, which include the ribozymes described hereafter, can be designed to target a *IL1RL1* nucleotide sequence, often a variant associated with osteoarthritis, or a substantially identical sequence thereof. Among the variants, minor alleles and major alleles can be targeted, and those associated with a higher risk of osteoarthritis are often designed, tested, and administered to subjects.

[0115] An antisense nucleic acid can be constructed using chemical synthesis and enzymatic ligation reactions using standard procedures. For example, an antisense nucleic acid (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Antisense nucleic acid also can be produced biologically using an expression vector into which a nucleic acid has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted nucleic acid will be of an antisense orientation to a target nucleic acid of interest, described further in the following subsection).

[0116] When utilized as therapeutics, antisense nucleic acids typically are administered to a subject (e.g., by direct injection at a tissue site) or generated in situ such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding a polypeptide and thereby inhibit expression of the polypeptide, for example, by inhibiting transcription and/or translation. Alternatively, antisense nucleic acid molecules can be modified to target selected cells and then are administered systemically. For systemic administration, antisense molecules can be modified such that they specifically bind to receptors or antigens expressed on a selected cell surface, for example, by linking antisense nucleic acid molecules to peptides or antibodies which bind to cell surface receptors or antigens. Antisense nucleic acid molecules can also be delivered to cells using the vectors described herein. Sufficient intracellular concentrations of antisense molecules are achieved by incorporating a strong promoter, such as a pol II or pol III promoter, in the vector construct.

[0117] Antisense nucleic acid molecules sometimes are alpha-anomeric nucleic acid molecules. An alpha-anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual beta-units, the strands run parallel to each other (Gaultier et al., Nucleic Acids. Res. 15: 6625-6641 (1987)). Antisense nucleic acid molecules can also comprise a 2'-o-methylribonucleotide (Inoue et al., Nucleic Acids Res. 15: 6131-6148 (1987)) or a chimeric RNA-DNA analogue (Inoue et al., FEBS Lett. 215: 327-330 (1987)). Antisense nucleic acids sometimes are composed of DNA or PNA or any other nucleic acid derivatives described previously.

[0118] In another embodiment, an antisense nucleic acid is a ribozyme. A ribozyme having specificity for a *IL1RL1* nucleotide sequence can include one or more sequences complementary to such a nucleotide sequence, and a sequence having a known catalytic region responsible for mRNA cleavage (see e.g., U.S. Pat. No. 5,093,246 or Haselhoff and Gerlach, Nature 334: 585-591 (1988)). For example, a derivative of a Tetrahymena L-19 IVS RNA is sometimes utilized in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in a mRNA (see e.g., Cech et al. U.S. Patent No. 4,987,071; and Cech et al. U.S. Patent No. 5,116,742). Also, target mRNA sequences

can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules (see e.g., Bartel & Szostak, Science 261: 1411-1418 (1993)).

[0119] Osteoarthritis directed molecules include in certain embodiments nucleic acids that can form triple helix structures with a *IL1RL1* nucleotide sequence, or a substantially identical sequence thereof, especially one that includes a regulatory region that controls expression of a polypeptide. Gene expression can be inhibited by targeting nucleotide sequences complementary to the regulatory region of a nucleotide sequence referenced herein or a substantially identical sequence (e.g., promoter and/or enhancers) to form triple helical structures that prevent transcription of a gene in target cells (see e.g., Helene, Anticancer Drug Des. 6(6): 569-84 (1991); Helene et al., Ann. N.Y. Acad. Sci. 660: 27-36 (1992); and Maher, Bioassays 14(12): 807-15 (1992). Potential sequences that can be targeted for triple helix formation can be increased by creating a so-called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizeable stretch of either purines or pyrimidines to be present on one strand of a duplex.

[0120] Osteoarthritis directed molecules include RNAi and siRNA nucleic acids. Gene expression may be inhibited by the introduction of double-stranded RNA (dsRNA), which induces potent and specific gene silencing, a phenomenon called RNA interference or RNAi. See, e.g., Fire et al., US Patent Number 6,506,559; Tuschl et al. PCT International Publication No. WO 01/75164; Kay et al. PCT International Publication No. WO 03/010180A1; or Bosher JM, Labouesse, Nat Cell Biol 2000 Feb;2(2):E31-6. This process has been improved by decreasing the size of the double-stranded RNA to 20-24 base pairs (to create small-interfering RNAs or siRNAs) that "switched off" genes in mammalian cells without initiating an acute phase response, i.e., a host defense mechanism that often results in cell death (see, e.g., Caplen et al. Proc Natl Acad Sci U S A. 2001 Aug 14;98(17):9742-7 and Elbashir et al. Methods 2002 Feb;26(2):199-213). There is increasing evidence of post-transcriptional gene silencing by RNA interference (RNAi) for inhibiting targeted expression in mammalian cells at the mRNA level, in human cells. There is additional evidence of effective methods for inhibiting the proliferation and migration of tumor cells in human patients, and for inhibiting metastatic cancer development (see, e.g., U.S. Patent Application No. US2001000993183; Caplen et al. Proc Natl Acad Sci U S A; and Abderrahmani et al. Mol Cell Biol 2001 Nov21(21):7256-67).

[0121] An "siRNA" or "RNAi" refers to a nucleic acid that forms a double stranded RNA and has the ability to reduce or inhibit expression of a gene or target gene when the siRNA is delivered to or expressed in the same cell as the gene or target gene. "siRNA" refers to short double-stranded RNA formed by the complementary strands. Complementary portions of the siRNA that hybridize to form the double stranded molecule often have substantial or complete identity to the target molecule sequence. In

one embodiment, an siRNA refers to a nucleic acid that has substantial or complete identity to a target gene and forms a double stranded siRNA.

[0122] When designing the siRNA molecules, the targeted region often is selected from a given DNA sequence beginning 50 to 100 nucleotides downstream of the start codon. See, e.g., Elbashir et al,. Methods 26:199-213 (2002). Initially, 5' or 3' UTRs and regions nearby the start codon were avoided assuming that UTR-binding proteins and/or translation initiation complexes may interfere with binding of the siRNP or RISC endonuclease complex. Sometimes regions of the target 23 nucleotides in length conforming to the sequence motif AA(N19)TT (N, an nucleotide), and regions with approximately 30% to 70% G/C-content (often about 50% G/C-content) often are selected. If no suitable sequences are found, the search often is extended using the motif NA(N21). The sequence of the sense siRNA sometimes corresponds to (N19) TT or N21 (position 3 to 23 of the 23-nt motif), respectively. In the latter case, the 3' end of the sense siRNA often is converted to TT. The rationale for this sequence conversion is to generate a symmetric duplex with respect to the sequence composition of the sense and antisense 3' overhangs. The antisense siRNA is synthesized as the complement to position 1 to 21 of the 23-nt motif. Because position 1 of the 23-nt motif is not recognized sequence-specifically by the antisense siRNA, the 3'-most nucleotide residue of the antisense siRNA can be chosen deliberately. However, the penultimate nucleotide of the antisense siRNA (complementary to position 2 of the 23-nt motif) often is complementary to the targeted sequence. For simplifying chemical synthesis, TT often is utilized. siRNAs corresponding to the target motif NAR(N17)YNN, where R is purine (A,G) and Y is pyrimidine (C,U), often are selected. Respective 21 nucleotide sense and antisense siRNAs often begin with a purine nucleotide and can also be expressed from pol III expression vectors without a change in targeting site. Expression of RNAs from pol III promoters often is efficient when the first transcribed nucleotide is a purine.

[0123] The sequence of the siRNA can correspond to the full length target gene, or a subsequence thereof. Often, the siRNA is about 15 to about 50 nucleotides in length (e.g., each complementary sequence of the double stranded siRNA is 15-50 nucleotides in length, and the double stranded siRNA is about 15-50 base pairs in length, sometimes about 20-30 nucleotides in length or about 20-25 nucleotides in length, e.g., 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length. The siRNA sometimes is about 21 nucleotides in length. Methods of using siRNA are well known in the art, and specific siRNA molecules may be purchased from a number of companies including Dharmacon Research, Inc.

[0124] Antisense, ribozyme, RNAi and siRNA nucleic acids can be altered to form modified nucleic acid molecules. The nucleic acids can be altered at base moieties, sugar moieties or phosphate backbone moieties to improve stability, hybridization, or solubility of the molecule. For example, the deoxyribose phosphate backbone of nucleic acid molecules can be modified to generate peptide nucleic acids (see Hyrup et al., Bioorganic & Medicinal Chemistry 4 (1): 5-23 (1996)). As used herein, the terms "peptide

nucleic acid" or "PNA" refers to a nucleic acid mimic such as a DNA mimic, in which the deoxyribose phosphate backbone is replaced by a pseudopeptide backbone and only the four natural nucleobases are retained. The neutral backbone of a PNA can allow for specific hybridization to DNA and RNA under conditions of low ionic strength. Synthesis of PNA oligomers can be performed using standard solid phase peptide synthesis protocols as described, for example, in Hyrup et al., (1996) supra and Perry-O'Keefe et al., Proc. Natl. Acad. Sci. 93: 14670-675 (1996).

[0125] PNA nucleic acids can be used in prognostic, diagnostic, and therapeutic applications. For example, PNAs can be used as antisense or antigene agents for sequence-specific modulation of gene expression by, for example, inducing transcription or translation arrest or inhibiting replication. PNA nucleic acid molecules can also be used in the analysis of single base pair mutations in a gene, (e.g., by PNA-directed PCR clamping); as "artificial restriction enzymes" when used in combination with other enzymes, (e.g., S1 nucleases (Hyrup (1996) supra)); or as probes or primers for DNA sequencing or hybridization (Hyrup et al., (1996) supra; Perry-O'Keefe supra).

[0126] In other embodiments, oligonucleotides may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across cell membranes (see e.g., Letsinger et al., Proc. Natl. Acad. Sci. USA 86: 6553-6556 (1989); Lemaitre et al., Proc. Natl. Acad. Sci. USA 84: 648-652 (1987); PCT Publication No. W088/09810) or the blood-brain barrier (see, e.g., PCT Publication No. W089/10134). In addition, oligonucleotides can be modified with hybridization-triggered cleavage agents (See, e.g., Krol et al., Bio-Techniques 6: 958-976 (1988)) or intercalating agents. (See, e.g., Zon, Pharm. Res. 5: 539-549 (1988)). To this end, the oligonucleotide may be conjugated to another molecule, (e.g., a peptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent).

[0127] Also included herein are molecular beacon oligonucleotide primer and probe molecules having one or more regions complementary to a *IL1RL1* nucleotide sequence, or a substantially identical sequence thereof, two complementary regions one having a fluorophore and one a quencher such that the molecular beacon is useful for quantifying the presence of the nucleic acid in a sample. Molecular beacon nucleic acids are described, for example, in Lizardi et al., U.S. Patent No. 5,854,033; Nazarenko et al., U.S. Patent No. 5,866,336, and Livak et al., U.S. Patent 5,876,930.

# **Antibodies**

[0128] The term "antibody" as used herein refers to an immunoglobulin molecule or immunologically active portion thereof, i.e., an antigen-binding portion. Examples of immunologically active portions of immunoglobulin molecules include F(ab) and F(ab')<sub>2</sub> fragments which can be generated by treating the antibody with an enzyme such as pepsin. An antibody sometimes is a polyclonal, monoclonal, recombinant (e.g., a chimeric or humanized), fully human, non-human (e.g.,

murine), or a single chain antibody. An antibody may have effector function and can fix complement, and is sometimes coupled to a toxin or imaging agent.

[0129] A full-length polypeptide or antigenic peptide fragment encoded by a nucleotide sequence referenced herein can be used as an immunogen or can be used to identify antibodies made with other immunogens, e.g., cells, membrane preparations, and the like. An antigenic peptide often includes at least 8 amino acid residues of the amino acid sequences encoded by a nucleotide sequence referenced herein, or substantially identical sequence thereof, and encompasses an epitope. Antigenic peptides sometimes include 10 or more amino acids, 15 or more amino acids, 20 or more amino acids, or 30 or more amino acids. Hydrophilic and hydrophobic fragments of polypeptides sometimes are used as immunogens.

[0130] Epitopes encompassed by the antigenic peptide are regions located on the surface of the polypeptide (e.g., hydrophilic regions) as well as regions with high antigenicity. For example, an Emini surface probability analysis of the human polypeptide sequence can be used to indicate the regions that have a particularly high probability of being localized to the surface of the polypeptide and are thus likely to constitute surface residues useful for targeting antibody production. The antibody may bind an epitope on any domain or region on polypeptides described herein.

[0131] Also, chimeric, humanized, and completely human antibodies are useful for applications which include repeated administration to subjects. Chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, can be made using standard recombinant DNA techniques. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson et al International Application No. PCT/US86/02269; Akira, et al European Patent Application 184,187; Taniguchi, M., European Patent Application 171,496; Morrison et al European Patent Application 173,494; Neuberger et al PCT International Publication No. WO 86/01533; Cabilly et al U.S. Patent No. 4,816,567; Cabilly et al European Patent Application 125,023; Better et al., Science 240: 1041-1043 (1988); Liu et al., Proc. Natl. Acad. Sci. USA 84: 3439-3443 (1987); Liu et al., J. Immunol. 139: 3521-3526 (1987); Sun et al., Proc. Natl. Acad. Sci. USA 84: 214-218 (1987); Nishimura et al., Canc. Res. 47: 999-1005 (1987); Wood et al., Nature 314: 446-449 (1985); and Shaw et al., J. Natl. Cancer Inst. 80: 1553-1559 (1988); Morrison, S. L., Science 229: 1202-1207 (1985); Oi et al., BioTechniques 4: 214 (1986); Winter U.S. Patent 5,225,539; Jones et al., Nature 321: 552-525 (1986); Verhoeyan et al., Science 239: 1534; and Beidler et al., J. Immunol. 141: 4053-4060 (1988).

[0132] Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Such antibodies can be produced using transgenic mice that are incapable of expressing endogenous immunoglobulin heavy and light chains genes, but which can express human heavy and light chain genes. See, for example, Lonberg and Huszar, Int. Rev. Immunol. 13: 65-93 (1995); and U.S.

Patent Nos. 5,625,126; 5,633,425; 5,569,825; 5,661,016; and 5,545,806. In addition, companies such as Abgenix, Inc. (Fremont, CA) and Medarex, Inc. (Princeton, NJ), can be engaged to provide human antibodies directed against a selected antigen using technology similar to that described above. Completely human antibodies that recognize a selected epitope also can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody (e.g., a murine antibody) is used to guide the selection of a completely human antibody recognizing the same epitope. This technology is described for example by Jespers et al., Bio/Technology 12: 899-903 (1994).

[0133] An antibody can be a single chain antibody. A single chain antibody (scFV) can be engineered (see, e.g., Colcher et al., Ann. N Y Acad. Sci. 880: 263-80 (1999); and Reiter, Clin. Cancer Res. 2: 245-52 (1996)). Single chain antibodies can be dimerized or multimerized to generate multivalent antibodies having specificities for different epitopes of the same target polypeptide.

[0134] Antibodies also may be selected or modified so that they exhibit reduced or no ability to bind an Fc receptor. For example, an antibody may be an isotype or subtype, fragment or other mutant, which does not support binding to an Fc receptor (e.g., it has a mutagenized or deleted Fc receptor binding region).

[0135] Also, an antibody (or fragment thereof) may be conjugated to a therapeutic moiety such as a cytotoxin, a therapeutic agent or a radioactive metal ion. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1 dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thiotepa chlorambucil, melphalan, carmustine (BCNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

[0136] Antibody conjugates can be used for modifying a given biological response. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a polypeptide such as tumor necrosis factor, gamma-interferon, alpha-interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or

other growth factors. Also, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, for example.

[0137] An antibody (e.g., monoclonal antibody) can be used to isolate target polypeptides by standard techniques, such as affinity chromatography or immunoprecipitation. Moreover, an antibody can be used to detect a target polypeptide (e.g., in a cellular lysate or cell supernatant) in order to evaluate the abundance and pattern of expression of the polypeptide. Antibodies can be used diagnostically to monitor polypeptide levels in tissue as part of a clinical testing procedure, e.g., to determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance (i.e., antibody labeling). Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase,  $\beta$ -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S or <sup>3</sup>H. Also, an antibody can be utilized as a test molecule for determining whether it can treat osteoarthritis, and as a therapeutic for administration to a subject for treating osteoarthritis.

[0138] An antibody can be made by immunizing with a purified antigen, or a fragment thereof, e.g., a fragment described herein, a membrane associated antigen, tissues, e.g., crude tissue preparations, whole cells, preferably living cells, lysed cells, or cell fractions.

[0139] Included herein are antibodies which bind only a native polypeptide, only denatured or otherwise non-native polypeptide, or which bind both, as well as those having linear or conformational epitopes. Conformational epitopes sometimes can be identified by selecting antibodies that bind to native but not denatured polypeptide. Also featured are antibodies that specifically bind to a polypeptide variant associated with osteoarthritis.

# Methods for Identifying Candidate Therapeutics for Treating Osteoarthritis

[0140] Current therapies for the treatment of osteoarthritis have limited efficacy, limited tolerability and significant mechanism-based side effects, and few of the available therapies adequately address underlying defects. Current therapeutic approaches were largely developed in the absence of defined molecular targets or even a solid understanding of disease pathogenesis. Therefore, provided are methods of identifying candidate therapeutics that target biochemical pathways related to the development of osteoarthritis.

- [0141] Thus, featured herein are methods for identifying a candidate therapeutic for treating osteoarthritis. The methods comprise contacting a test molecule with a target molecule in a system. A "target molecule" as used herein refers to a *IL1RL1* nucleic acid, a substantially identical nucleic acid thereof, or a fragment thereof, and an encoded polypeptide of the foregoing. The methods also comprise determining the presence or absence of an interaction between the test molecule and the target molecule, where the presence of an interaction between the test molecule and the nucleic acid or polypeptide identifies the test molecule as a candidate osteoarthritis therapeutic. The interaction between the test molecule and the target molecule may be quantified.
- [0142] Test molecules and candidate therapeutics include, but are not limited to, compounds, antisense nucleic acids, siRNA molecules, ribozymes, polypeptides or proteins encoded by a IL1RL1 nucleotide sequence, or a substantially identical sequence or fragment thereof, and immunotherapeutics (e.g., antibodies and HLA-presented polypeptide fragments). A test molecule or candidate therapeutic may act as a modulator of target molecule concentration or target molecule function in a system. A "modulator" may agonize (i.e., up-regulates) or antagonize (i.e., down-regulates) a target molecule concentration partially or completely in a system by affecting such cellular functions as DNA replication and/or DNA processing (e.g., DNA methylation or DNA repair), RNA transcription and/or RNA processing (e.g., removal of intronic sequences and/or translocation of spliced mRNA from the nucleus), polypeptide production (e.g., translation of the polypeptide from mRNA), and/or polypeptide posttranslational modification (e.g., glycosylation, phosphorylation, and proteolysis of pro-polypeptides). A modulator may also agonize or antagonize a biological function of a target molecule partially or completely, where the function may include adopting a certain structural conformation, interacting with one or more binding partners, ligand binding, catalysis (e.g., phosphorylation, dephosphorylation, hydrolysis, methylation, and isomerization), and an effect upon a cellular event (e.g., effecting progression of osteoarthritis).
- [0143] As used herein, the term "system" refers to a cell free *in vitro* environment and a cell-based environment such as a collection of cells, a tissue, an organ, or an organism. A system is "contacted" with a test molecule in a variety of manners, including adding molecules in solution and allowing them to interact with one another by diffusion, cell injection, and any administration routes in an animal. As used herein, the term "interaction" refers to an effect of a test molecule on test molecule, where the effect sometimes is binding between the test molecule and the target molecule, and sometimes is an observable change in cells, tissue, or organism.
- [0144] There are many standard methods for detecting the presence or absence of interaction between a test molecule and a target molecule. For example, titrametric, acidimetric, radiometric, NMR, monolayer, polarographic, spectrophotometric, fluorescent, and ESR assays probative of a target molecule interaction may be utilized.

[0145] Test molecule/target molecule interactions can be detected and/or quantified using assays known in the art. For example, an interaction can be determined by labeling the test molecule and/or the target molecule, where the label is covalently or non-covalently attached to the test molecule or target molecule. The label is sometimes a radioactive molecule such as <sup>125</sup>I, <sup>131</sup>I, <sup>35</sup>S or <sup>3</sup>H, which can be detected by direct counting of radioemission or by scintillation counting. Also, enzymatic labels such as horseradish peroxidase, alkaline phosphatase, or luciferase may be utilized where the enzymatic label can be detected by determining conversion of an appropriate substrate to product. In addition, presence or absence of an interaction can be determined without labeling. For example, a microphysiometer (e.g., Cytosensor) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indication of an interaction between a test molecule and target molecule (McConnell, H. M. et al., Science 257: 1906-1912 (1992)).

[0146] In cell-based systems, cells typically include a *IL1RL1* nucleic acid, an encoded polypeptide, or substantially identical nucleic acid or polypeptide thereof, and are often of mammalian origin, although the cell can be of any origin. Whole cells, cell homogenates, and cell fractions (*e.g.*, cell membrane fractions) can be subjected to analysis. Where interactions between a test molecule with a target polypeptide are monitored, soluble and/or membrane bound forms of the polypeptide may be utilized. Where membrane-bound forms of the polypeptide are used, it may be desirable to utilize a solubilizing agent. Examples of such solubilizing agents include non-ionic detergents such as noctylglucoside, n-dodecylglucoside, n-dodecylmaltoside, octanoyl-N-methylglucamide, decanoyl-N-methylglucamide, Triton® X-100, Triton® X-114, Thesit®, Isotridecypoly(ethylene glycol ether)<sub>n</sub>, 3-[(3-cholamidopropyl)dimethylamminio]-1-propane sulfonate (CHAPSO), or N-dodecyl-N,N-dimethyl-3-ammonio-1-propane sulfonate.

[0147] An interaction between a test molecule and target molecule also can be detected by monitoring fluorescence energy transfer (FET) (see, e.g., Lakowicz et al., U.S. Patent No. 5,631,169; Stavrianopoulos et al. U.S. Patent No. 4,868,103). A fluorophore label on a first, "donor" molecule is selected such that its emitted fluorescent energy will be absorbed by a fluorescent label on a second, "acceptor" molecule, which in turn is able to fluoresce due to the absorbed energy. Alternately, the "donor" polypeptide molecule may simply utilize the natural fluorescent energy of tryptophan residues. Labels are chosen that emit different wavelengths of light, such that the "acceptor" molecule label may be differentiated from that of the "donor". Since the efficiency of energy transfer between the labels is related to the distance separating the molecules, the spatial relationship between the molecules can be assessed. In a situation in which binding occurs between the molecules, the fluorescent emission of the

"acceptor" molecule label in the assay should be maximal. An FET binding event can be conveniently measured through standard fluorometric detection means well known in the art (e.g., using a fluorimeter).

[0148] In another embodiment, determining the presence or absence of an interaction between a test molecule and a target molecule can be effected by monitoring surface plasmon resonance (see, e.g., Sjolander & Urbaniczk, Anal. Chem. 63: 2338-2345 (1991) and Szabo et al., Curr. Opin. Struct. Biol. 5: 699-705 (1995)). "Surface plasmon resonance" or "biomolecular interaction analysis (BIA)" can be utilized to detect biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore). Changes in the mass at the binding surface (indicative of a binding event) result in alterations of the refractive index of light near the surface (the optical phenomenon of surface plasmon resonance (SPR)), resulting in a detectable signal which can be used as an indication of real-time reactions between biological molecules.

[0149] In another embodiment, the target molecule or test molecules are anchored to a solid phase, facilitating the detection of target molecule/test molecule complexes and separation of the complexes from free, uncomplexed molecules. The target molecule or test molecule is immobilized to the solid support. In an embodiment, the target molecule is anchored to a solid surface, and the test molecule, which is not anchored, can be labeled, either directly or indirectly, with detectable labels discussed herein.

[0150] It may be desirable to immobilize a target molecule, an anti-target molecule antibody, and/or test molecules to facilitate separation of target molecule/test molecule complexes from uncomplexed forms, as well as to accommodate automation of the assay. The attachment between a test molecule and/or target molecule and the solid support may be covalent or non-covalent (see, e.g., U.S. Patent No. 6,022,688 for non-covalent attachments). The solid support may be one or more surfaces of the system, such as one or more surfaces in each well of a microtiter plate, a surface of a silicon wafer, a surface of a bead (see, e.g., Lam, Nature 354: 82-84 (1991)) that is optionally linked to another solid support, or a channel in a microfluidic device, for example. Types of solid supports, linker molecules for covalent and non-covalent attachments to solid supports, and methods for immobilizing nucleic acids and other molecules to solid supports are well known (see, e.g., U.S. Patent Nos. 6,261,776; 5,900,481; 6,133,436; and 6,022,688; and WIPO publication WO 01/18234).

[0151] In an embodiment, target molecule may be immobilized to surfaces via biotin and streptavidin. For example, biotinylated target polypeptide can be prepared from biotin-NHS (N-hydroxy-succinimide) using techniques known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, IL), and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). In another embodiment, a target polypeptide can be prepared as a fusion polypeptide. For example, glutathione-S-transferase/target polypeptide fusion can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, MO) or glutathione derivitized microtiter plates, which are then combined with a test molecule

under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components, or the matrix is immobilized in the case of beads, and complex formation is determined directly or indirectly as described above. Alternatively, the complexes can be dissociated from the matrix, and the level of target molecule binding or activity is determined using standard techniques.

[0152] In an embodiment, the non-immobilized component is added to the coated surface containing the anchored component. After the reaction is complete, unreacted components are removed (e.g., by washing) under conditions such that a significant percentage of complexes formed will remain immobilized to the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of manners. Where the previously non-immobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface, e.g., by adding a labeled antibody specific for the immobilized component, where the antibody, in turn, can be directly labeled or indirectly labeled with, e.g., a labeled anti-Ig antibody.

[0153] In another embodiment, an assay is performed utilizing antibodies that specifically bind target molecule or test molecule but do not interfere with binding of the target molecule to the test molecule. Such antibodies can be derivitized to a solid support, and unbound target molecule may be immobilized by antibody conjugation. Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies reactive with the target molecule, as well as enzyme-linked assays which rely on detecting an enzymatic activity associated with the target molecule.

[0154] Cell free assays also can be conducted in a liquid phase. In such an assay, reaction products are separated from unreacted components, by any of a number of standard techniques, including but not limited to: differential centrifugation (see, e.g., Rivas, G., and Minton, Trends Biochem Sci Aug; 18(8): 284-7 (1993)); chromatography (gel filtration chromatography, ion-exchange chromatography); electrophoresis (see, e.g., Ausubel et al., eds. Current Protocols in Molecular Biology, J. Wiley: New York (1999)); and immunoprecipitation (see, e.g., Ausubel et al., eds., supra). Media and chromatographic techniques are known to one skilled in the art (see, e.g., Heegaard, J Mol. Recognit. Winter; 11(1-6): 141-8 (1998); Hage & Tweed, J. Chromatogr. B Biomed. Sci. Appl. Oct 10; 699 (1-2): 499-525 (1997)). Further, fluorescence energy transfer may also be conveniently utilized, as described herein, to detect binding without further purification of the complex from solution.

[0155] In another embodiment, modulators of target molecule expression are identified. For example, a cell or cell free mixture is contacted with a candidate compound and the expression of target mRNA or target polypeptide is evaluated relative to the level of expression of target mRNA or target

polypeptide in the absence of the candidate compound. When expression of target mRNA or target polypeptide is greater in the presence of the candidate compound than in its absence, the candidate compound is identified as an agonist of target mRNA or target polypeptide expression. Alternatively, when expression of target mRNA or target polypeptide is less (e.g., less with statistical significance) in the presence of the candidate compound than in its absence, the candidate compound is identified as an antagonist or inhibitor of target mRNA or target polypeptide expression. The level of target mRNA or target polypeptide expression can be determined by methods described herein.

[0156] In another embodiment, binding partners that interact with a target molecule are detected. The target molecules can interact with one or more cellular or extracellular macromolecules, such as polypeptides *in vivo*, and these interacting molecules are referred to herein as "binding partners." Binding partners can agonize or antagonize target molecule biological activity. Also, test molecules that agonize or antagonize interactions between target molecules and binding partners can be useful as therapeutic molecules as they can up-regulate or down-regulated target molecule activity *in vivo* and thereby treat osteoarthritis.

[0157] Binding partners of target molecules can be identified by methods known in the art. For example, binding partners may be identified by lysing cells and analyzing cell lysates by electrophoretic techniques. Alternatively, a two-hybrid assay or three-hybrid assay can be utilized (see, e.g., U.S. Patent No. 5,283,317; Zervos et al., Cell 72:223-232 (1993); Madura et al., J. Biol. Chem. 268: 12046-12054 (1993); Bartel et al., Biotechniques 14: 920-924 (1993); Iwabuchi et al., Oncogene 8: 1693-1696 (1993); and Brent WO94/10300). A two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. The assay often utilizes two different DNA constructs. In one construct, a IL1RL1 nucleic acid (sometimes referred to as the "bait") is fused to a gene encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In another construct, a DNA sequence from a library of DNA sequences that encodes a potential binding partner (sometimes referred to as the "prey") is fused to a gene that encodes an activation domain of the known transcription factor. Sometimes, a IL1RL1 nucleic acid can be fused to the activation domain. If the "bait" and the "prey" molecules interact in vivo, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ) which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected and cell colonies containing the functional transcription factor can be isolated and used to identify the potential binding partner.

[0158] In an embodiment for identifying test molecules that antagonize or agonize complex formation between target molecules and binding partners, a reaction mixture containing the target molecule and the binding partner is prepared, under conditions and for a time sufficient to allow complex formation. The reaction mixture often is provided in the presence or absence of the test molecule. The

test molecule can be included initially in the reaction mixture, or can be added at a time subsequent to the addition of the target molecule and its binding partner. Control reaction mixtures are incubated without the test molecule or with a placebo. Formation of any complexes between the target molecule and the binding partner then is detected. Decreased formation of a complex in the reaction mixture containing test molecule as compared to in a control reaction mixture indicates that the molecule antagonizes target molecule/binding partner complex formation. Alternatively, increased formation of a complex in the reaction mixture containing test molecule as compared to in a control reaction mixture indicates that the molecule agonizes target molecule/binding partner complex formation. In another embodiment, complex formation of target molecule/binding partner can be compared to complex formation of mutant target molecule/binding partner (e.g., amino acid modifications in a target polypeptide). Such a comparison can be important in those cases where it is desirable to identify test molecules that modulate interactions of mutant but not non-mutated target gene products.

[0159] The assays can be conducted in a heterogeneous or homogeneous format. In heterogeneous assays, target molecule and/or the binding partner are immobilized to a solid phase, and complexes are detected on the solid phase at the end of the reaction. In homogeneous assays, the entire reaction is carried out in a liquid phase. In either approach, the order of addition of reactants can be varied to obtain different information about the molecules being tested. For example, test compounds that agonize target molecule/binding partner interactions can be identified by conducting the reaction in the presence of the test molecule in a competition format. Alternatively, test molecules that agonize preformed complexes, e.g., molecules with higher binding constants that displace one of the components from the complex, can be tested by adding the test compound to the reaction mixture after complexes have been formed.

[0160] In a heterogeneous assay embodiment, the target molecule or the binding partner is anchored onto a solid surface (e.g., a microtiter plate), while the non-anchored species is labeled, either directly or indirectly. The anchored molecule can be immobilized by non-covalent or covalent attachments. Alternatively, an immobilized antibody specific for the molecule to be anchored can be used to anchor the molecule to the solid surface. The partner of the immobilized species is exposed to the coated surface with or without the test molecule. After the reaction is complete, unreacted components are removed (e.g., by washing) such that a significant portion of any complexes formed will remain immobilized on the solid surface. Where the non-immobilized species is pre-labeled, the detection of label immobilized on the surface is indicative of complex. Where the non-immobilized species is not pre-labeled, an indirect label can be used to detect complexes anchored to the surface; e.g., by using a labeled antibody specific for the initially non-immobilized species. Depending upon the order of addition of reaction components, test compounds that inhibit complex formation or that disrupt preformed complexes can be detected.

[0161] In another embodiment, the reaction can be conducted in a liquid phase in the presence or absence of test molecule, where the reaction products are separated from unreacted components, and the complexes are detected (e.g., using an immobilized antibody specific for one of the binding components to anchor any complexes formed in solution, and a labeled antibody specific for the other partner to detect anchored complexes). Again, depending upon the order of addition of reactants to the liquid phase, test compounds that inhibit complex or that disrupt preformed complexes can be identified.

[0162] In an alternate embodiment, a homogeneous assay can be utilized. For example, a preformed complex of the target gene product and the interactive cellular or extracellular binding partner product is prepared. One or both of the target molecule or binding partner is labeled, and the signal generated by the label(s) is quenched upon complex formation (e.g., U.S. Patent No. 4,109,496 that utilizes this approach for immunoassays). Addition of a test molecule that competes with and displaces one of the species from the preformed complex will result in the generation of a signal above background. In this way, test substances that disrupt target molecule/binding partner complexes can be identified.

[0163] Candidate therapeutics for treating osteoarthritis are identified from a group of test molecules that interact with a target molecule. Test molecules are normally ranked according to the degree with which they modulate (e.g., agonize or antagonize) a function associated with the target molecule (e.g., DNA replication and/or processing, RNA transcription and/or processing, polypeptide production and/or processing, and/or biological function/activity), and then top ranking modulators are selected. Also, pharmacogenomic information described herein can determine the rank of a modulator. The top 10% of ranked test molecules often are selected for further testing as candidate therapeutics, and sometimes the top 15%, 20%, or 25% of ranked test molecules are selected for further testing as candidate therapeutics. Candidate therapeutics typically are formulated for administration to a subject.

#### Therapeutic Formulations

[0164] Formulations and pharmaceutical compositions typically include in combination with a pharmaceutically acceptable carrier one or more target molecule modulators. The modulator often is a test molecule identified as having an interaction with a target molecule by a screening method described above. The modulator may be a compound, an antisense nucleic acid, a ribozyme, an antibody, or a binding partner. Also, formulations may comprise a target polypeptide or fragment thereof in combination with a pharmaceutically acceptable carrier.

[0165] As used herein, the term "pharmaceutically acceptable carrier" includes solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions. Pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

[0166] A pharmaceutical composition typically is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[0167] Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, *e.g.*, gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

[0168] Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL<sup>TM</sup> (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, sodium chloride in

the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[0169] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0170] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[0171] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art. Molecules can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[0172] In one embodiment, active molecules are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations will be apparent to those skilled in the art. Materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

[0173] It is advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity

of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0174] Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD<sub>50</sub>/ED<sub>50</sub>. Molecules which exhibit high therapeutic indices are preferred. While molecules that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

[0175] The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such molecules lies preferably within a range of circulating concentrations that include the  $ED_{50}$  with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any molecules used in the methods described herein, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the  $IC_{50}$  (*i.e.*, the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

[0176] As defined herein, a therapeutically effective amount of protein or polypeptide (*i.e.*, an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, sometimes about 0.01 to 25 mg/kg body weight, often about 0.1 to 20 mg/kg body weight, and more often about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The protein or polypeptide can be administered one time per week for between about 1 to 10 weeks, sometimes between 2 to 8 weeks, often between about 3 to 7 weeks, and more often for about 4, 5, or 6 weeks. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments.

[0177] With regard to polypeptide formulations, featured herein is a method for treating osteoarthritis in a subject, which comprises contacting one or more cells in the subject with a first polypeptide, where the subject comprises a second polypeptide having one or more polymorphic variations associated with cancer, and where the first polypeptide comprises fewer polymorphic

variations associated with cancer than the second polypeptide. The first and second polypeptides are encoded by a nucleic acid which comprises a nucleotide sequence in SEQ ID NO: 1-4; a nucleotide sequence which encodes a polypeptide consisting of an amino acid sequence encoded by a nucleotide sequence referenced in SEQ ID NO: 1-4; a nucleotide sequence which encodes a polypeptide that is 90% or more identical to an amino acid sequence encoded by a nucleotide sequence of SEQ ID NO: 1-4 and a nucleotide sequence 90% or more identical to a nucleotide sequence in SEQ ID NO: 1-4. The subject often is a human.

[0178] For antibodies, a dosage of 0.1 mg/kg of body weight (generally 10 mg/kg to 20 mg/kg) is often utilized. If the antibody is to act in the brain, a dosage of 50 mg/kg to 100 mg/kg is often appropriate. Generally, partially human antibodies and fully human antibodies have a longer half-life within the human body than other antibodies. Accordingly, lower dosages and less frequent administration is often possible. Modifications such as lipidation can be used to stabilize antibodies and to enhance uptake and tissue penetration (e.g., into the brain). A method for lipidation of antibodies is described by Cruikshank et al., J. Acquired Immune Deficiency Syndromes and Human Retrovirology 14:193 (1997).

[0179] Antibody conjugates can be used for modifying a given biological response, the drug moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a polypeptide such as tumor necrosis factor, alpha-interferon, beta-interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator; or, biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors. Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate as described by Segal in U.S. Patent No. 4,676,980.

[0180] For compounds, exemplary doses include milligram or microgram amounts of the compound per kilogram of subject or sample weight, for example, about 1 microgram per kilogram to about 500 milligrams per kilogram, about 100 micrograms per kilogram to about 5 milligrams per kilogram, or about 1 microgram per kilogram to about 50 micrograms per kilogram. It is understood that appropriate doses of a small molecule depend upon the potency of the small molecule with respect to the expression or activity to be modulated. When one or more of these small molecules is to be administered to an animal (e.g., a human) in order to modulate expression or activity of a polypeptide or nucleic acid described herein, a physician, veterinarian, or researcher may, for example, prescribe a relatively low dose at first, subsequently increasing the dose until an appropriate response is obtained. In addition, it is understood that the specific dose level for any particular animal subject will depend upon a variety of

factors including the activity of the specific compound employed, the age, body weight, general health, gender, and diet of the subject, the time of administration, the route of administration, the rate of excretion, any drug combination, and the degree of expression or activity to be modulated.

[0181] With regard to nucleic acid formulations, gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see, e.g., U.S. Patent 5,328,470) or by stereotactic injection (see e.g., Chen et al., (1994) Proc. Natl. Acad. Sci. USA 91:3054-3057). Pharmaceutical preparations of gene therapy vectors can include a gene therapy vector in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells (e.g., retroviral vectors) the pharmaceutical preparation can include one or more cells which produce the gene delivery system. Examples of gene delivery vectors are described herein.

## Therapeutic Methods

[0182] A therapeutic formulation described above can be administered to a subject in need of a therapeutic for inducing a desired biological response.. Therapeutic formulations can be administered by any of the paths described herein. With regard to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from pharmacogenomic analyses described herein.

[0183] As used herein, the term "treatment" is defined as the application or administration of a therapeutic formulation to a subject, or application or administration of a therapeutic agent to an isolated tissue or cell line from a subject with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect osteoarthritis, symptoms of osteoarthritis or a predisposition towards osteoarthritis. A therapeutic formulation includes, but is not limited to, small molecules, peptides, antibodies, ribozymes and antisense oligonucleotides. Administration of a therapeutic formulation can occur prior to the manifestation of symptoms characteristic of osteoarthritis, such that osteoarthritis is prevented or delayed in its progression. The appropriate therapeutic composition can be determined based on screening assays described herein.

[0184] As discussed, successful treatment of osteoarthritis can be brought about by techniques that serve to agonize target molecule expression or function, or alternatively, antagonize target molecule expression or function. These techniques include administration of modulators that include, but are not limited to, small organic or inorganic molecules; antibodies (including, for example, polyclonal, monoclonal, humanized, anti-idiotypic, chimeric or single chain antibodies, and Fab, F(ab')<sub>2</sub> and Fab expression library fragments, scFV molecules, and epitope-binding fragments thereof); and peptides, phosphopeptides, or polypeptides.

[0185] Further, antisense and ribozyme molecules that inhibit expression of the target gene can also be used to reduce the level of target gene expression, thus effectively reducing the level of target gene activity. Still further, triple helix molecules can be utilized in reducing the level of target gene activity. Antisense, ribozyme and triple helix molecules are discussed above. It is possible that the use of antisense, ribozyme, and/or triple helix molecules to reduce or inhibit mutant gene expression can also reduce or inhibit the transcription (triple helix) and/or translation (antisense, ribozyme) of mRNA produced by normal target gene alleles, such that the concentration of normal target gene product present can be lower than is necessary for a normal phenotype. In such cases, nucleic acid molecules that encode and express target gene polypeptides exhibiting normal target gene activity can be introduced into cells via gene therapy method. Alternatively, in instances in that the target gene encodes an extracellular polypeptide, it can be preferable to co-administer normal target gene polypeptide into the cell or tissue in order to maintain the requisite level of cellular or tissue target gene activity.

[0186] Another method by which nucleic acid molecules may be utilized in treating or preventing osteoarthritis is use of aptamer molecules specific for target molecules. Aptamers are nucleic acid molecules having a tertiary structure which permits them to specifically bind to ligands (see, e.g., Osborne, et al., Curr. Opin. Chem. Biol. 1(1): 5-9 (1997); and Patel, D. J., Curr. Opin. Chem. Biol. Jun; 1(1): 32-46 (1997)).

[0187] Yet another method of utilizing nucleic acid molecules for osteoarthritis treatment is gene therapy, which can also be referred to as allele therapy. Provided herein is a gene therapy method for treating osteoarthritis in a subject, which comprises contacting one or more cells in the subject or from the subject with a nucleic acid having a first nucleotide sequence (e.g., the first nucleotide sequence is identical to or substantially identical to a nucleotide sequence of SEQ ID NO: 1-4). Genomic DNA in the subject comprises a second nucleotide sequence having one or more polymorphic variations associated with osteoarthritis (e.g., the second nucleotide sequence is identical to or substantially identical to a nucleotide sequence of SEQ ID NO: 1). The first and second nucleotide sequences typically are substantially identical to one another, and the first nucleotide sequence comprises fewer polymorphic variations associated with osteoarthritis than the second nucleotide sequence. The first nucleotide sequence may comprise a gene sequence that encodes a full-length polypeptide or a fragment thereof. The subject is often a human. Allele therapy methods often are utilized in conjunction with a method of first determining whether a subject has genomic DNA that includes polymorphic variants associated with osteoarthritis.

[0188] In another allele therapy embodiment, provided herein is a method which comprises contacting one or more cells in the subject or from the subject with a polypeptide encoded by a nucleic acid having a first nucleotide sequence (e.g., the first nucleotide sequence is identical to or substantially identical to the nucleotide sequence of SEQ ID NO: 1-4). Genomic DNA in the subject comprises a

second nucleotide sequence having one or more polymorphic variations associated with osteoarthritis (e.g., the second nucleotide sequence is identical to or substantially identical to a nucleotide sequence of SEQ ID NO: 1). The first and second nucleotide sequences typically are substantially identical to one another, and the first nucleotide sequence comprises fewer polymorphic variations associated with osteoarthritis than the second nucleotide sequence. The first nucleotide sequence may comprise a gene sequence that encodes a full-length polypeptide or a fragment thereof. The subject is often a human.

[0189] For antibody-based therapies, antibodies can be generated that are both specific for target molecules and that reduce target molecule activity. Such antibodies may be administered in instances where antagonizing a target molecule function is appropriate for the treatment of osteoarthritis.

[0190] In circumstances where stimulating antibody production in an animal or a human subject by injection with a target molecule is harmful to the subject, it is possible to generate an immune response against the target molecule by use of anti-idiotypic antibodies (see, e.g., Herlyn, Ann. Med.;31(1): 66-78 (1999); and Bhattacharya-Chatterjee & Foon, Cancer Treat. Res.; 94: 51-68 (1998)). Introducing an anti-idiotypic antibody to a mammal or human subject often stimulates production of anti-anti-idiotypic antibodies, which typically are specific to the target molecule. Vaccines directed to osteoarthritis also may be generated in this fashion.

[0191] In instances where the target molecule is intracellular and whole antibodies are used, internalizing antibodies may be preferred. Lipofectin or liposomes can be used to deliver the antibody or a fragment of the Fab region that binds to the target antigen into cells. Where fragments of the antibody are used, the smallest inhibitory fragment that binds to the target antigen is preferred. For example, peptides having an amino acid sequence corresponding to the Fv region of the antibody can be used. Alternatively, single chain neutralizing antibodies that bind to intracellular target antigens can also be administered. Such single chain antibodies can be administered, for example, by expressing nucleotide sequences encoding single-chain antibodies within the target cell population (see, e.g., Marasco et al., Proc. Natl. Acad. Sci. USA 90: 7889-7893 (1993)).

[0192] Modulators can be administered to a patient at therapeutically effective doses to treat osteoarthritis. A therapeutically effective dose refers to an amount of the modulator sufficient to result in amelioration of symptoms of osteoarthritis. Toxicity and therapeutic efficacy of modulators can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD<sub>50</sub> (the dose lethal to 50% of the population) and the ED<sub>50</sub> (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD<sub>50</sub>/ED<sub>50</sub>. Modulators that exhibit large therapeutic indices are preferred. While modulators that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such molecules to the site of affected tissue in order to minimize potential damage to uninfected cells, thereby reducing side effects.

[0193] Data obtained from cell culture assays and animal studies can be used in formulating a range of dosages for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED<sub>50</sub> with little or no toxicity. The dosage can vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the methods described herein, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC<sub>50</sub> (*i.e.*, the concentration of the test compound that achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

[0194] Another example of effective dose determination for an individual is the ability to directly assay levels of "free" and "bound" compound in the serum of the test subject. Such assays may utilize antibody mimics and/or "biosensors" that have been created through molecular imprinting techniques. Molecules that modulate target molecule activity are used as a template, or "imprinting molecule", to spatially organize polymerizable monomers prior to their polymerization with catalytic reagents. The subsequent removal of the imprinted molecule leaves a polymer matrix which contains a repeated "negative image" of the compound and is able to selectively rebind the molecule under biological assay conditions. A detailed review of this technique can be seen in Ansell et al., Current Opinion in Biotechnology 7: 89-94 (1996) and in Shea, Trends in Polymer Science 2: 166-173 (1994). Such "imprinted" affinity matrixes are amenable to ligand-binding assays, whereby the immobilized monoclonal antibody component is replaced by an appropriately imprinted matrix. An example of the use of such matrixes in this way can be seen in Vlatakis, et al., Nature 361: 645-647 (1993). Through the use of isotope-labeling, the "free" concentration of compound which modulates target molecule expression or activity readily can be monitored and used in calculations of IC<sub>50</sub>. Such "imprinted" affinity matrixes can also be designed to include fluorescent groups whose photon-emitting properties measurably change upon local and selective binding of target compound. These changes readily can be assayed in real time using appropriate fiberoptic devices, in turn allowing the dose in a test subject to be quickly optimized based on its individual IC<sub>50</sub>. An example of such a "biosensor" is discussed in Kriz et al., Analytical Chemistry 67: 2142-2144 (1995).

[0195] The examples set forth below are intended to illustrate but not limit the invention.

#### **Examples**

[0196] In the following studies a group of subjects was selected according to specific parameters relating to osteoarthritis. Nucleic acid samples obtained from individuals in the study group were subjected to genetic analysis, which identified associations between osteoarthritis and a polymorphism in

the *IL1RL1* gene on chromosome two. The polymorphism was genotyped again in two replication cohorts consisting of individuals selected for OA. In addition, SNPs proximal to the incident polymorphism were identified and allelotyped in OA case and control pools. Methods are described for producing *IL1RL1* polypeptide and *IL1RL1* polypeptide variants in vitro or in vivo, *IL1RL1* nucleic acids or polypeptides and variants thereof are utilized for screening test molecules for those that interact with *IL1RL1* molecules. Test molecules identified as interactors with *IL1RL1* molecules and *IL1RL1* variants are further screened *in vivo* to determine whether they treat osteoarthritis.

# Example 1 Samples and Pooling Strategies

#### Sample Selection

[0197] Blood samples were collected from individuals diagnosed with knee osteoarthritis, which were referred to as case samples. Also, blood samples were collected from individuals not diagnosed with knee osteoarthritis as gender and age-matched controls. A database was created that listed all phenotypic trait information gathered from individuals for each case and control sample. Genomic DNA was extracted from each of the blood samples for genetic analyses.

## **DNA Extraction from Blood Samples**

[0198] Six to ten milliliters of whole blood was transferred to a 50 ml tube containing 27 ml of red cell lysis solution (RCL). The tube was inverted until the contents were mixed. Each tube was incubated for 10 minutes at room temperature and inverted once during the incubation. The tubes were then centrifuged for 20 minutes at 3000 x g and the supernatant was carefully poured off. 100-200 µl of residual liquid was left in the tube and was pipetted repeatedly to resuspend the pellet in the residual supernatant. White cell lysis solution (WCL) was added to the tube and pipetted repeatedly until completely mixed. While no incubation was normally required, the solution was incubated at 37°C or room temperature if cell clumps were visible after mixing until the solution was homogeneous. 2 ml of protein precipitation was added to the cell lysate. The mixtures were vortexed vigorously at high speed for 20 sec to mix the protein precipitation solution uniformly with the cell lysate, and then centrifuged for 10 minutes at 3000 x g. The supernatant containing the DNA was then poured into a clean 15 ml tube, which contained 7 ml of 100% isopropanol. The samples were mixed by inverting the tubes gently until white threads of DNA were visible. Samples were centrifuged for 3 minutes at 2000 x g and the DNA was visible as a small white pellet. The supernatant was decanted and 5 ml of 70% ethanol was added to each tube. Each tube was inverted several times to wash the DNA pellet, and then centrifuged for 1 minute at 2000 x g. The ethanol was decanted and each tube was drained on clean absorbent paper. The

DNA was dried in the tube by inversion for 10 minutes, and then 1000 µl of 1X TE was added. The size of each sample was estimated, and less TE buffer was added during the following DNA hydration step if the sample was smaller. The DNA was allowed to rehydrate overnight at room temperature, and DNA samples were stored at 2-8°C.

[0199] DNA was quantified by placing samples on a hematology mixer for at least 1 hour. DNA was serially diluted (typically 1:80, 1:160, 1:320, and 1:640 dilutions) so that it would be within the measurable range of standards. 125 µl of diluted DNA was transferred to a clear U-bottom microtitre plate, and 125 µl of 1X TE buffer was transferred into each well using a multichannel pipette. The DNA and 1X TE were mixed by repeated pipetting at least 15 times, and then the plates were sealed. 50 µl of diluted DNA was added to wells A5-H12 of a black flat bottom microtitre plate. Standards were inverted six times to mix them, and then 50 µl of 1X TE buffer was pipetted into well A1, 1000 ng/ml of standard was pipetted into well A2, 500 ng/ml of standard was pipetted into well A3, and 250 ng/ml of standard was pipetted into well A4. PicoGreen (Molecular Probes, Eugene, Oregon) was thawed and freshly diluted 1:200 according to the number of plates that were being measured. PicoGreen was vortexed and then 50µl was pipetted into all wells of the black plate with the diluted DNA. DNA and PicoGreen were mixed by pipetting repeatedly at least 10 times with the multichannel pipette. The plate was placed into a Fluoroskan Ascent Machine (microplate fluorometer produced by Labsystems) and the samples were allowed to incubate for 3 minutes before the machine was run using filter pairs 485 nm excitation and 538 nm emission wavelengths. Samples having measured DNA concentrations of greater than 450 ng/μl were re-measured for conformation. Samples having measured DNA concentrations of 20 ng/μl or less were re-measured for confirmation.

### Pooling Strategies - Discovery Cohort

[0200] Samples were derived from the Nottingham knee OA family study (UK) where index cases were identified through a knee replacement registry. Siblings were approached and assessed with knee x-rays and assigned status as affected or unaffected. In all 1,157 individuals were available. In order to create same-sex pools of appropriate sizes, 335 unrelated female individuals with OA from the Nottingham OA sample were selected for the case pool. The control pool was made up of unrelated female individuals from the St. Thomas twin study (England) with normal knee x-rays and without other indications of OA, regardless of anatomical location, as well as lacking family history of OA. The St. Thomas twin study consists of Caucasian, female participants from the St. Thomas' Hospital, London, adult-twin registry, which is a voluntary registry of >4,000 twin pairs ranging from 18 to 76 years of age. The female case samples and female control samples are described further in Table 1 below.

[0201] A select set of samples from each group were utilized to generate pools, and one pool was created for each group. Each individual sample in a pool was represented by an equal amount of genomic DNA. For example, where 25 ng of genomic DNA was utilized in each PCR reaction and there were 200 individuals in each pool, each individual would provide 125 pg of genomic DNA. Inclusion or exclusion of samples for a pool was based upon the following criteria: the sample was derived from an individual characterized as Caucasian; the sample was derived from an individual of British paternal and maternal descent; case samples were derived from individuals diagnosed with specific knee osteoarthritis (OA) and were recruited from an OA knee replacement clinic. Control samples were derived from individuals free of OA, family history of OA, and rheumatoid arthritis. Also, sufficient genomic DNA was extracted from each blood sample for all allelotyping and genotyping reactions performed during the study. Phenotype information from each individual was collected and included age of the individual, gender, family history of OA, general medical information (e.g., height, weight, thyroid disease, diabetes, psoriasis, hysterectomy), joint history (previous and current symptoms, joint-related operations, age at onset of symptoms, date of primary diagnosis, age of individual as of primary diagnosis and order of involvement), and knee-related findings (crepitus, restricted passive movement, bony swelling/deformity). Additional knee information included knee history, current symptoms, any major knee injury, menisectomy, knee replacement surgery, age of surgery, and treatment history (including hormone replace therapy (HRT)). Samples that met these criteria were added to appropriate pools based on disease status.

[0202] The selection process yielded the pools set forth in Table 1, which were used in the studies that follow:

TABLE 1

	Female case	Female control
Pool size (Number)	335	335
Pool Criteria (ex: case/control)	control	case
Mean Age (ex: years)	57.21	69.95

# Example 2 Association of Polymorphic Variants with Osteoarthritis

[0203] A whole-genome screen was performed to identify particular SNPs associated with occurrence of osteoarthritis. As described in Example 1, two sets of samples were utilized, which included samples from female individuals having knee osteoarthritis (osteoarthritis cases), and samples

from female individuals not having knee osteoarthritis (female controls). The initial screen of each pool was performed in an allelotyping study, in which certain samples in each group were pooled. By pooling DNA from each group, an allele frequency for each SNP in each group was calculated. These allele frequencies were then compared to one another. Particular SNPs were considered as being associated with osteoarthritis when allele frequency differences calculated between case and control pools were statistically significant. SNP disease association results obtained from the allelotyping study were then validated by genotyping each associated SNP across all samples from each pool. The results of the genotyping then were analyzed, allele frequencies for each group were calculated from the individual genotyping results, and a p-value was calculated to determine whether the case and control groups had statistically significant differences in allele frequencies for a particular SNP. When the genotyping results agreed with the original allelotyping results, the SNP disease association was considered validated at the genetic level.

#### SNP Panel Used for Genetic Analyses

[0204] A whole-genome SNP screen began with an initial screen of approximately 25,000 SNPs over each set of disease and control samples using a pooling approach. The pools studied in the screen are described in Example 1. The SNPs analyzed in this study were part of a set of 25,488 SNPs confirmed as being statistically polymorphic as each is characterized as having a minor allele frequency of greater than 10%. The SNPs in the set reside in genes or in close proximity to genes, and many reside in gene exons. Specifically, SNPs in the set are located in exons, introns, and within 5,000 base-pairs upstream of a transcription start site of a gene. In addition, SNPs were selected according to the following criteria: they are located in ESTs; they are located in Locuslink or Ensembl genes; and they are located in Genomatix promoter predictions. SNPs in the set were also selected on the basis of even spacing across the genome, as depicted in Table 2.

[0205] A case-control study design using a whole genome association strategy involving approximately 28,000 single nucleotide polymorphisms (SNPs) was employed. Approximately 25,000 SNPs were evenly spaced in gene-based regions of the human genome with a median inter-marker distance of about 40,000 base pairs. Additionally, approximately 3,000 SNPs causing amino acid substitutions in genes described in the literature as candidates for various diseases were used. The case-control study samples were of female Caucasian origin (British paternal and maternal descent) 670 individuals were equally distributed in two groups: female controls and female cases. The whole genome association approach was first conducted on 2 DNA pools representing the 2 groups. Significant markers were confirmed by individual genotyping.

**TABLE 2** 

General Stat	istics	Spacing Statistics		
Total # of SNPs # of Exonic SNPs	25,488 >4,335 (17%)	Median Minimum*	37,058 bp 1,000 bp	
# SNPs with refSNP ID	20,776 (81%)	Maximum*	3,000,000 bp	
Gene Coverage	>10,000	Mean	122,412 bp	
Chromosome Coverage	All	Std Deviation *Excludes outliers	373,325 bp	

### Allelotyping and Genotyping Results

[0206] The genetic studies summarized above and described in more detail below identified an allelic variant in the IL1RL1 gene that is associated with osteoarthritis.

# Assay for Verifying, Allelotyping, and Genotyping SNPs

[0207] A MassARRAY<sup>TM</sup> system (Sequenom, Inc.) was utilized to perform SNP genotyping in a high-throughput fashion. This genotyping platform was complemented by a homogeneous, single-tube assay method (hME<sup>TM</sup> or homogeneous MassEXTEND<sup>TM</sup> (Sequenom, Inc.)) in which two genotyping primers anneal to and amplify a genomic target surrounding a polymorphic site of interest. A third primer (the MassEXTEND<sup>TM</sup> primer), which is complementary to the amplified target up to but not including the polymorphism, was then enzymatically extended one or a few bases through the polymorphic site and then terminated.

[0208] For each polymorphism, SpectroDESIGNER<sup>TM</sup> software (Sequenom, Inc.) was used to generate a set of PCR primers and a MassEXTEND<sup>TM</sup> primer which where used to genotype the polymorphism. Other primer design software could be used or one of ordinary skill in the art could manually design primers based on his or her knowledge of the relevant factors and considerations in designing such primers. Table 3 shows PCR primers and Table 4 shows extension primers used for analyzing polymorphisms. The initial PCR amplification reaction was performed in a 5 μl total volume containing 1X PCR buffer with 1.5 mM MgCl<sub>2</sub> (Qiagen), 200 μM each of dATP, dGTP, dTTP (Gibco-BRL), 2.5 ng of genomic DNA, 0.1 units of HotStar DNA polymerase (Qiagen), and 200 nM each of forward and reverse PCR primers specific for the polymorphic region of interest.

**TABLE 3: PCR Primers** 

SNP Reference	Forward PCR primer	Reverse PCR primer
rs1041973	ACGTTGGATGGGGACTTCTGACAATACAGG	ACGTTGGATGAATCGTGTGTTTGCCTCAGG

[0209] Samples were incubated at 95°C for 15 minutes, followed by 45 cycles of 95°C for 20 seconds, 56°C for 30 seconds, and 72°C for 1 minute, finishing with a 3 minute final extension at 72°C. Following amplification, shrimp alkaline phosphatase (SAP) (0.3 units in a 2 μl volume) (Amersham Pharmacia) was added to each reaction (total reaction volume was 7 μl) to remove any residual dNTPs that were not consumed in the PCR step. Samples were incubated for 20 minutes at 37°C, followed by 5 minutes at 85°C to denature the SAP.

[0210] Once the SAP reaction was complete, a primer extension reaction was initiated by adding a polymorphism-specific MassEXTEND™ primer cocktail to each sample. Each MassEXTEND™ cocktail included a specific combination of dideoxynucleotides (ddNTPs) and deoxynucleotides (dNTPs) used to distinguish polymorphic alleles from one another. Methods for verifying, allelotyping and genotyping SNPs are disclosed, for example, in U.S. Pat. No. 6,258,538, the content of which is hereby incorporated by reference. In Table 4, ddNTPs are shown and the fourth nucleotide not shown is the dNTP.

**TABLE 4: Extension Primers** 

SNP Reference	Extend Probe	Termination Mix
rs1041973	ATACCAGAATCAGCAACT	ACT

[0211] The MassEXTEND<sup>TM</sup> reaction was performed in a total volume of 9 μl, with the addition of 1X ThermoSequenase buffer, 0.576 units of ThermoSequenase (Amersham Pharmacia), 600 nM MassEXTEND<sup>TM</sup> primer, 2 mM of ddATP and/or ddCTP and/or ddGTP and/or ddTTP, and 2 mM of dATP or dCTP or dGTP or dTTP. The deoxy nucleotide (dNTP) used in the assay normally was complementary to the nucleotide at the polymorphic site in the amplicon. Samples were incubated at 94°C for 2 minutes, followed by 55 cycles of 5 seconds at 94°C, 5 seconds at 52°C, and 5 seconds at 72°C.

[0212] Following incubation, samples were desalted by adding 16 μl of water (total reaction volume was 25 μl), 3 mg of SpectroCLEAN<sup>TM</sup> sample cleaning beads (Sequenom, Inc.) and allowed to incubate for 3 minutes with rotation. Samples were then robotically dispensed using a piezoelectric dispensing device (SpectroJET<sup>TM</sup> (Sequenom, Inc.)) onto either 96-spot or 384-spot silicon chips containing a matrix that crystallized each sample (SpectroCHIP<sup>TM</sup> (Sequenom, Inc.)). Subsequently, MALDI-TOF mass spectrometry (Biflex and Autoflex MALDI-TOF mass spectrometers (Bruker Daltonics) can be used) and SpectroTYPER RT<sup>TM</sup> software (Sequenom, Inc.) were used to analyze and interpret the SNP genotype for each sample.

### Genetic Analysis

[0213] Minor allelic frequencies for the polymorphisms set forth in Table A were verified as being 10% or greater using the extension assay described above in a group of samples isolated from 92 individuals originating from the state of Utah in the United States, Venezuela and France (Coriell cell repositories).

[0214] Genotyping results are shown for female pools in Table 5. In Table 5, "AF" refers to allelic frequency; and "F case" and "F control" refer to female case and female control groups, respectively.

 SNP Reference
 AF F case
 AF F control
 p-value

 rs1041973
 A = 0.189 A = 0.233 C = 0.767
 0.0539

**TABLE 5: Genotyping Results** 

[0215] All of the single marker alleles set forth in Table A were considered validated, since the genotyping data agreed with the allelotyping data and each SNP significantly associated with osteoarthritis. Particularly significant associations with osteoarthritis are indicated by a calculated p-value of less than 0.05 for genotype results. SNP rs1041973 has a calculated p-value of greater than 0.05 (0.0539), but was included because it is an exonic SNP located in the IL1RL1 gene, which plays a role in inflammation and is a compelling target for osteoarthritis.

# Example 3

### Association of Polymorphic Variants with Osteoarthritis in Replication Cohorts

[0216] The single marker polymorphism set forth in Table A was genotyped again in two replication cohorts consisting of individuals selected for OA.

### Sample Selection and Pooling Strategies – Replication Sample 1

[0217] A second case control sample (replication sample #1) was created by using 100 Caucasian female cases from Chingford, UK, and 148 unrelated female cases from the St. Thomas twin study. Cases were defined as having Kellgren-Lawrence (KL) scores of at least 2 in at least one knee x-ray. In addition, 199 male knee replacement cases from Nottingham were included. (For a cohort description, see the Nottingham description provided in Example 1). The control pool was made up of unrelated female individuals from the St. Thomas twin study (England) with normal knee x-rays and without other indications of OA, regardless of anatomical location, as well as lacking family history of OA. The St. Thomas twin study consists of Caucasian, female participants from the St. Thomas' Hospital, London, adult-twin registry, which is a voluntary registry of >4,000 twin pairs ranging from 18 to 76 years of age.

The replication sample 1 cohort was used to replicate the initial results. Table 6 below summarizes the selected phenotype data collected from the case and control individuals.

**TABLE 6** 

Phenotype	Female cases (n=248): median (range)/ (n,%)	Male cases (n=199): median (range)/ (n,%)	Female controls (n=313): mean (range)/ (n,%)
Age	59 (39- 73)	66 (45- 73)	55 (50- 72)
Height (cm)	162 (141- 178)	175 (152- 198)	162 (141- 176)
Weight (kg)	68 (51- 123)	86 (62- 127)	64 (40- 111)
Body mass index (kg/m²)	26 (18- 44)	29 (21- 41)	24 (18- 46)
Kellgren- Lawrence* left knee	0 (63, 26%), 1 (20, 8%), 2 (105, 43%), 3 (58, 23%), 4 (1, 0%)	NA	NA
Kellgren- Lawrence* right knee	0 (43, 7%), 1 (18, 7%), 2 (127, 52%), 3 (57, 23%), 4 (1, 0%)	NA	NA
KL* >2 both knees	No (145, 59%), Yes (101, 41%)	NA	NA
KL* >2 either knee	No (0, 0%), Yes (248, 100%)	NA	NA

<sup>\* 0:</sup> normal, 1: doubtful, 2: definite osteophyte (bony protuberance), 3: joint space narrowing (with or without osteophyte), 4: joint deformity

### Sample Selection and Pooling Strategies – Replication Sample 2

[0218] A third case control sample (replication sample #2) was created by using individuals with symptoms of OA from Newfoundland, Canada. These individuals were recruited and examined by rheumatologists. Affected joints were x-rayed and a final diagnosis of definite or probable OA was made according to American College of Rheumatology criteria by a single rheumatologist to avoid any interexaminer diagnosis variability. Controls were recruited from volunteers without any symptoms from the musculoskeletal system based on a normal joint exam performed by a rheumatologist. Only cases with a diagnosis of definite OA were included in the study. Only individuals of Caucasian origin were included. The cases consisted of 228 individuals with definite knee OA, 106 individuals with definite hip OA, and 74 individuals with hip OA.

TABLE 7

Phenotype	Case	Control
Age at Visit	62.7	52.5

Phenotype	Case	Control
Sex (Female/Male)	227/119	174/101
Knee OA Xray: No	35% (120)	80% (16)
Unknown	1% (4)	0% (0)
Yes	64% (221)	20% (4)
Hip OA Xray: No	63% (215)	80% (16)
Unknown	2% (7)	0% (0)
Yes	35% (121)	20% (4)

### Assay for Verifying, Allelotyping, and Genotyping SNPs

[0219] Genotyping of the replication cohorts described in Tables 6 and 7 was performed using the same methods used for the original genotyping, as described herein. A MassARRAY<sup>TM</sup> system (Sequenom, Inc.) was utilized to perform SNP genotyping in a high-throughput fashion. This genotyping platform was complemented by a homogeneous, single-tube assay method (hME<sup>TM</sup> or homogeneous MassEXTEND<sup>TM</sup> (Sequenom, Inc.)) in which two genotyping primers anneal to and amplify a genomic target surrounding a polymorphic site of interest. A third primer (the MassEXTEND<sup>TM</sup> primer), which is complementary to the amplified target up to but not including the polymorphism, was then enzymatically extended one or a few bases through the polymorphic site and then terminated.

[0220] For each polymorphism, SpectroDESIGNER<sup>TM</sup> software (Sequenom, Inc.) was used to generate a set of PCR primers and a MassEXTEND<sup>TM</sup> primer which where used to genotype the polymorphism. Other primer design software could be used or one of ordinary skill in the art could manually design primers based on his or her knowledge of the relevant factors and considerations in designing such primers. Table 3 shows PCR primers and Table 4 shows extension probes used for analyzing (*e.g.*, genotyping) polymorphisms in the replication cohorts. The initial PCR amplification reaction was performed in a 5 μl total volume containing 1X PCR buffer with 1.5 mM MgCl<sub>2</sub> (Qiagen), 200 μM each of dATP, dGTP, dCTP, dTTP (Gibco-BRL), 2.5 ng of genomic DNA, 0.1 units of HotStar DNA polymerase (Qiagen), and 200 nM each of forward and reverse PCR primers specific for the polymorphic region of interest.

[0221] Samples were incubated at 95°C for 15 minutes, followed by 45 cycles of 95°C for 20 seconds, 56°C for 30 seconds, and 72°C for 1 minute, finishing with a 3 minute final extension at 72°C. Following amplification, shrimp alkaline phosphatase (SAP) (0.3 units in a 2 μl volume) (Amersham Pharmacia) was added to each reaction (total reaction volume was 7 μl) to remove any residual dNTPs

that were not consumed in the PCR step. Samples were incubated for 20 minutes at 37°C, followed by 5 minutes at 85°C to denature the SAP.

[0222] Once the SAP reaction was complete, a primer extension reaction was initiated by adding a polymorphism-specific MassEXTEND™ primer cocktail to each sample. Each MassEXTEND™ cocktail included a specific combination of dideoxynucleotides (ddNTPs) and deoxynucleotides (dNTPs) used to distinguish polymorphic alleles from one another. Methods for verifying, allelotyping and genotyping SNPs are disclosed, for example, in U.S. Pat. No. 6,258,538, the content of which is hereby incorporated by reference. In Table 7, ddNTPs are shown and the fourth nucleotide not shown is the dNTP.

[0223] The MassEXTEND<sup>TM</sup> reaction was performed in a total volume of 9 μl, with the addition of 1X ThermoSequenase buffer, 0.576 units of ThermoSequenase (Amersham Pharmacia), 600 nM MassEXTEND<sup>TM</sup> primer, 2 mM of ddATP and/or ddCTP and/or ddGTP and/or ddTTP, and 2 mM of dATP or dCTP or dGTP or dTTP. The deoxy nucleotide (dNTP) used in the assay normally was complementary to the nucleotide at the polymorphic site in the amplicon. Samples were incubated at 94°C for 2 minutes, followed by 55 cycles of 5 seconds at 94°C, 5 seconds at 52°C, and 5 seconds at 72°C.

[0224] Following incubation, samples were desalted by adding 16 μl of water (total reaction volume was 25 μl), 3 mg of SpectroCLEAN<sup>TM</sup> sample cleaning beads (Sequenom, Inc.) and allowed to incubate for 3 minutes with rotation. Samples were then robotically dispensed using a piezoelectric dispensing device (SpectroJET<sup>TM</sup> (Sequenom, Inc.)) onto either 96-spot or 384-spot silicon chips containing a matrix that crystallized each sample (SpectroCHIP<sup>TM</sup> (Sequenom, Inc.)). Subsequently, MALDI-TOF mass spectrometry (Biflex and Autoflex MALDI-TOF mass spectrometers (Bruker Daltonics) can be used) and SpectroTYPER RT<sup>TM</sup> software (Sequenom, Inc.) were used to analyze and interpret the SNP genotype for each sample.

#### Genetic Analysis

[0225] Genotyping results for replication cohorts #1 and #2 are provided in Tables 8 and 9, respectively.

TABLE 8

rsID	Replication #1 rsID (Mixed Male/Female cases and Female controls)				
	AF OA Con	AF OA Cas	Delta	P-value	P-value
rs1041973	0.77	0.79	-0.02	0.357	NA

**TABLE 9** 

rsID	Replication #2 (Newfoundland) (Male/Female cases and controls)				Meta-analysis Disc. + Rep #2
	AF OA Con	AF OA Cas	Delta	P-value	Not Done
rs1041973	0.78	0.79	-0.016	0.510	

[0226] To combine the evidence for association from multiple sample collections, a meta-analysis procedure was employed. The allele frequencies were compared between cases and controls within the discovery sample, as well as within the replication cohort #1 using the DerSimian-Laird approach (DerSimonian, R. and N. Laird. 1986. Meta-analysis in clinical trials. Control Clin Trials 7: 177-188.)

[0227] The absence of a statistically significant association in one or more of the replication cohorts should not be interpreted as minimizing the value of the original finding. There are many reasons why a biologically derived association identified in a sample from one population would not replicate in a sample from another population. The most important reason is differences in population history. Due to bottlenecks and founder effects, there may be common disease predisposing alleles present in one population that are relatively rare in another, leading to a lack of association in the candidate region. Also, because common diseases such as arthritis-related disorders are the result of susceptibilities in many genes and many environmental risk factors, differences in population-specific genetic and environmental backgrounds could mask the effects of a biologically relevant allele. For these and other reasons, statistically strong results in the original, discovery sample that did not replicate in one or more of the replication samples may be further evaluated in additional replication cohorts and experimental systems.

# Example 4

### IL1RL1 Region Proximal SNPs

[0228] It has been discovered that SNP rs1041973 in Interleukin 1 receptor-like 1 isoform 1 (*IL1RL1*) is associated with occurrence of osteoarthritis in subjects. Interleukin 1 receptor-like 1 isoform 1 is a member of the interleukin 1 receptor family with no known ligand (orphan receptor). IL1RL1 exists in both a soluble and transmembrane form, suggesting that it may have ligand or ligand scavenging activity. Studies of the similar gene in mouse suggested that this receptor can be induced by proinflammatory stimuli. This gene and four other interleukin 1 receptor family genes, including interleukin 1 receptor, type I (IL1R1), interleukin 1 receptor, type II (IL1R2), interleukin 1 receptor-like 2 (IL1RL2), and interleukin 18 receptor 1 (IL18R1), form a cytokine receptor gene cluster.

[0229] Ninety-one additional allelic variants proximal to rs1041973 were identified and subsequently allelotyped in osteoarthritis case and control sample sets as described in Examples 1 and 2.

The polymorphic variants are set forth in Table 10. The chromosome positions provided in column four of Table 10 are based on Genome "Build 34" of NCBI's GenBank.

TABLE 10

dbSNP Position in SEO Chromosome Allele					
dbSNP rs#	Chromosome	ID NO: 1	Chromosome Position	Allele Variants	
rs884517	2	207	102527857	c/t	
rs1476984	2	6019	102533669	a/g	
rs951774	2	6414	102534064	a/c	
rs2041737	2	7341	102534991	a/g	
rs1420091	2	10984	102538634	a/g	
rs2110660	2	12351	102540001	c/g	
rs1362347	2	13335	102540985	a/g	
rs3073968	2	16584	102544234	-/tgtg/tgtgag	
rs4090473	2	16737	102544387	c/g	
rs1558622	2	23897	102551547	c/t	
rs1558621	2	24057	102551707	c/t	
rs1558620	2	25145	102552795	a/g	
rs1558619	2	25300	102552950	a/c	
rs950881	2	26262	102553912	a/c	
rs950880	2	26312	102553962	g/t	
rs1362346	2	26589	102554239	c/t	
rs1968171	2	27302	102554952	a/g	
rs1813299	2	27358	102555008	a/t	
rs1813298	2	27451	102555101	c/g	
rs1968170	2	27552	102555202	c/t	
rs974389	2	30731	102558381	c/t	
rs971764	2	32085	102559735	a/g	
rs1420089	2	32139	102559789	a/g	
rs1420088	2	33184	102560834	a/g	
rs1420103	2	42382	102570032	g/t	
rs1420102	2	42569	102570219	a/g	
rs1997467	2	44823	102572473	c/t	
rs1997466	2	45217	102572867	c/g	
rs1362350	2	45548	102573198	c/g	
rs2310220	2	45601	102573251	a/g	
rs1362349	2	45722	102573372	c/g	
rs3755278	2	45967	102573617	a/g	
rs3771180	2	47367	102575017	a/c	
rs3771179	2	47642	102575292	a/c	
rs985523	2	48126	102575776	c/t	
rs1041973	2	49218	102576868	a/c	
rs3214363	2	49274	102576924	-/a	
rs873022	2	49433	102577083	g/t	
rs3771177	2	49610	102577260	a/c	
rs3732129	2	51282	102578932	a/g	
rs1420101	2	51466	102579116	a/g	
rs12905	2	53757	102581407	a/g	

dbSNP rs#	Chromosome	Position in SEQ ID NO: 1	Chromosome Position	Allele Variants
rs3771175	2	53960	102581610	a/t
rs3821204	2	54031	102581681	c/g
rs2160203	2	54574	102582224	c/t
rs1946131	2	55679	102583329	a/g
rs1054096	2	56100	102583750	c/t
rs2287038	2	56182	102583832	c/t
rs1921622	2	59817	102587467	a/g
rs1861246	2	60533	102588183	a/g
rs1861245	2	60656	102588306	a/g
rs3755276	2	72209	102599859	a/g
rs2287037	2	72778	102600428	a/g
rs1420099	2	74293	102601943	c/g
rs3771174	2	77335	102604985	a/g
rs1420098	2	78029	102605679	a/g
rs1362348	2	78374	102606024	c/g
rs1882348	2	78421	102606071	a/t
rs1558627	2	78434	102606084	c/t
rs2058622	2	79174	102606824	c/t
rs3836110	2	79397	102607047	-/g
rs3771172	2	79562	102607212	a/g
rs3771171	2	79700	102607350	a/g
rs3771170	2	79730	102607380	a/t
rs2160202	2	79904	102607554	c/t
rs2058623	2	79920	102607570	a/g
rs3771167	2	79938	102607588	c/t
rs3771166	2	79972	102607622	c/t
rs1974675	2	80125	102607775	c/t
rs1465321	2	80368	102608018	a/g
rs2041740	2	83484	102611134	c/t
rs3771164	2	85536	102613186	a/t
rs2270298	2	85829	102613479	c/t
rs2270297	2	86425	102614075	a/g
rs2041739	2	88083	102615733	a/g
rs2080289	2	88770	102616420	c/t
rs3821203	2	90622	102618272	a/g
rs3771162	2	90924	102618574	a/t
rs3213733	2	91634	102619284	g/t
rs3213732	2	92029	102619679	c/t
rs1035130	2	95152	102622802	a/g
rs3752659	2	95348	102622998	c/t
rs3755274	2	96145	102623795	c/t
rs2241117	2	96793	102624443	a/g
rs2241116	2	97015	102624665	g/t
rs881890	2	97064	102624714	c/t
rs3771161	2	97711	102625361	g/t
rs3771160	2	97855	102625505	a/c
, , , , , , , , , , , , , , , , , , , ,		, 0.000		u

dbSNP rs#	Chromosome	Position in SEQ ID NO: 1	Chromosome Position	Allele Variants
rs1420104	2	not mapped	not mapped	c/t
rs2041738	2	not mapped	not mapped	a/c

# Assay for Verifying and Allelotyping SNPs

[0230] The methods used to verify and allelotype the 91 proximal SNPs of Table 10 are the same methods described in Examples 1 and 2 herein. The primers and probes used in these assays are provided in Table 11 and Table 12, respectively.

TABLE 11

dbSNP rs#	Forward PCR primer	Reverse PCR primer
rs884517	ACGTTGGATGCATTTTCTGGTGTGACTCCC	ACGTTGGATGATGTTCGGTCACTTGTGAGC
rs1476984	ACGTTGGATGTGAGAGAGTTGAAGAATGGG	ACGTTGGATGCCAAGAAGTGATTTCCTTCC
rs951774	ACGTTGGATGTCAGCCAGAGGTCTTTACTC	ACGTTGGATGTTAGAAGTCTCTTGGGTGGG
rs2041737	ACGTTGGATGGAGATGGAGTTTCCCTCTTG	ACGTTGGATGAAACCAAGAGGTGGAGGTTG
rs1420091	ACGTTGGATGCACCCCTATTATAAAACCCAC	ACGTTGGATGACCAGAAATGGCATCTATGG
rs2110660	ACGTTGGATGTCTCCCGAGATGAGGAATC	ACGTTGGATGGTGATCTCCTCAGTACTCTG
rs1362347	ACGTTGGATGTTCTTTGGTAATGAGGTAGG	ACGTTGGATGTGCTTGCCCTCTATTTATGG
rs3073968	ACGTTGGATGGAATGATGAGGAAGGAAGGG	ACGTTGGATGTAAAGCCACATGTTCACCCG
rs4090473	ACGTTGGATGTAGTGTTTCACTCTTCCC	ACGTTGGATGTCAAGCACCTCTGTTAACTC
rs1558622	ACGTTGGATGATACTTCCTGGTTTTCTGGG	ACGTTGGATGGGCTCAAAGTCATCACCCAA
rs1558621	ACGTTGGATGACAGTGGCGATGCCAACATT	ACGTTGGATGCCTGTAGTAGGACCCTACTG
rs1558620	ACGTTGGATGTTGCAGGTGTCTGGTGATAG	ACGTTGGATGAGTTTGCCTTTCTTCATGGC
rs1558619	ACGTTGGATGCCCTAATTAGGATTCCGCAC	ACGTTGGATGCTCCATCACACTTTGACTGC
rs950881	ACGTTGGATGCTTATCTCAGTCTGCCAGTG	ACGTTGGATGGGTGAGTGAATTAGTCCTGG
rs950880	ACGTTGGATGTGCCAAAGACAATCAAATCC	ACGTTGGATGCACTCACCTCTGATTTCTAG
rs1362346	ACGTTGGATGTTCCTCAGGTTCACCAAGAG	ACGTTGGATGTCCCGAACCTCATCTCATAC
rs1968171	ACGTTGGATGAATGTTTCAGCCCAGCATGG	ACGTTGGATGATCTCCTGACCTCATGATCC
rs1813299	ACGTTGGATGAATTCCAGCACTTTGGGAGG	ACGTTGGATGTTTCACCGTGTTAGCCAGGA
rs1813298	ACGTTGGATGTTACTGCAAGCTCCACCTCC	ACGTTGGATGATTAACTGGGCGTGGTGGTG
rs1968170	ACGTTGGATGAGCTTGCAGTAAGCCCAGAT	ACGTTGGATGTGTTAGGGTAATTACAGTGC
rs974389	ACGTTGGATGCTCTAGCCCAATATGTCTCC	ACGTTGGATGACTGGAGATGTGAACCCATC
rs971764	ACGTTGGATGGAGATTAAGAGG	ACGTTGGATGAGTTGTTTGACTTCGGACTG
rs1420089	ACGTTGGATGAGACAGCACATATCAATGAC	ACGTTGGATGTATTGTGCGGTTCGCTATAG
rs1420088	ACGTTGGATGGGATGACTGTCAAAAACATC	ACGTTGGATGTAATTTTTCAGGAGCAAGGC
rs1420103	ACGTTGGATGTCCATTGGAATATGACCTCC	ACGTTGGATGCCAGGCACATGAGCTATATC
rs1420102	ACGTTGGATGGATTGGTCAGGAACTCAAAC	ACGTTGGATGTGGGTTGCTTCTAGCTATTG
rs1997467	ACGTTGGATGTGAATTTCAGTGAGTCAGGC	ACGTTGGATGTGAGGGGAAAAAAAAACATCC
rs1997466	ACGTTGGATGATAGGCACATACAGGATTTC	ACGTTGGATGCTCCCTTTTCAGATTAATCTC
rs1362350	ACGTTGGATGGAGAACATTCTCTATACCAG	ACGTTGGATGTGCCTGAATAGTGAGAAGCC
rs2310220	ACGTTGGATGGGTTGAAACCAGACTTGCTG	ACGTTGGATGCAGCCTAATCTCTGGTATAG
rs1362349	ACGTTGGATGCAATACTCTGTGGTACTTATC	ACGTTGGATGTAAACAGTCTTATCCTTGGG
rs3755278	ACGTTGGATGAGTGCTGAATAGGTTTGTTC	ACGTTGGATGGCCTAGTTTAAGAATGAATGC

dbSNP rs#	Forward PCR primer	Reverse PCR primer
rs3771180	ACGTTGGATGGTCAACATCAAGAATTCTTAG	ACGTTGGATGCCTGAAATTTGATTTGTGGC
rs3771179	ACGTTGGATGGTCTTCATAATTCATGATTG	ACGTTGGATGTCTTAAAATATAAGGGGAAG
rs985523	ACGTTGGATGTCCTATGGAAGTTTTGGGTC	ACGTTGGATGCTGCGAAGTAGCATGATAAC
rs1041973	ACGTTGGATGGGGACTTCTGACAATACAGG	ACGTTGGATGAATCGTGTGTTTGCCTCAGG
rs3214363	ACGTTGGATGCAGGCAATCAACCACTGAAG	ACGTTGGATGCTGCAGTTGCTGATTCTGGT
rs873022	ACGTTGGATGCCTAGTCCTTTCTGGAACAG	ACGTTGGATGATCCCTGCAACTGTAAATCC
rs3771177	ACGTTGGATGAAGGTTAGAAGCCCCTTTTC	ACGTTGGATGGGCTGGAATTAAGAACAAAC
rs3732129	ACGTTGGATGCTAATTCAAAGCCACATCTG	ACGTTGGATGTAAGTTAGCATTCAGATTGC
rs1420101	ACGTTGGATGCAACATTTATGTACACCATAG	ACGTTGGATGTTAGTAATACTCATTGGATT
rs12905	ACGTTGGATGCTCCCAGCAAACAGGAACAG	ACGTTGGATGATCAAGACAATGGGAATGGC
rs3771175	ACGTTGGATGAAAGAGCACAAAAGAACACG	ACGTTGGATGTTATGAACTCCCTCTGTGTC
rs3821204	ACGTTGGATGCATGTTGTAAGCATGGTCCG	ACGTTGGATGACTTTACCACCCTCGCTAAC
rs2160203	ACGTTGGATGACACAGACCCAAACCATACC	ACGTTGGATGTTCCCGTGTGTTCCATGTAC
rs1946131	ACGTTGGATGGGGAACTCAGGGTTTAACAC	ACGTTGGATGTACACTCATCACTCCTCAGG
rs1054096	ACGTTGGATGATCAAGGTGCTATGTGAGGG	ACGTTGGATGAAAGCAGGAGTACACAAGGG
rs2287038	ACGTTGGATGAATGTCCCTGGTTACCTATG	ACGTTGGATGACAAATAAGCTAGAAGGAGC
rs1921622	ACGTTGGATGGCCACTTCTTAATTCTGTCC	ACGTTGGATGATTTCAGCTAGTGCCTATGG
rs1861246	ACGTTGGATGCACAAGCTCTTCACCTCTTC	ACGTTGGATGTGGCTGAGGAGAGTGTAAC
rs1861245	ACGTTGGATGTGCTGCCTTCAATGTGTGAC	ACGTTGGATGAGGAAAGGTCAGAGGACATG
rs3755276	ACGTTGGATGCCAGCACTCACTAACATGTG	ACGTTGGATGAAACTCATATGGGCAGCCAC
rs2287037	ACGTTGGATGCAGATTCAGCCAAAGCTTTC	ACGTTGGATGAAAAATCTGTGTGCCAGAAG
rs1420099	ACGTTGGATGTTACACACTCTCCAGAGGTG	ACGTTGGATGAAAGCTTCTAGCTGCCTGAG
rs3771174	ACGTTGGATGACCCAGATTCTCTGGCTTTG	ACGTTGGATGTACCACAAGTGCCGAAAGAG
rs1420098	ACGTTGGATGGGGACGTGAAGTACAAGAT	ACGTTGGATGGGAGACCAAAAAAAGTTACC
rs1362348	ACGTTGGATGCATGTCATAGGAAGAGTAGG	ACGTTGGATGTCAGCAACTCAAATATGCAG
rs1882348	ACGTTGGATGCCTACTCTTCCTATGACATG	ACGTTGGATGCCCTAAAAGGAAATCCTATC
rs1558627	ACGTTGGATGCCCTAAAAGGAAATCCTATC	ACGTTGGATGCCTACTCTTCCTATGACATG
rs2058622	ACGTTGGATGCTGTGAAACCTTGGTAGCAC	ACGTTGGATGTTTCTGATGCCTGGGAGTTC
rs3836110	ACGTTGGATGACTCACAAATGGGGTAAAGG	ACGTTGGATGTGCCTTCATTCAATCAGGAG
rs3771172	ACGTTGGATGCAGAAGCAAATGGCATTGGC	ACGTTGGATGCCATTGTTGCTTCCTAAGCC
rs3771171	ACGTTGGATGAGGGTAGCAGATAGGAGATG	ACGTTGGATGAAGCTGCTTCTCCTCATC
rs3771170	ACGTTGGATGCAAGGCCATTGTCAAAGCTG	ACGTTGGATGGTGTCCCAGAGTGGATATTG
rs2160202	ACGTTGGATGAGCAGTATTTACTGCAGATG	ACGTTGGATGCCCACATCAAACTGCAAAGG
rs2058623	ACGTTGGATGATTTACTGCAGATGTGTGTG	ACGTTGGATGTGTTCACTGATAGATCCCAC
rs3771167	ACGTTGGATGCTAACTTAAGTGTGTAACCC	ACGTTGGATGCTAACGGGAAATTTTCAGGTG
rs3771166	ACGTTGGATGGTGAACAGACTTTACACCTG	ACGTTGGATGCCTCAGTGGCATTTGATTAT
rs1974675	ACGTTGGATGACTAAGAAGGAAGGGGATAC	ACGTTGGATGGTACATTTCCCTCTACCTTC
rs1465321	ACGTTGGATGTCACAGCTTTGGGTCAGTTG	ACGTTGGATGTCAACAACACACTGCACCTG
rs2041740	ACGTTGGATGCATCCATGTCCCTACAAAAG	ACGTTGGATGAAAGCTCTTATACACCATGG
rs3771164	ACGTTGGATGCCTGTGACATGTATGGAAATG	ACGTTGGATGTCAAATCCATAGGTACACTC
rs2270298	ACGTTGGATGTGAAGTAGTGTTCTCTCTC	ACGTTGGATGAATATGAGCACTGTAGCTGC
rs2270297	ACGTTGGATGTTTCCTGCCAAAAAGAAAGG	ACGTTGGATGGACCACCACTAGTTCAAA
rs2041739	ACGTTGGATGTAGACCCTGAAGTTTCCCAC	ACGTTGGATGCACCTAGAGGTTCCTTTTGC
rs2080289	ACGTTGGATGTGGAGAATGTCAACTGAGTC	ACGTTGGATGATACAAACAAGAGGCCATGG
rs3821203		ACGTTGGATGGGATCCAGAGAAAGGTAGTC
rs3771162	ACGTTGGATGTGAGTGAGAC	ACGTTGGATGTGGCACTTTCTGAGA
rs3213733	ACGTTGGATGTGAAAGCACCTTGTATCTGG	ACGTTGGATGCATCTTCCTCTGCCTTTTAG

dbSNP rs#	Forward PCR primer	Reverse PCR primer
rs3213732	ACGTTGGATGGTCAGGTTAAAAGTGGCAAC	ACGTTGGATGTGACACTGGATACACATTTC
rs1035130	ACGTTGGATGTTAGGATCCGATCCATTTTC	ACGTTGGATGCTCTGCTTTGCTGAATGAAG
rs3752659	ACGTTGGATGTGCATAATGCGTCCACCTAG	ACGTTGGATGGGCTGATGTGTATTTTGGGC
rs3755274	ACGTTGGATGTATCAAAGGTGTGTGCACCC	ACGTTGGATGAGGGGTAGAAAACCACAGTG
rs2241117	ACGTTGGATGTGGCTGGAAGATCATGATGC	ACGTTGGATGCCCCAAGTTGTTAGGAAGAG
rs2241116	ACGTTGGATGAATGCAGGCAACATCACAGC	ACGTTGGATGAGTAGGCTCTGTTCGTTACC
rs881890	ACGTTGGATGATGCCATTTGCCTTCTGGAG	ACGTTGGATGTCTCAGGGTAACGAACAGAG
rs3771161	ACGTTGGATGCCATCAGGTGAGCACTGAAA	ACGTTGGATGTCATTGCCTCCTGAACTTGG
rs3771160	ACGTTGGATGAGAAATGGCTGTGACTGGAG	ACGTTGGATGTATCCAGGGAGTTGATGGTG
rs3771159	ACGTTGGATGCAGGTGATGGTCCAACAAAG	ACGTTGGATGTGCTGTGGTCCACTCACTTG
rs1420104	ACGTTGGATGTATTCTGGAGGCTGAGGTGG	ACGTTGGATGTGGAGTGCAGTGGTGATC
rs2041738	ACGTTGGATGTGGTGAAACCCCATCTCTAC	ACGTTGGATGTTTCAAGCTATTCTCCTGCC

**TABLE 12** 

dbSNP rs#	Extend Primer	Term Mix
rs884517	GGTGTGACTCCCAGACCAA	ACT
rs1476984	ATGGGTAGTTAATGGTGGAAATTT	ACT
rs951774	CAAAGTAGTTGACTTGTCTTTCT	ACT
rs2041737	CCAGGCTAGTGCAGTGGC	ACT
rs1420091	CCCACATTATATTGTCATTACTTT	ACG
rs2110660	ATGAGGAATCAGAGCTGGGA	ACT
rs1362347	GTAATGAGGTAGGAATAATATTG	ACT
rs3073968	GGCAATTGTGTGTGTGTG	CGT
rs4090473	CTTACTCCTATTCCAAAGTTCA	ACT
rs1558622	ACTGCAAGGGAGAGCCCC	ACT
rs1558621	AGTGTGTGTGTGCGTGC	ACT
rs1558620	GTCTGGTGATAGTTGGGTGC	ACG
rs1558619	GATTCCGCACATCCTATGCCT	ACT
rs950881	GATGGTTTGTGCCTCTGGTC	ACT
rs950880	ATTTAAGAATGCTTTCGTCATAAG	ACT
rs1362346	GAATATCTATGCCCACCAGAT	ACG
rs1968171	GCCCAGCATGGTGGCTCA	ACG
rs1813299	GTGGATCATGAGGTCAGGAG	CGT
rs1813298	GCCTCAGCCTCCCGAGTA	ACT
rs1968170	AGCCTGGGTGACAGAGCC	ACT
rs974389	GTCTCCTGAATTTCAGAAGCA	ACT
rs971764	GTCAAGGTAAAAACATTATTGTG	ACG
rs1420089	GCACATATCAATGACAAGACTA	ACT
rs1420088	CATGTTATGTAACTCTGAGTTC	ACT
rs1420103	GAATATGACCTCCAGAAGGCAA	ACT
rs1420102	GAACTCAAACAAATACTTGGACAC	ACG
rs1997467	TTCAGTGACTCTCACAATAAGC	ACG
rs1997466	AAGAAAAAGCTGGTTCAATGAG	ACT
rs1362350	ACATTCTCTATACCAGAGATTAGG	ACT

dbSNP rs#	Extend Primer	Term Mix
rs2310220	CTGAACTTCAAAGTCAAGCTTTT	ACG
rs1362349	CTGTGGTACTTATCATTAACATCA	ACT
rs3755278	ACTCGGAATTCTTTTACATTTGGT	ACT
rs3771180	CATCAAGAATTCTTAGTACATGAT	ACT
rs3771179	TATGTTAGTAAATTTCTATGTTGG	ACT
rs985523	CATATAGCTTTCACAATGATCATG	ACG
rs1041973	ATACCAGAATCAGCAACT	ACT
rs3214363	GAGCAGGGTGAAAGAAGATGGG	ACT
rs873022	TTCTAGGAATACTATCAGGTTGA	ACT
rs3771177	TTTTCACCTACTAGAGGCCC	CGT
rs3732129	GCCACATCTGTTCTTTATTCTTT	ACG
rs1420101	CCATCACAAAGCCTCTCATTA	ACT
rs12905	AGACAGCAAACAACATCC	ACG
rs3771175	CACAAAAGAACACGTTCAGTTT	CGT
rs3821204	TAAGCATGGTCCGTTCTATAC	ACT
rs2160203	CCACACACATTATCATTGTTA	ACT
rs1946131	TTAACACTCTTTGGCTATTTGACA	ACT
rs1054096	TCCATCCAGCCTGCCCAC	ACG
rs2287038	TACCTATGTGTTTGAATTATCTTC	ACT
rs1921622	GAAAGAGGACTTAAAAATTGATGA	ACT
rs1861246	CTTCACCTCTTCTTTTCAGTC	ACG
rs1861245	CTGGAATGGTTTTCTACTTCC	ACG
rs3755276	GTGTGTATGCATGTGTTCGC	ACT
rs2287037	ACAAAAGTGTGCCTATCTTATGAA	ACT
rs1420099	GGTGGGAGGTTGATAATTGAAA	ACT
rs3771174	CTGACCATCATCTACCCAGG	ACT
rs1420098	ACGTGAAGTACAAGATTCTTCA	ACT
rs1362348	GAGTAGGAAAGAAAAGGATGTG	ACT
rs1882348	TCCTATGACATGAAATACATTCT	CGT
rs1558627	AAGCAGAGAGAGATAAACTTATT	ACG
rs2058622	AAACCTTGGTAGCACTTCTGT	ACT
rs3836110	AACAAACACCGCCCCCC	CGT
rs3771172	GCATTGGCCATCTTTCTGATA	ACG
rs3771171	GAGGTGTCCCAGAGTGGATA	ACG
rs3771170	CAAAGCTGCTTCTCCTCA	CGT
rs2160202	TATACACATATGTGTTCTAACTTA	ACT
rs2058623	ACTTAGGTGTGTAACCCTTTG	ACG
rs3771167	CTTTGTAGTTTGATGTGGGATCT	ACT
rs3771166	ACTTTACACCTGAAAATTTCCC	ACT
rs1974675	GAAGGGATACAAAAGGGATA	ACT
rs1465321	CAGTTGGCCTCAGTGTTAACCC	ACG
rs2041740	GAACTCATGCTTTTTTATGGCTG	ACG
rs3771164	GACATGTATGGAAATGTGTGTG	CGT
rs2270298	CTCTCTCTGCATGTGTGT	ACT

dbSNP rs#	Extend Primer	Term Mix
rs2270297	AGCCAAGTAGAGGAGCACC	ACT
rs2041739	CTCCTGAGTTCCTGTGAATAC	ACT
rs2080289	TCTCAGGACTCCACTCAAATGTC	ACT
rs3821203	GGCAGGAGGCAATTTCGGT	ACT
rs3771162	CAGTGAGACTCAGGAGTGC	CGT
rs3213733	TGTATCTGGTTTTCTCTCACTCA	ACT
rs3213732	CAACATTCAAAAAATGGCACTCTT	ACG
rs1035130	TCCGATCCATTTTCTTCCCC	ACT
rs3752659	CCTAGGGTATGGCCACTATAATTA	ACG
rs3755274	CACCCAACTATAAAGAAAGACCTC	ACG
rs2241117	ATCATGATGCTAAGTTGAAAATAT	ACT
rs2241116	TCAAGCATTTTAAACATGTGAATT	CGT
rs881890	TGCCTTCTGGAGTCCTGTAA	ACT
rs3771161	GTGAGCACTGAAAAACTTTAAGA	ACT
rs3771160	GCCAGAAAGCTGTGATTTCCA	ACT
rs3771159	CCAACAAAGATTTGAGCCCC	ACT
rs1420104	CTGGGAGGTGGAGACTGCA	ACT
rs2041738	AAAAATACAAAAATTAGCTGGGC	ACT

### Genetic Analysis

[0231] Allelotyping results from the discovery cohort are shown for cases and controls in Table 13. The allele frequency for the A2 allele is noted in the fifth and sixth columns for osteoarthritis case pools and control pools, respectively, where "AF" is allele frequency. The allele frequency for the A1 allele can be easily calculated by subtracting the A2 allele frequency from 1 (A1 AF = 1-A2 AF). For example, the SNP rs951774 has the following case and control allele frequencies: case A1 (A) = 0.24; case A2 (C) = 0.76; control A1 (A) = 0.20; and control A2 (C) = 0.80, where the nucleotide is provided in paranthesis. Some SNPs are labeled "untyped" because of failed assays.

**TABLE 13** 

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs884517	207	102527857	C/T			
rs1476984	6019	102533669	A/G	0.83	0.83	0.973
rs951774	6414	102534064	A/C	0.76	0.80	0.099
rs2041737	7341	102534991	A/G	0.38	0.32	0.146
rs1420091	10984	102538634	A/G	0.33	0.35	0.388
rs2110660	12351	102540001	C/G	0.41	0.40	0.753
rs1362347	13335	102540985	A/G	0.83	0.83	0.895
rs3073968	16584	102544234	- /TGTG/T GTGAG	0.48	0.48	0.878
rs4090473	16737	102544387	C/G	0.42	0.43	0.633
rs1558622	23897	102551547	С/Т	0.40	0.39	0.879

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs1558621	24057	102551707	C/T	0.32	0.31	0.795
rs1558620	25145	102552795	A/G	0.37	0.37	0.998
rs1558619	25300	102552950	A/C	0.46	0.47	0.556
rs950881	26262	102553912	A/C	0.75	0.74	0.636
rs950880	26312	102553962	G/T	0.45	0.48	0.285
rs1362346	26589	102554239	C/T			
rs1968171	27302	102554952	A/G	0.43	0.43	0.891
rs1813299	27358	102555008	A/T			
rs1813298	27451	102555101	C/G			
rs1968170	27552	102555202	С/Т	0.65	0.65	0.941
rs974389	30731	102558381	C/T	0.41	0.42	0.734
rs971764	32085	102559735	A/G	0.45	0.44	0.738
rs1420089	32139	102559789	A/G	0.16	0.19	0.099
rs1420088	33184	102560834	A/G	0.41	0.40	0.869
rs1420103	42382	102570032	G/T	0.68	0.68	0.952
rs1420102	42569	102570219	A/G	0.48	0.46	0.349
rs1997467	44823	102572473	C/T			
rs1997466	45217	102572867	C/G	0.46	0.46	0.693
rs1362350	45548	102573198	C/G	0.48	0.46	0.475
rs2310220	45601	102573251	A/G	0.40	0.41	0.480
rs1362349	45722	102573372	C/G	0.41	0.42	0.893
rs3755278	45967	102573617	A/G	0.07	0.08	0.876
rs3771180	47367	102575017	A/C	0.91	0.90	0.669
rs3771179	47642	102575292	A/C	0.08	0.08	0.986
rs985523	48126	102575776	C/T	0.17	0.13	0.064
rs1041973	49218	102576868	A/C	- 0.,,	0.10	0.001
rs3214363	49274	102576924	-/A	1		
rs873022	49433	102577083	G/T	0.53	0.56	0.321
rs3771177	49610	102577260	A/C	0.33	0.31	0.278
rs3732129	51282	102578932	A/G	0.46	0.50	0.127
rs1420101	51466	102579116	A/G	0.55	0.57	0.257
rs12905	53757	102581407	A/G	0.30	0.27	0.262
rs3771175	53960	102581610	A/T	0.84	0.82	0.174
rs3821204	54031	102581681	C/G	0.26	0.23	0.222
rs2160203	54574	102582224	C/T	0.21	0.26	0.033
rs1946131	55679	102583329	A/G	0.73	0.74	0.710
rs1054096	56100	102583750	C/T	0.69	0.65	0.137
rs2287038	56182	102583832	C/T	0.98	0.95	0.207
rs1921622	59817	102587467	A/G	0.40	0.43	0.218
rs1861246	60533	102588183	A/G	0.22	0.18	0.068
rs1861245	60656	102588306	A/G	0.35	0.37	0.377
rs3755276	72209	102599859	A/G	0.51	0.48	0.355
rs2287037	72778	102600428	A/G	0.49	0.53	0.195
rs1420099	74293	102601943	C/G	0.58	0.56	0.416
rs3771174	77335	102604985	A/G	1	3.00	
rs1420098	78029	102605679	A/G	0.33	0.32	0.532
rs1362348	78374	102606024	C/G	0.02	0.03	0.590
rs1882348	78421	102606071	A/T	0.36	0.35	0.596
rs1558627	78434	102606084	С/Т	0.62	0.65	0.219
rs2058622	79174	102606824	С/Т	0.57	0.59	0.528
rs3836110	79397	102607047	-/G	0.72	0.73	0.856
rs3771172	79562	102607212	Ā/G	0.28	0.25	0.261
rs3771171	79700	102607212	A/G	J	0.20	3,231
rs3771170	79730	102607380	A/T	0.24	0.23	0.533
rs2160202	79904	102607554	C/T	0.55	0.62	0.061
rs2058623	79920	102607570	A/G	0.67	0.68	0.631
	1 0020	102001310	_ ~0	1 0.07	1 0.00	0.001

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs3771166	79972	102607622	С/Т	0.55	0.53	0.624
rs1974675	80125	102607775	С/Т	0.57	0.55	0.470
rs1465321	80368	102608018	A/G	0.27	0.26	0.614
rs2041740	83484	102611134	С/Т	0.26	0.25	0.622
rs3771164	85536	102613186	A/T	0.76	0.73	0.197
rs2270298	85829	102613479	C/T	0.23	0.21	0.329
rs2270297	86425	102614075	A/G	0.60	0.60	0.900
rs2041739	88083	102615733	A/G	0.43	0.40	0.235
rs2080289	88770	102616420	C/T	0.56	0.59	0.322
rs3821203	90622	102618272	A/G	0.58	0.62	0.194
rs3771162	90924	102618574	A/T	0.30	0.28	0.260
rs3213733	91634	102619284	G/T	0.76	0.73	0.287
rs3213732	92029	102619679	С/Т	0.44	0.42	0.507
rs1035130	95152	102622802	A/G	0.58	0.61	0.234
rs3752659	95348	102622998	С/Т	0.80	0.80	0.957
rs3755274	96145	102623795	C/T	0.26	0.25	0.549
rs2241117	96793	102624443	A/G	0.71	0.75	0.077
rs2241116	97015	102624665	G/T	0.16	0.15	0.469
rs881890	97064	102624714	С/Т			
rs3771161	97711	102625361	G/T	0.70	0.68	0.348
rs3771160	97855	102625505	A/C			
rs3771159	98708	102626358	A/G	0.38	0.40	0.294
rs1420104	not mapped	not mapped	C/T			
rs2041738	not mapped	not mapped	A/C	<u>l</u>		

[0232] The *IL1RL1* proximal SNPs were also allelotyped in the replication cohorts using the methods described herein and the primers provided in Tables 11 and 12. The replication allelotyping results for replication cohort #1 and replication cohort #2 are provided in Tables 14 and 15, respectively.

**TABLE 14** 

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs884517	207	102527857	C/T			
rs1476984	6019	102533669	A/G	0.82	0.81	0.878
rs951774	6414	102534064	A/C	0.76	0.82	0.024
rs2041737	7341	102534991	A/G	0.36	0.32	0.382
rs1420091	10984	102538634	A/G	0.32	0.35	0.340
rs2110660	12351	102540001	C/G	0.39	0.39	0.951
rs1362347	13335	102540985	A/G	0.83	0.83	0.766
			- /TGTG/T			
rs3073968	16584	102544234	GTGAG	0.47	0.48	0.822
rs4090473	16737	102544387	C/G	0.40	0.41	0.663
rs1558622	_23897	102551547	C/T_	0.38	0.38	0.943
rs1558621	24057	102551707	С/Т	0.33	0.32	0.631
rs1558620	25145	102552795	A/G_	0.34	0.34	0.957
rs1558619	25300	102552950	A/C	0.44	0.47	0.368
rs950881	26262	102553912	A/C	0.76	0.74	0.476
rs950880	26312	102553962	G/T	0.42	0.47	0.199
rs1362346	26589	102554239	C/T			
rs1968171	27302	102554952	A/G	0.43	0.44	0.641
rs1813299	27358	102555008	A/T			

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs1813298	27451	102555101	C/G	<del> </del>		
rs1968170	27552	102555202	C/T	0.64	0.65	0.797
rs974389	30731	102558381	C/T	0.39	0.41	0.678
rs971764	32085	102559735	A/G	0.47	0.46	0.710
rs1420089	32139	102559789	A/G	0.16	0.21	0.075
rs1420088	33184	102560834	A/G	0.41	0.40	0.869
rs1420103	42382	102570032	G/T	0.69	0.72	0.268
rs1420102	42569	102570219	A/G	0.50	0.47	0.329
rs1997467	44823	102572473	C/T			
rs1997466	45217	102572867	C/G	0.49	0.47	0.675
rs1362350	45548	102573198	C/G	0.51	0.47	0.308
rs2310220	45601	102573251	A/G	0.40	0.44	0.282
rs1362349	45722	102573372	C/G	0.42	0.43	0.730
rs3755278	45967	102573617	A/G	0.08	0.08	0.902
rs3771180	47367	102575017	A/C	0.93	0.92	0.591
rs3771179	47642	102575292	A/C	0.08	0.08	0.936
rs985523	48126	102575776	C/T	0.17	0.13	0.156
rs1041973	49218	102576868	A/C			
rs3214363	49274	102576924	-/A			
rs873022	49433	102577083	G/T	0.51	0.56	0.138
rs3771177	49610	102577260	A/C	0.36	0.32	0.125
rs3732129	51282	102578932	A/G	0.43	0.50	0.048
rs1420101	51466	102579116	A/G	0.50	0.55	0.132
rs12905	53757	102581407	A/G	0.33	0.28	0.127
rs3771175	53960	102581610	A/T	0.86	0.83	0.217
rs3821204	54031	102581681	C/G	0.29	0.23	0.071
rs2160203 rs1946131	54574 55679	102582224	C/T A/G	0.19 0.72	0.26	0.016
rs1054096		102583329	C/T	0.72	0.73 0.65	0.771
rs2287038	56100 56182	102583750 102583832	C/T	0.70	NA	0.079 0.975
rs1921622	59817	102583832	A/G	0.93	0.41	0.973
rs1861246	60533	102588183	A/G	0.37	0.15	0.031
rs1861245	60656	102588306	A/G	0.34	0.39	0.149
rs3755276	72209	102599859	A/G	0.53	0.46	0.072
rs2287037	72778	102600428	A/G	0.45	0.51	0.069
rs1420099	74293	102601943	C/G	0.59	0.55	0.312
rs3771174	77335	102604985	A/G		0.00	0.0
rs1420098	78029	102605679	A/G	0.35	0.32	0.328
rs1362348	78374	102606024	C/G	0.02	NA	0.025
rs1882348	78421	102606071	A/T	0.40	0.37	0.399
rs1558627	78434	102606084	C/T	0.64	0.69	0.118
rs2058622	79174	102606824	C/T	0.59	0.62	0.491
rs3836110	79397	102607047	-/G	0.74	0.75	0.625
rs3771172	79562	102607212	A/G	0.31	0.27	0.200
rs3771171	79700	102607350	A/G			
rs3771170	79730	102607380	A/T_	0.22	0.20	0.346
rs2160202	79904	102607554	C/T	0.55	0.60	0.217
rs2058623	79920	102607570	A/G	0.69	0.72	0.266
rs3771167	79938	102607588	C/T			
rs3771166	79972	102607622	С/Т	0.57	untyped	NA
rs1974675	80125	102607775	С/Т	0.58	0.54	0.297
rs1465321	80368	102608018	A/G	0.25	0.23	0.471
rs2041740	83484	102611134	C/T	0.25	0.22	0.450
rs3771164	85536	102613186	A/T	0.77	0.72	0.073
rs2270298	85829	102613479	C/T	0.25	0.22	0.324
rs2270297	86425	102614075	A/G	0.63	0.64	0.589

dbSNP rs#	Position in SEQ ID NO: 1	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs2080289	88770	102616420	С/Т	0.53	0.58	0.114
rs3821203	90622	102618272	A/G	0.55	0.61	0.104
rs3771162	90924	102618574	A/T	0.34	0.29	0.261
rs3213733	91634	102619284	G/T	0.77	0.73	0.260
rs3213732	92029	102619679	C/T	0.48	0.41	0.026
rs1035130	95152	102622802	A/G	0.55	0.60	0.152
rs3752659	95348	102622998	C/T	0.81	0.80	0.760
rs3755274	96145	102623795	C/T	0.25	0.22	0.319
rs2241117	96793	102624443	A/G	0.72	0.80	0.024
rs2241116	97015	102624665	G/T	0.18	NA	NA
rs881890	97064	102624714	C/T			
rs3771161	97711	102625361	G/T	0.71	0.66	0.146
rs3771160	97855	102625505	A/C			
rs3771159	98708	102626358	A/G	0.38	0.42	0.175
rs1420104	not mapped	not mapped	С/Т			
rs2041738	not mapped	not mapped	A/C			

**TABLE 15** 

dbSNP rs# Position in SEQ ID NO:		Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value	
rs884517	207	102527857	С/Т				
rs1476984			A/G	0.84	0.85	0.759	
rs951774	6414	102533669 102534064	A/C	0.77	0.75	0.730	
rs2041737	7341	102534991	A/G	0.40	NA NA		
rs1420091	10984	102538634	A/G	0.34 0.35		0.819	
rs2110660	12351	102540001	C/G	0.43	0.42	0.665	
rs1362347	13335	102540985	A/G	0.82	0.83	0.576	
rs3073968	16584	102544234	/TGTG/T GTGAG	0.49	0.49	0.997	
rs4090473	16737	102544387	C/G	0.44	0.45	0.732	
rs1558622	23897	102551547	C/T	0.42	0.42	0.976	
rs1558621	24057	102551707	C/T	0.31	0.31	0.926	
rs1558620			A/G	0.42	0.42	0.867	
	rs1558619 25300 103		A/C	0.48	0.48	0.930	
rs950881	26262	102553912	A/C	0.73	0.73	0.955	
rs950880	26312	102553962	G/T	0.48	0.49	0.919	
rs1362346	26589	102554239	C/T				
rs1968171	27302	102554952	A/G	0.43	0.42	0.717	
rs1813299			A/T				
	rs1813298 27451		C/G				
rs1968170 27552		102555202 102558381	C/T	0.67	0.66	0.830	
	rs974389 30731		C/T	0.44	0.45	0.857	
rs971764	32085	102559735	A/G	0.43	0.42	0.845	
rs1420089	32139	102559789	A/G	0.15	0.16	0.809	
rs1420088	33184	102560834	A/G				
rs1420103	42382	102570032	G/T 0.68		0.63	0.178	
rs1420102	42569	102570219	A/G	0.45	0.44	0.722	
rs1997467	44823	102572473	C/T				
rs1997466	45217	102572867	C/G	0.44	0.43	0.805	
rs1362350	45548	102573198	C/G	0.45	0.46	0.890	
rs2310220	45601	102573251	A/G	0.38	0.37	0.661	
rs1362349	45722	102573372	C/G	0.41	0.40	0.762	
rs3755278	45967	102573617	A/G	0.07	0.07	0.984	

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value	
rs3771180	771180 47367		A/C	0.88	0.87	0.812	
rs3771179	47642	102575017 102575292	A/C	0.07	0.07	0.868	
rs985523	48126	102575776	C/T	0.16	0.13	0.270	
rs1041973	49218	102576868	A/C				
rs3214363	49274	102576924	-/A				
rs873022	49433	102577083	G/T	0.57	0.56	0.868	
rs3771177	49610	102577260	A/C	0.29	0.29	0.988	
rs3732129	51282	102578932	A/G	0.51	0.52	0.795	
rs1420101	51466	102579116	A/G	0.60	0.61	0.864	
rs12905	53757	102581407	A/G	0.26	0.26	0.994	
rs3771175	53960	102581610	A/T	0.82	0.80	0.444	
rs3821204	54031	102581681	C/G	0.22	0.23	0.769	
rs2160203	54574	102582224	C/T	0.23	0.25	0.732	
rs1946131	55679	102583329	A/G	0.75	0.77	0.723	
rs1054096	56100	102583750	C/T	0.66	0.65	0.807	
rs2287038	56182	102583832	С/Т	0.97	0.00		
rs1921622	59817	102587467	A/G	0.44	0.46	0.472	
rs1861246	60533	102588183	A/G	0.23	0.24	0.824	
rs1861245	60656	102588306	A/G	0.36	0.34	0.590	
rs3755276	72209	102599859	A/G	0.48	0.51	0.423	
rs2287037	72778	102600428	A/G	0.55	0.54	0.941	
rs1420099	74293	102601943	C/G	0.58	0.58	0.904	
rs3771174	77335	102604985	A/G				
rs1420098	78029	102605679	A/G	0.30	0.31	0.827	
rs1362348	78374	102606024	C/G	0.04	-0.01		
rs1882348	78421	102606071	A/T	0.31	0.31	0.968	
rs1558627	78434	102606084	C/T	0.60	0.59	0.839	
rs2058622	79174	102606824	C/T	0.56	0.55	0.961	
rs3836110	79397	102607047	-/G	0.70	0.69	0.643	
rs3771172	79562	102607212	A/G	0.24	0.22	0.675	
rs3771171	79700	102607350	A/G				
rs3771170	79730	102607380	A/T	0.26	0.27	0.700	
rs2160202	79904	102607554	C/T	untyped	0.64	NA	
rs2058623	79920	102607570	A/G	0.65	0.62	0.389	
rs3771167	79938	102607588	C/T				
rs3771166	79972	102607622	C/T	0.53	0.53	0.820	
rs1974675	80125	102607775	С/Т	0.55	0.56	0.842	
rs1465321	80368	102608018	A/G	0.29	0.30	0.781	
rs2041740	83484	102611134	C/T	0.28	0.30	0.658	
rs3771164	85536	102613186	A/T	0.73	0.74	0.905	
rs2270298	85829	102613479	C/T	0.19	0.18	0.654	
rs2270297	86425	102614075	A/G	0.57	0.53	0.249	
rs2041739	88083	102615733	A/G	0.42	0.41	0.892	
rs2080289	88770	102616420	C/T	0.61	0.60	0.840	
rs3821203	90622	102618272	A/G	0.62	0.63	0.927	
rs3771162	90924	102618574	A/T	0.26	0.25	0.621	
rs3213733	91634	102619284	G/T	0.75	0.74	0.728	
rs3213732	92029	102619679	С/Т	0.39	0.45	0.176	
rs1035130	95152	102622802	A/G	0.62	0.63	0.792	
rs3752659	95348	102622998	С/Т	0.79	0.80	0.826	
rs3755274	96145	102623795	C/T	0.27	0.29	0.618	
rs2241117	96793	102624443	A/G	0.70	0.67	0.480	
rs2241116	97015	102624665	G/T	0.15	0.15	0.849	
rs881890	97064	102624714	C/T				
rs3771161	97711	102625361	G/T	0.68	0.70	0.681	
rs3771160	97855	102625505	A/C				
rs3771159	98708	102626358	A/G	0.37	0.37	0.970	

dbSNP rs#	Position in SEQ ID NO:	Chromosome Position	A1/A2 Allele	F A2 Case AF	F A2 Control AF	F p- Value
rs1420104	not mapped	not mapped	С/Т			
rs2041738	not mapped	not mapped	A/C			

[0233] Allelotyping results were considered particularly significant with a calculated p-value of less than or equal to 0.05 for allelotype results. These values are indicated in bold. The allelotyping p-values were plotted in Figure 1 for the discovery cohort. The position of each SNP on the chromosome is presented on the x-axis. The y-axis gives the negative logarithm (base 10) of the p-value comparing the estimated allele in the case group to that of the control group. The minor allele frequency of the control group for each SNP designated by an X or other symbol on the graphs in Figure 1 can be determined by consulting Tables 13. For example, the left-most X on the left graph is at position 102527857. By proceeding down the Table from top to bottom and across the graphs from left to right the allele frequency associated with each symbol shown can be determined.

[0234] To aid the interpretation, multiple lines have been added to the graph. The broken horizontal lines are drawn at two common significance levels, 0.05 and 0.01. The vertical broken lines are drawn every 20kb to assist in the interpretation of distances between SNPs. Two other lines are drawn to expose linear trends in the association of SNPs to the disease. The light gray line (or generally bottommost curve) is a nonlinear smoother through the data points on the graph using a local polynomial regression method (W.S. Cleveland, E. Grosse and W.M. Shyu (1992) Local regression models. Chapter 8 of Statistical Models in S eds J.M. Chambers and T.J. Hastie, Wadsworth & Brooks/Cole.). The black line provides a local test for excess statistical significance to identify regions of association. This was created by use of a 10kb sliding window with 1kb step sizes. Within each window, a chi-square goodness of fit test was applied to compare the proportion of SNPs that were significant at a test wise level of 0.01, to the proportion that would be expected by chance alone (0.05 for the methods used here). Resulting p-values that were less than 10<sup>-8</sup> were truncated at that value.

[0235] Finally, the exons and introns of the genes in the covered region are plotted below each graph at the appropriate chromosomal positions. The gene boundary is indicated by the broken horizontal line. The exon positions are shown as thick, unbroken bars. An arrow is place at the 3' end of each gene to show the direction of transcription.

### Example 5

# Gene expression profiling in IL-1 beta and PMA stimulated SW1353 cells

[0236] The human chondrosarcoma cell line, SW1353, (ATCC HTB-94) was grown in L-15 media containing 10% FCS. Culture conditions were at 37 degrees with 0% CO2 with media changes every 2-3

days. SW1353 cells were grown to 80-90% confluence in 10 cm dishes and then stimulated with either 10ng/ml IL-1 beta (human recombinant, Research Diagnostics) or with 200nm PMA (Sigma). IL-1 beta stimulation was for 3 and 24 hours and PMA stimulation was for 3 and 24 hours. Control cells were grown and extracted in parallel with treated cells. As shown in Figure 6, IL1RL1 was seen to be upregulated by IL1-beta and by phorbol esters in a human chondrocyte cell line model (SW1353 monolayer cell line).

[0237] The expression profiling in IL-1 beta and PMA stimulated SW1353 cells grown in 3-D alginate cultures W1353 cells were cultured as above and then resupended in 1.2% alginate beads at a density of 4 millions cells/ml according to the manufacturer (Cambrex). Cells were grown for 2 weeks and an alginate bead was removed from culture and tested for the presence of proteoglycans by Alcian Blue staining (Sigma). Positive staining indicated that the chondrocytes were expressing ECM proteins. Alginate cultures were then stimulated with IL-1 beta for 24 hours or with PMA for 3 hours. Control cells were grown and extracted in parallel with treated cells. As shown in Figure 7, IL1RL1 was seen to be upregulated by IL1-beta and by phorbol esters in a human chondrocyte cell line model (SW1353 3-D alginate cell line).

### RNA extraction and cDNA synthesis

[0238] Cells from control chondrocytes and stimulated chondrocytes were isolated at the appropriate time period. mRNA was isolated from total cell lysates using poly dT beads according to the manufacturer (Dynal). Isolated mRNA was used to generate cDNA using SuperScript II reverse transcriptase according to the manufacturer (Invitrogen).

### Expression profiling using semi-quantitative PCR

[0239] cDNA levels were normalized using the housekeeping gene, GAPDH. Specific primers corresponding to MMP8 and MMP13 were used in semi-quantitative PCR as positive indicators of induction of an osteoarthritic phenotype. All specific primers used, including MMP8, MMP13, BVES, CHDC1 and IL1RL1 (transmembrane form, soluble form, soluble isoform 1 and soluble isoform 2) for semi-quantitative PCR and are listed in Table 16.

		1
Gene	Forward primer	Reverse primer
GAPDH	ATCATCTCTGCCCCCTCTG	GAGGATTGCTGATGATCTTC
MMP8	CAATACTGGGCTCTGAGTGG	GGAAAGGCACCTGATATGC
MMP13	ATATCTGAACTGGGTCTTCC	GACAGCATCTACTTTATCACC
BVES	AACAGTATAGCCAGCTCC	ATCATCATCTTCTGCTCC
CHDC1	CCAAAGATCAGGACATGGATA	TGCTGTTTGTGGTAGGAGAG
IL1RL1 (TM)	CCACTCTGCTCTGGAGAGAC	GCCTGCTCTTTCGTATGTTG
IL1RL1 (Sol)	TCCGTCACTGACTCCAAGTT	TTGCTGCTGTGGAATACATG

**TABLE 16: Primer Sequences for Expression Profiling** 

Gene	Forward primer	Reverse primer			
IL1RL1 (ST2_3)	AGGCTTTTCTCTGTTTCC	GTTGAATTCTTGGTTCACC			
IL1RL1 (ST2_2)	TAATGTGATGACTGAGGACG	TGCAGAAACTCTGACACC			

[0240] In a human chondrocyte cell line model, IL1RL1 was seen to be upregulated by IL1-beta and by phorbol esters. IL1RL1 has an unknown function, but it may possibly mediated inflammatory responses that can contribute to the development of OA. IL1RL1 is druggable by antibodies or by protein agents.

### Example 6

# In Vitro Production of Target Polypeptides

[0241] cDNA is cloned into a pIVEX 2.3-MCS vector (Roche Biochem) using a directional cloning method. A cDNA insert is prepared using PCR with forward and reverse primers having 5' restriction site tags (in frame) and 5-6 additional nucleotides in addition to 3' gene-specific portions, the latter of which is typically about twenty to about twenty-five base pairs in length. A Sal I restriction site is introduced by the forward primer and a Sma I restriction site is introduced by the reverse primer. The ends of PCR products are cut with the corresponding restriction enzymes (*i.e.*, Sal I and Sma I) and the products are gel-purified. The pIVEX 2.3-MCS vector is linearized using the same restriction enzymes, and the fragment with the correct sized fragment is isolated by gel-purification. Purified PCR product is ligated into the linearized pIVEX 2.3-MCS vector and *E. coli* cells transformed for plasmid amplification. The newly constructed expression vector is verified by restriction mapping and used for protein production.

[0242] E. coli lysate is reconstituted with 0.25 ml of Reconstitution Buffer, the Reaction Mix is reconstituted with 0.8 ml of Reconstitution Buffer; the Feeding Mix is reconstituted with 10.5 ml of Reconstitution Buffer; and the Energy Mix is reconstituted with 0.6 ml of Reconstitution Buffer. 0.5 ml of the Energy Mix was added to the Feeding Mix to obtain the Feeding Solution. 0.75 ml of Reaction Mix, 50 µl of Energy Mix, and 10 µg of the template DNA is added to the E. coli lysate.

[0243] Using the reaction device (Roche Biochem), 1 ml of the Reaction Solution is loaded into the reaction compartment. The reaction device is turned upside-down and 10 ml of the Feeding Solution is loaded into the feeding compartment. All lids are closed and the reaction device is loaded into the RTS500 instrument. The instrument is run at 30°C for 24 hours with a stir bar speed of 150 rpm. The pIVEX 2.3 MCS vector includes a nucleotide sequence that encodes six consecutive histidine amino acids on the C-terminal end of the target polypeptide for the purpose of protein purification. Target polypeptide is purified by contacting the contents of reaction device with resin modified with Ni<sup>2+</sup> ions. Target polypeptide is eluted from the resin with a solution containing free Ni<sup>2+</sup> ions.

### Example 7

# Cellular Production of Target Polypeptides

[0244] Nucleic acids are cloned into DNA plasmids having phage recombination cites and target polypeptides are expressed therefrom in a variety of host cells. Alpha phage genomic DNA contains short sequences known as attP sites, and *E. coli* genomic DNA contains unique, short sequences known as attB sites. These regions share homology, allowing for integration of phage DNA into *E. coli* via directional, site-specific recombination using the phage protein Int and the *E. coli* protein IHF. Integration produces two new att sites, L and R, which flank the inserted prophage DNA. Phage excision from *E. coli* genomic DNA can also be accomplished using these two proteins with the addition of a second phage protein, Xis. DNA vectors have been produced where the integration/excision process is modified to allow for the directional integration or excision of a target DNA fragment into a backbone vector in a rapid *in vitro* reaction (Gateway™ Technology (Invitrogen, Inc.)).

[0245] A first step is to transfer the nucleic acid insert into a shuttle vector that contains attL sites surrounding the negative selection gene, ccdB (e.g. pENTER vector, Invitrogen, Inc.). This transfer process is accomplished by digesting the nucleic acid from a DNA vector used for sequencing, and to ligate it into the multicloning site of the shuttle vector, which will place it between the two attL sites while removing the negative selection gene ccdB. A second method is to amplify the nucleic acid by the polymerase chain reaction (PCR) with primers containing attB sites. The amplified fragment then is integrated into the shuttle vector using Int and IHF. A third method is to utilize a topoisomerase-mediated process, in which the nucleic acid is amplified via PCR using gene-specific primers with the 5' upstream primer containing an additional CACC sequence (e.g., TOPO® expression kit (Invitrogen, Inc.)). In conjunction with Topoisomerase I, the PCR amplified fragment can be cloned into the shuttle vector via the attL sites in the correct orientation.

[0246] Once the nucleic acid is transferred into the shuttle vector, it can be cloned into an expression vector having attR sites. Several vectors containing attR sites for expression of target polypeptide as a native polypeptide, N-fusion polypeptide, and C-fusion polypeptides are commercially available (e.g., pDEST (Invitrogen, Inc.)), and any vector can be converted into an expression vector for receiving a nucleic acid from the shuttle vector by introducing an insert having an attR site flanked by an antibiotic resistant gene for selection using the standard methods described above. Transfer of the nucleic acid from the shuttle vector is accomplished by directional recombination using Int, IHF, and Xis (LR clonase). Then the desired sequence can be transferred to an expression vector by carrying out a one hour incubation at room temperature with Int, IHF, and Xis, a ten minute incubation at 37°C with proteinase K, transforming bacteria and allowing expression for one hour, and then plating on selective media. Generally, 90% cloning efficiency is achieved by this method. Examples of expression vectors

are pDEST 14 bacterial expression vector with att7 promoter, pDEST 15 bacterial expression vector with a T7 promoter and a N-terminal GST tag, pDEST 17 bacterial vector with a T7 promoter and a N-terminal polyhistidine affinity tag, and pDEST 12.2 mammalian expression vector with a CMV promoter and neo resistance gene. These expression vectors or others like them are transformed or transfected into cells for expression of the target polypeptide or polypeptide variants. These expression vectors are often transfected, for example, into murine-transformed a adipocyte cell line 3T3-L1, (ATCC), human embryonic kidney cell line 293, and rat cardiomyocyte cell line H9C2.

[0247] Modifications may be made to the foregoing without departing from the basic aspects of the invention. Although the invention has been described in substantial detail with reference to one or more specific embodiments, those of skill in the art will recognize that changes may be made to the embodiments specifically disclosed in this application, yet these modifications and improvements are within the scope and spirit of the invention, as set forth in the claims which follow. All publications or patent documents cited in this specification are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference.

[0248] Citation of the above publications or documents is not intended as an admission that any of the foregoing is pertinent prior art, nor does it constitute any admission as to the contents or date of these publications or documents. U.S. patents and other publications referenced herein are hereby incorporated by reference.

# Nucleotide and Amino Acid Sequence Embodiments

[0249] Table A includes information pertaining to the incident polymorphic variant associated with osteoarthritis identified herein. Public information pertaining to the polymorphism and the genomic sequence that includes the polymorphism are indicated. The genomic sequences identified in Table A may be accessed at the http address www.ncbi.nih.gov/entrez/query.fcgi, for example, by using the publicly available SNP reference number (e.g., rs1041973). The chromosome position refers to the position of the SNP within NCBI's Genome Build 34, which may be accessed at the following http address: www.ncbi.nlm.nih.gov/mapview/map\_search.cgi?chr=hum\_chr.inf&query=. The "Contig Position" provided in Table A corresponds to a nucleotide position set forth in the contig sequence (see "Contig Accession No."), and designates the polymorphic site corresponding to the SNP reference number. The sequence containing the polymorphisms also may be referenced by the "Nucleotide Accession No." set forth in Table A. The "Sequence Identification" corresponds to cDNA sequence that encodes associated target polypeptides (e.g., IL1RL1) of the invention. The position of the SNP within the cDNA sequence is provided in the "Sequence Position" column of Table A. If the SNP falls within an exon, the corresponding amino acid position (and amino acid change, if applicable) is provided as

well. Also, the allelic variation at the polymorphic site and the allelic variant identified as associated with osteoarthritis is specified in Table A. All nucleotide and polypeptide sequences referenced and accessed by the parameters set forth in Table A are incorporated herein by reference.

TABLE A

RS_ID	Chromo -some	Chrom Position	Contig Accession No. [1]	Contig Position		Sequence	Amino Acid Position	Locus	Locus ID	A [3]	Allelic Variability	OA Assoc. Allele
1041973	2	102576868	Hs2_22327_34:13	5021492	NM_003856	coding- nonsynon	E78A	IL1RL1	9173	R	[A/C]	С

[1] Contig Accession Number which can be found in the NCBI Database: http address: www.ncbi.nih.gov/entrez/query.fcgi

[2] Sequence Identification or Nucleotide Accession Number which can be found in the NCBI Database:

http address: www.ncbi.nih.gov/entrez/query.fcgi

[3] "A" column is the sequence orientation ("F" is forward, "R" is reverse).

[0250] The following is a genomic nucleotide sequence for an *IL1RL1* region. The following nucleotide representations are used throughout: "A" or "a" is adenosine, adenine, or adenylic acid; "C" or "c" is cytidine, cytosine, or cytidylic acid; "G" or "g" is guanosine, guanine, or guanylic acid; "T" or "t" is thymidine, thymine, or thymidylic acid; and "I" or "i" is inosine, hypoxanthine, or inosinic acid. Exons are indicated in italicized lower case type, introns are depicted in normal text lower case type, and polymorphic sites are depicted in bold upper case type. SNPs are designated by the following convention: "R" represents A or G, "M" represents A or C; "W" represents A or T; "Y" represents C or T; "S" represents C or G; "K" represents G or T; "V" represents A, C or G; "H" represents A, C, or T; "D" represents A, G, or T; "B" represents C, G, or T; and "N" represents A, G, C, or T.

## IL1RL1 Genomic Sequence (SEQ ID NO: 1)

>2:102527651-102626600

```
1
       tgcaggattt gtatgtaagt atccaacaga caacactgg ggtgatcctg tgaatgtaga
61
       ggaaqacqtq atcttaaatq qaccaqqqaa qtaaqacctt aqqatqtacc aqccaqctqa
       gtggcaggta atattgattc tcatgtttca ccccatgttc ggtcacttgt gagcagattc
121
181
       ttcagttcaa ttctctggct agttaaYttg gtctgggagt cacaccagaa aatgtttctg
241
       aaaataggta cctattgtga aaagtataat cctctttgct attgtgagta gtgctgcaat
301
       aaacatacat gtgcatgtgt ctttatagca gcatgattta tattcccttg ggtatatacc
361
       cagtaatggg atggctgggt caaatgatat ttctggttct agatccctga ggggagcggg
421
       gagggatagc attaggagat atacctaatg taaatgacga gttactgggt gcagcacacc
       aacgtggcac atgtatacat atgtaacaaa cctccacgtt atgcacatgt accctagaac
481
541
       ttaaaqtata ataaaaaata tatataaaqt ataatcatca aattqttaaa atcatqattc
601
       tcatgaagta gattgtgtct gttgaagttt aattaatgaa ggaaacaaca ggatgtcagg
661
       actgaatcca taaggatctg agaaactggt atttccataa gatgttaaga taatattgat
721
       aactgggata caaaactgct taatgtccta caatgtaata aaacaatttc tgttgtgttt
781
       tgcatgagat agaaacaaaa atacctccca ttggtcagtt ttctacaaga taatctttaa
```

```
841
       ctttacaggg ttgtaaaata gctatgcctg aattgttagt taaaacccca ttttttttt
901
       tagagaaaac aatgateett agacteaett actagaaata tgttetagtt tttttataaa
       961
1021
       aataaaataa aaaatattta ttttaaatat tttttcaag catcagtata ccttattctt
1081
       gttcaacttg gggaatggcc ttattcaaat tgggaatcct cagctcatct ttgtctttgg
1141
       cattccagag aatgaatgtg tttgggaaat tgtgtcagct gagcagtgta tacaqaaaqq
1201
       gacaqqaqac cactqaaatq qttcaqqtta taqttaqata qttcctqcca ctqqttaqat
1261
       caggagagtt agatggttat agttagatag ttctttatgc tggcagtggg aaacaagaga
1321
       agaggacata tgagggagat gctcatgtta ttgggaacaa tatttggctc ctggtagaat
       gttaggggcg agggagaaga ttctaatgtg ttgagttcgg gagccgctgc gccattaata
1381
1441
       aaaccagaaa cettetacag tgetecaggt gggagecatt agcateaaag gegattetaa
1501
       aatcagtcct gaaagctatt ctatcccttg ttctgcatgc agaacagctc ctgagggcct
1561
       gttgagggcc cagggaaact ttggcccagg gccaaggaag gactggtgtt cccttttgqt
1621
       ttagetetge teacagacaa etgeecagaa ggtecaggtt ecaagteact gtgeaagetg
1681
       acacqtqtqa qacataqqaa qctqttqqqa ctacaqaacc caqqqactcc aaaccctqtt
1741
       taagtaccac acctgagggg aagaagaccc tgcttcccaa ttagtcagct gggtagaagg
       ctgaggattt ttgaaaagag gtagttggtt tagcctcagc tttctttaga ataatattgt
1801
1861
       ggtttgaaag tgtgcaacct cctagtgtca tcatttcctt ttgtccctga cagaactgtc
1921
       ttcacgctgg aaataaaagt atgggacagg cttgaagcag agttgtgaag tagaagtttg
1981
       ctaggaatcc attittatc gaaagcccca ggcaactctg attaatttgg tctgggagtc
2041
       atacatggag aaatactgtt ctaggagctc attcagctaa cacctctctc acctagaggt
2101
       catgogggga aagaaatgtg tattgcatta taattcatag tagactgtga ataactaata
2161
       ggaaaagatt cctttcagct tctcctgtct agttaagttc cagcagggac acacctttag
2221
       cttaaaggaa tgcattttag tggtgatggt ggtggtggtg gtggtgatgg tggtggtggt
2281
       ggtggtggtg gtgtgtcttc ctaggtcctc tcaaatcaca tcacatggtg gagggataaa
2341
       aaqaqqcaqq taaaaaataq catttqttct acaattqatq aatqactaaa tqtcctcttq
2401
       aacataaaga atttatattg gtaaaagctt ctctgctatg tgcattagta aaaacaacaa
2461
       categtgccg ggcgcggtgg ctcacgcctg taattccagc gctttgggag gccaaggcgg
2521
       gtggatcaca aggtcaggag atcgagacca tcctggctaa cacggtgaaa tcccqtctct
2581
       actaaaaata caaaaaatta gccagcgtgc tggcaggtgc ctgtagtccc agctactcgq
2641
       gaggctgagg caggagaatg gtgtaagtga accegggagg eggageetge agtgageega
2701
       2761
       tegtttgetg atttagaact ettettaaag ageeteeata tgtttaaaga geeatgeett
2821
       atgactgtat gactataatt ttataaagcg ctttccctac agtatcttac taaatccttt
2881
       cagcaatcct ggaagtacca tcattatctt tattttgcat attcaagttc tgaaagcctg
2941
       gtttgctcca gtaagtagta gagctaggac caaaacccaa gttttggttg ggttttcagc
3001
       cctgaatctt tcaqccctqa accttttcat aqctqcctct qacttcacaq ctcactqcat
3061
       aaagccttct tcaqacccat cctctqtcaq catcqtctaq tctaqcctqt caqqtatttt
3121
       ggcctccaca ggagatccta ttctcaatgt ttgtaaatat ttagctaaat ctgtagcaga
3181
       catctggttt cataatgaga aagagcattt ttgttttcat tagggaaaaa gagcatgact
3241
       gggcgcagta actgcagttt tcatcgtgga gaaaaagaga gtgagacttg aacttgaaaa
3301
       agtgaagcca agcagtgcag ttggcaggtg gagggtgggg aggagggaga ataaagacaa
3361
       gtcaggagga gagagggaa aatgatgcac agagaaagag ggaagaggca aagagaaagg
3421
       aagggagcct ggacagcaga ggggtctcct tcaggaaaca gagttgaagt gaaatagtgt
3481
       aatccccaqq tqqcaqqtac aacatttctc aqcacaatqa ctctqaacaa cttaattctt
3541
       tottagggga agtagttaca oggaaagtgt aatgtgtgct gcacatttgt agttgtgata
3601
       ccatcagaaa gaggtacttg ctgtggaaag aacttctcaa cccctccatg taatgaccct
3661
       tggagaatgc attatcaaag cagagtcctg cagcctgaaa gtaaacagca tctacttaga
3721
       aatacagtga catgtaaact ttccccaaat caacagtgaa accagtagca ataactttga
3781
       tgaaacaaag gggcattgcc gagctgtcag gacacaacag agcacaacag aacagcccca
3841
       ggtaaatatt gataaattag gttgatctgt gtattcttag tcatgggaag ctggggcaga
       taacgaaacc ttgtatgtgt cctttatagc tctgtctgat gagtttgggc gacggagtct
3901
3961
       tgctctgtcg cccaggctgg agtgcaatgg cgtgatctca gctcactgca acctctgcct
4021
       ccctgattca agtgattctc ctgcctcagc ctcctgagta gctgggatta caggcacgtg
       ccqccacqcc aggctaattt ttgtattttt agtagagaca gggtttcacc atgttgatca
4081
4141
       ggctggtctc gaactcctaa actcatgatc cgcccacctc agcctcctga agtgctggqa
4201
       ttacaagcct gagccaccgt gcccagcctt gtttttttt aagtgatttt tgagggcagt
4261
       tttatgggca gaaaagactc ttcgtgaatt tattttgatt ttaaaattac tattcaattt
4321
       ttcaaagtta aaaatgtaaa aaatctagtt tattataaat tgaagaccag cttaaaaatc
4381
       atataatacc atttatatgt ctagcacatt ttaacaaact tcaacactag gttactccac
4441
       ttacaaattt qqcaaactaa atattttaaq qtattttaaa ccqcatttat acatttactq
4501
       tatttgattt ttcttcttaa gaactttggc taaaaccttc ccaagtaatg aattaattct
4561
       tataaattta agttototta gaaattttga ogotacatat aagottaaat attggatata
4621
       aacttagtaa ggatttcact taaaatcgta aatctaaaca atttccaatt ataccttttt
```

```
4681
       aaaaaagagt tacatgactt attctaatgg gtcaatattt taggcatcac aaatacataa
4741
       ttgcaaatat atagattcct taaattatac catggtcttg attcttatta catagatgtg
4801
       tacatacctt tctacccagg tctaaatacc actgaattca aaatgagtct tgcattctaa
4861
       gtacttcgaa attcttaatg ggtattaaaa agtgacccag ttgcttatgg ataaacctca
4921
       taatttgcct tgatacttgg tcacaaatta aaggtcattc tactgacaaa ttacttgtat
4981
       acaatctaca aaacaaataq catqcaattc tcattacaqc acacqtqttc aqactttaac
5041
       aaagctgttt gaagtctaca tcatctcctg agttagagaa gccccaattg ccacatacag
5101
       atacttagta ctgtcataca gagtcacata tctactggat tctacctgga tcttaggatc
5161
       aagatgcttg tactcagaca aacaaattag gaaagtagcc tcctttgcag agtggtctag
5221
       gggtgttacg agccaaaaaa tcattcacgg ttttatggtt aagaagtgag atgatctgta
5281
       atgaaacctc acagagattc tgttaaataa caagtgccca aaagctactt ggaaaaggag
5341
       tagactgaag agaaaatgag agaatcaaaa atagttttaa tttcctgtta gaatcatctc
5401
       ctttacttta ctcttgcatg tgtaactgtg gaggagattg taaacattcc cttagctaac
       tgtacttgtt actcttgatc aacatccctt agtaactgtg tatatgtgat gcttcatgca
5461
5521
       ttagtcagag ttgagaacaa aaacagagta gctcaaacaa aacataggtt tatctgtccc
5581
       tcatgtaaca acacatgggt agccaggtgc tctagggatg gtagatggat ctgcttcatg
       aaqtcactca qaatccccaa ttcctcctac cttgctgctc tgccctgtgt ggtcctcatc
5641
5701
       tgtctcaccg ctgaagccac cttgcagttc acctagaaga gaatatgtac tgtttctgcc
5761
       aaatctgtgc tatagatact agatgctttt aatggagagg ttagccaaga aacatgacat
5821
       ttctcttata ggtaataaaa tacttctctc taaaqaaata cttagtttaa aattttattc
5881
       tgctatagtt ttagcagtag gataaaagct tagtttgcag aagcggaaat tatcctatat
5941
       ttaggcaaag tgagctatct gtgtgttatg attagagaga ataccaagaa gtgatttcct
6001
       tccttaaaat aacttctcYa aatttccacc attaactacc cattcttcaa ctctctcaat
6061
       ttttcttatt cctcccatgg agggagtgca tgtatttaat tactaacttt actagtatta
6121
       tataatctat gctccaaaat ggttagtcta caattagtag agggataagt ctttagggtt
6181
       ggaaagaaaa aacagttatt aaagatctac tattgctgga tgtcatacta actcttctta
       cttattatgt tatttatacc acctaacagt ttgataatat tacccatccc cccgcttttt
6241
6301
       tttttaagag ataaggcata tatgacttat atagagttta ggggattttc ctcaagatta
6361
       gacagttaga agtctcttgg gtgggagagc agttctaggg gaaaaggaaa gatMagaaag
6421
       acaagtcaac tactttgagt aaagacctct ggctgagcag gtcattgtgt attcccagaa
6481
       tttgtcacca actagttgtg tggttttgga catacccctt cacatcacct caatgtcacc
6541
       tttggtaaag tgaaagaatt ggttgtagat attctttaaa ttatcttta ttcaggcttt
6601
       ctacaqccca attctctcaa ttgttgactt ttacttaqaa ttttgqqcaa acactaqctq
6661
       atacaatcca tttaccaaga tgtttatctg atgacatttt catcaagttt ctaaagatgc
6721
       catattgtct ttctggcagg cttttggaaa gtgttcaaga tctcctgttt tattttcaca
6781
       gttggcccct gagcctcaag tgtaagcaga gctgggctga gaggcttagg tcctcctggc
6841
       tgatgtcact aggctgaggg gccttgggga agaaaacaaa tttctcccag attcaaccct
6901
       ccaaccaact gcaggtcctg gaacagtgag taactccttt actcagaagc tgaacattaa
6961
       aagtetttte tgattettae caacttggea cataagaaat acagteattt eetcaaatae
7021
       atttttatta tgagggtact gaacattgta tttattggtc atttgcattt tttgtaaatt
       gtagattttt aaaaacctct ttgtctgggt gcggtggctc acacctgtaa tcccagcact
7081
7141
       ttgggaggcc gaggaaggtg gatcatctga ggtcaggagt ttgagaccag cctgaccaac
7201
       atggtgaaac cccatctcta ctaaaaatac aaaaattagc tgggcttggt ggcgtgcgcc
7261
       tgtaatccca gttactccgg aggctgaggc aggagaatag cttgaaacca agaggtggag
7321
       gttgcagtga gccaagatcg Ygccactgca ctagcctggg caacaagagg gaaactccat
7381
       ctcaaaaaac aaaaacaaac aaacaaacaa aaacattttt gtattaacta gtaaagtctc
7441
       ctgatacatt aagaatatta aagacaggca tatagtgagg ttttttttgtt tagtttccat
7501
       atgeetttta atttgtttaa ggatatagte atttgtttag aetataggta aggaatagtt
7561
       atatcttttq ttccaqaaaq tttcaqattt tqaqqqaata ttactattcc taaqqtttta
7621
       ttgacatata aattttaaaa atattatgta atgagagcta tatatatata ttctttttt
7681
       gtgcatttta tcatgggtac taggttaagt gtctgtttct atgaacatac aacttactta
7741
       acatatgact tggaaacaga acatgcatag taaatgtttc tcaagtaaaat aagtaaaagt
7801
       atatgagaaa caaatgtgct gtttacatgt tgtcatagga atcatcttag cactaaatgg
7861
       ataattagga tocaggttca toottgggot tggcacaaat gagottggac aatotgaaat
7921
       atggatggtt gataataggt aattaggttt aaattatacc aqqaaactga aqaacattca
7981
       gaaaatggtt acagataatt acagaaggtt gcattacaga ggaaaaattt ctcacagtga
8041
       gagacaattg gcctcacatg cctttactga ggcatacatt gtgtcataat ttccttttat
8101
       tgcaccttca acattttgaa gggaattaat tgtctgaatt aggctagatg tcatttttt
8161
       cttggaatta gaagaaagat tgagtaaccc cttatgttta tctggaacat gtctttctgt
       qcatttaaaa atctcagagc ccttcagaac cttgcccgtg acccgtcatt gggaatatct
8221
8281
       taaaaatgtg tgtgtgtggg tgggtgtgtg tgtgtgaata taggaatgag ggaacactgt
8341
       cgtagacaga tatatctgaa gcttatcttt tggtttaggg catttggtct tgcatcacct
8401
        tttaagtott atgotatoot tgacttttga acagaacaaa ottotagatt tttgtatgtt
8461
       ctaagttctg cctttgcagt attagaggtc tattttccca gactagggac tggagtgttt
```

```
8521
       aatcatcact gtcatgtaga cctaatatag aaatacaaat gcaataagaa aaatgttgag
8581
       aaattaaagt cagaaacata tttgttgtga tacatgttca aaaataatat atttcattgc
8641
       aggaaatagc aaatccagag aaataaaaag aaagaaaaaa aataatctgc atttctacgc
8701
       atgcagctaa ccattgctga gattttggag tccttgtttc ctaggcacgt tttcctatgt
8761
       ctgtgtgcac acatgtatgg gatgttgcat atgtgtatgt gtgcatgtgt ttacacatgt
8821
       atttgcaggt atgtgttgat tattatacat tgagtcataa tgtatattac atttaaaatt
8881
       tcctgttttt acggctttat taaggtataa ttgagatatg aaaaattgca tacacttaaa
8941
       ttgtacagtc taagttttgt catatgtata cacaccagtg aaacccttac cacaatcaag
9001
       ataatgaaca tacccatccc ccgtaaagct tctacacccc tttcqaagtc catccttcaa
9061
       cttctcctac cgaatccaca ggcatccact gacctgattt ctgttactat aagtttgttc
9121
       acattttcta gaaattgata taaatggaat aatacagcat gtgcttgttt ctctctggct
9181
       tottotatto tgogtaatta ttgtgaaagt catccatatg attgcattta tcagtacatt
9241
       catactattg catttatcaa tatgtcattc ttgttgattg tttatcatat tcccctatta
9301
       tettettaae atetatagaa aettagtggt gteaaetete tttttettga tattagtaat
9361
       ttgtgtcttc tttctttttt cctcagtcag tgtgactaga gagttatcaa ctttattagt
9421
       cttctcaaca aaccagettt tggcatcatt gatttttett tattgttttt etgttecate
9481
       teactgtttt etgeattgat ttgtattatt teattttett etgttttetg tgetttettt
9541
       gctcttcttt ttctagtttt ttaaggtggt aaccaatatt gatttatgac ttttattatt
9601
       ttctaatgta tgcctttagt gttgtaaatt tccctcttta gtggcatatc ccaaattttg
9661
       atatgtagtg tttttgtttc catttagttc acaatacttt ctaactgccc ttttgatttt
9721
       ttgtttgcac catgggttat ttaaaagtgt attatttagt ttttgcacat ttacaaagtt
9781
       tctatatatt tcttttattg attttgtgat ttaattccac tgttgtcaaa taacacactt
9841
       tgtatgcctt gatccattta acttattgag acttggtttt gggactggaa tatgctctct
9901
       catggtaaat gatccttgta tgcttaagaa gagtgcttat tccagtgctg tcaggtggag
9961
       agttctataa ataacaatta ggtctacttg cttaacaatg gttttcaaat tttctatatc
10021
       tttacagatt tttctgtttc ttctatcaat tactgaaaga agaatattga catctctgac
10081
       tgtaactcaa gatttgtcta tttctccttt catttttatc agttttttcc tcatgtattt
10141
       tgaagctcaa ttattagaag cataaacagt attqttatat tcttcagata aattgaccc
10201
       tttatcatta tggaataact ccctttattc ctggttatat ataatatatg gaataactct
10261
       ctttattcct ggttatacat aatatatgga ataactccct ttattcctgg ttatatataa
10321
       tatatagaat aactctcttt attcctggtt atgtataatt ttagagcaat atataatatg
10381
       catatttttt gtctaatctt attatagcca ctccagccac gttttgattg gcattaatat
10441
       qqaatatctt ttttctttct tttqctttqa aacttqttqt qtctttqtat ttaaaqttqq
10501
       attettgttg geaaaatgtt gttgggtttt cetttttaaa teeaatetga gaatetetag
10561
       cttttagttg gggtgtttgg agatcattta tgttaatgtg attgatagtt tgggcttaaa
10621
       attettetet tgetatttgt tttetatttg teccatetgt tetttgttee teatttatee
10681
       ttttttgtgtc ttattttgga ttacttgttt ttttttataa ttccaaattt tctccattgt
10741
       aagatcgttc tctatagctc tttgttgtga tattgtagtg gtttctctag ggttatctgg
10801
       agttagtata catctttaaa taacacagtc caccttcaag tgatattgta ccaattcact
10861
       gataatttaa gagetttaca acagtgaact tecatttete ttetetggte tttgtgetgt
10921
       tgctgtcata catgttgctt ctacacccct attataaaac ccacattata ttgtcattac
10981
       tttYgcttcg aacaatcaat tatcttttaa ataatttaaa caataagaaa taaaagtctt
11041
       tttatttacc ccatagatgc catttctggt gctctgcatc ccttcctgtg gatcagatct
11101
       ctgtctggta tcagttcttt ctacttggag gacgttcctt tcacatgtct tgtagtactt
11161
       gtctgttact gacaaattgc ttcagctgtt gtatgcctga aaaagttttt atttcgtctt
11221
       cacttttgaa caatattttt attaagtaaa aaattctaag gtgacttttt ctttcagtca
11281
       tttaaagatt ttattctgct tgcattgttt ctgatgagaa atctgatgtc accctctccc
11341
       tagttcctct atagtaacat ggttttttct tttctcactt ttaagagttt ttctttatca
11401
       ctatttttag tgtttgagtt tctgggtact tcagggtttt atacttttca tcaaatttgc
11461
       aatattttgg cgttatttct tcacataact ttttttctgt ccttttgttc actcctttcc
11521
       tcctggaact tcaattatac ttacattaga tgactcgaaa tttccccaca attcacagat
11581
       gctgtatttt tttcagatct gttttctttc tgtattttat tttgcgtagt ttctattcca
11641
       tacaatatat ttttcatctg aaatgtaget tttatctgta gatgttttat ttaggtettt
11701
       ttaaaatctg ccatatctct gcttaacaca atatttcata taggtttttt tttcacatat
11761
       ggaatttgga tataatagct qttttcatgt ccttttctac ttattctatt tatctatgtt
11821
       aattetttige tagttttgat tgattgatte tietetttat taagggtagt attitetget
11881
       cctttttatg cttgaaaatt ttttattgta tgccagacaa tgttaatttt aacatattga
11941
       gtgctgaata tttttctatt tctataaata tttgagctgt gctctgaatt actgttatgt
12001
       tgccctgaaa caagtttgac atttttgggt cttcctttta agtttttttt atgtgtaacc
12061
       agagtgacat ttagcctgtg gttaatttat actactattg agacaaaact catgagttat
12121
       gtcattttcc attttgaata gtqqqcacaq gtactatttc tqaccctqta aqaqttqtqq
12181
       taattgttcc ctctgatcgt tttgggtggt tctttctttg accacagatc acttcctcag
12241
       aagcatgaac tcgtcagtac tgagctgagt aactgaagag ggcgctgcag atatctctgt
12301
       agtgatetee teagtactet gtgaacteaa gteacettag cetttecaga Steecagete
```

```
12361
       tgattcctca tctcggagag accactggac ttcatctctc ttcctcttct ctttgccaga
12421
       gttcagaaac tctccaggca gtaaccaggg caattgccag cctcttttt tgcttcccat
12481
       ctctcaaggt tcaccgtcgt ccactgcctg atgcttatat cttgaaagct gttgattcat
12541
       acgittigtt cagigittag ttatttaagt caggagggta aatcitatic ciattactci
12601
       atcttggctg gaagtagaaa tcacaaaact tgcctctaaa caaagtatat cacaaacatt
12661
       ttctatttat ttaaatatta ttagtctaca tcattttgga tgactacaaa gtattttatt
12721
       atttaaaaaat accatatatt aattaatact ttatttttta tatttcgttt tttcctgcta
12781
       ttgttacaat gttgtgaatg agcagataaa tatttttgaa tatccatgat tatctcaact
12841
       gtattagtag aagagaaatt gctgagttaa ggggtatttt gatgcagatt tccaaattat
12901
       cgtctaaaaa ggttgcaccg atttatatac ttattatctc tccaaatatc tgggtattat
12961
       tattataatt tttttaaacc tccaagcttc ataagtaaaa atggcacatt tattggtttt
13021
       aattagcatt tttatttctt ttgatgtata atattttaat tagtttattg aaaatctctg
13081
       teetteettt tigaattiag titatatett tigeatatti taaagtitag attgitgiet
13141
       tttctgttat taatttataa aatatttgtg ttgagtgtgt gagtgcatgt gcatgcatgt
13201
       gtatgcgtat gtgtgttt gtaacatttt catcagtttt ttgcttgccc tctatttatg
13261
       gtattttcag atgtgattag atttatttgg tctttgacat tatgatttct gtctctatca
13321
       ttgagtttag aaggYcaata ttattcctac ctcattacca aagaaatact taattctatc
13381
       tgcttacaga tttaactttt ttctttttcc ttttaaatct ctaatacatt tgqaatttaa
13441
       ttttcttact gattaaactt tacttgtata tgtttatatt ttataatgat aatattaaac
       catttgtaat gcttaaatta aatattaaat ggtcataaca ccatttattt aaagcctatt
13501
13561
       ctttctttca ggtttgatat actgactttt tcatcatgtt gataatcttg tgatatattc
13621
       ctgttaatgc tccattggtt tctacatttt ttatcctgtt cttataagtt gtctgtctcc
13681
       tcctgtggct gttctatttt attttaatta ctgaaacttt attggacaat agtaagctca
13741
       gaagagaata tetgtaggae aacagtagge teaaggagaa ateatetgtt eteetaatet
13801
       tttactattt ttcgtatttt cctgactgat cttcagtatt tctggaaaat tgacaagcaa
13861
       tgcagatttt ggataatagt gcaggcttca ctttcattgt gcctgggttc aaatcccagc
13921
       ttcttagctg actggcaatg tgacccagca aatttattta ctttggcatg gctcagtttt
13981
       cttatctgta aagtaagggt aatagtagtc cctatattat aagtttatta tgatgattgt
14041
       gtgggtggaa cccgtgtgtg gcacacaccg gcacttagtg agaagttagc tttgccattt
14101
       teetttatte tteaetttaa ttttagaatt attttgteaa etteaaaaaa gtteattatg
14161
       aatettatta etgagtette tetagaagea tgacatgact etteatteae ttgttttett
14221
       ttgcgtctct cgggaagttt tgcattttct gttgtgtata tataatattt gtacattata
14281
       totaactato tatotatota totatotato tatotatota tagatatata tatatacttg
14341
       cacaattata tccacataca ctctaagaaa taaggccagc agaaaatatg tgggtatgtt
14401
       agactataat tatctctgag tattaggtta tgagaatttt ttcatttgta gtattctata
14461
       cattttaaaa cttattaact gagcattact gatataaatc aataattttt aagtgacttt
14521
       tgtgccctac tacgctggtg ctgtgtgctt cagactgaag ggcagtgcct gttagatctt
14581
       atgtgctaca tgttacacaa gtactttggg aaataagctg ttgttttaaa aaatgtgtca
14641
       tcaatttcta gagaactgga cataactttg aagcaaaaca attttaagtg gatgactaaa
14701
       atcettectg aaaacgcatg cetgtggtee etgtgaaate acettttete eccettteet
14761
       aatctccttt ccctcctaaa tctaaatttt cttgtaatgc ttcaaaatat ctggacttga
       tctcagaaga tattctgtca tgcatattga caggactggg tactttcagc aaaacaaagg
14821
14881
       acatetttaa atttttaeca geeccagata gaagtgteta gagttetaga ttttaggaga
14941
       aatacagcca tgaaggaaaa tgttgacatt tatttcaacg tttcaacagg aggaaagaat
15001
       gagaacacaa tqtacaqatc aqcctttcct ctcatqttct tacaatqqqa qaaqqaqaqa
15061
       gggtatttga gacaccaaag catgaaagcc atgacacagg gaaaatactg agaacataac
15121
       ctacagatca gattttcctt ttatgctctt accatgtcat aaggcaaagt gcttgtttga
15181
       gatttcaggg cacaagagcc actacagcag gaaggcgtcg catcatcgga acatttgtga
15241
       tcaaagagaa gactgtctgc acaaagcacg gactgcagcc aaggggtcct tcacccactg
15301
       tteteacaaa tqettqeett tqetateaca actqetattq ecaqetqqae qetqaqtqte
15361
       ccctcagggt tcatagagat ctgtctccct gaggtctctt aggtgctgct gctgaccagt
15421
       agtaacttgg tgaagaccct cgattgtgag gacattgtgg tgaaatgttg ctttcttgac
15481
       aaaacaaatg tgtttccttc tgatccatga acaaagatct gttgaataaa tattatgagc
15541
       gtgtagaagg gagagacttt atctgatcag actagggaaa gggtgctttt aatatctgtg
15601
       gaacaatctg taagatcgga cgtttcctct gataacagaa actccagagt tacaagagaa
15661
       atggcttcct gacatcacag acactaaaaq gccttctgag taggaqtaqc aqttqaattt
15721
       15781
       catttettac tagcaagtac ccctgactte tacgtatttg ttgttagate cacaaagaat
15841
       atagtataac ttcattttct atcaactatt agggatctat agtgccataa caaagtactc
15901
       taaaatttat cagcttaaac agtaaacctt tattatctca catggttcct gagggtcagg
15961
       aatctgacag tgacctagct tggtgattct gcctcggggg tgctcatgaa gttgtggtca
16021
       agotgttgac tacacttttt catctcggaa gacttgactg caactgaggg acttgctgct
16081
       agggtcactc atgtggctgt tggcaggagg cttcaggacc tgtctgtgtg gagcactctc
16141
       tggggctgcc ctcgagatgg cttctctcag agggggtgat tcaagagaca gagtcccaga
```

```
16201
       caagtacagg atgccttaca taacctaatc ttggaaatgg cacgtgatca cttattccat
16261
       attatattga tgacatagac ctgacctggt acagtgaggg aaaagactac acaaggatat
16321
       aaacaccagg atgcagagat cagggggcca tcttggaggc tgcctgccac agctattaaa
16381
       ggtaatttgt tgacaactca caatttctag ctacccagtt gctcattctt ttgcttaaac
16441
       tattcaaggc tggtggggga catgtgaatc ctgttaaaat gtggaaagat ctaaatgtgt
16501
       tttatagaaa aaattgagta aacttttttt gcttaataag caaaagaatg atgaggaagg
16561
       aaggggcaat tgtgtgtgtg tgtNagagag agagagagag agagaatata aaaatctaat
16621
       tttttaccaa aattatttt tttcttccgg gtgaacatgt ggctttaaca taacttctat
16681
       tccacaaact caaaatcaag cacctctgtt aactcttaac taaacataga ctttttStga
16741
       actttggaat aggagtaagg gggaagagtg aaacacacta gctttcatgc cttgtttaat
16801
       ttctaatttc ttaaagagtg gacaagaaag agaatgaatt cagcaaagtc tttgatccct
16861
       acagttcata ttcagacaaa taggagccaa ttattaaggc taactgcctc agtaaaataa
16921
       aaaacagcca atttaacttc tgaacacagg aaggtttttt acccattatg gaatcatgaa
16981
       agatctgaat gcacatgata tggtgtaacc ataaaataag gttgagtttt tgcttcctgt
17041
       taaatctcat gagaacaagc attctgagtg aggggtgcgg taaagacaca cagggctagt
17101
       cttaaaagaa ggtaaagttt ctggctatgt attaggcttg attagaattt tctggttaaa
17161
       acgaattctc cttcagtttt gtttctttca tagatgctgg aattcctata aaagccattc
17221
       agacatttat atttatgact tttctacttt ttaaaaaact atactcctcg agagaataaa
17281
       ggagtaggtt ttactagaca ccatatttaa tgggttaaag caaccaagta agaagaggga
17341
       atgttctaga agtgagggaa aggggattca tgcagacatg aaagtgaagc aaaggtaaat
17401
       17461
       cttccttcct tcctccctcc ctccctccct ccctcccct ctctctctt
17521
       ctttctttct ttctttcttt ctttctttct ttctttcttt ctttctttct ttctttctt
17581
       17641
       egeceagget ggagtgeaat ggeaggatet tggeteaetg egacetetge eteceaggtt
17701
       caaqtqattc ttctqcctca qtctcctqaq taqctqqqat aacaqqcatc tqccaccacq
17761
       cccggctcat ttttgtattt ttagtaaaga cgggatttca ccatcttggt caggctggtc
17821
       ttgaactcct gacctcaggt tatccaccca cctcggcctc ccaaagtgct ggcattgcag
17881
       gcatgagcca ccatgcctgg actatctgcc ttctttctaa aagagataaa gctttctagc
17941
       tgcctagtac cttggtgaat ttgggctgag gccattcagc gacaagcact ggaaagaatt
18001
       gaagtgctcc caagtacctc tgcaagggaa gaggggtgat gagcagggga gaggagaga
18061
       ggaagaggac ctggtggatg gtaggaggcc atcaggtctt ccatgtgatg tgcgaatctt
18121
       tagctaggta caccgtattt tagaaatgtt tggtgtaaat ggatttettt tgcaaactgg
18181
       agaggggact gagttttggg ctgatgctga gtggaacaga gatgggactc ctggagcttg
18241
       agacacagga tettggagaa aatgageee agaaateage agatgteatg aggggaetet
18301
       gggttatcat aagctgtgga attggtatcg gatctgaact tgtttcttca aagtgcaagg
18361
       aagaccaget aatetettgg ttttgccaat ttagggcaat tttcaacatt acgaaaatte
18421
       atttagtttq qqtqcaqaqc tcaqatqtcc tctccatqtq gacacaaccc tacatqtqca
18481
       cacacaaaca agcacttgag cacacatgtg tgtgcacaaa ccccactttt ctgcttttcc
18541
       aggagcaccg tatgttattt cagacaaata agtttggcca actttgatct ttccattcag
18601
       gtcactgatt gctgaattgc cacagatgtt gttctgccct ccttaggttc tagggcattt
       tcattaagga gagttacctc caagaggggt catggcaggc gagcaaacca tacttggggt
18661
18721
       gcaccattaa agacctccac ccctggggag cgatcctcac agcggagcag gacttggccc
18781
       cttacaagtt tgcctagtct ctccctgaca ggtgggtaca gcagcaggag tctgaattcc
18841
       acattttgca qaqccctqqc ataqacqqat cattcaaqaq aaacttcctt aaqctqccca
18901
       cggctgctca ggtcctcatg tctgcattca gatgtggcct tctttattac taactcagga
18961
       ggaccttgat ggatgttagt ctccctggat ttgaaaacta tttatgggag cactgaaaat
19021
       ggcattattc gttttgatgg caatcaagta cataaatgcc aatgatagag actgaaggat
19081
       aagttccaca gcggatgtgg ttcaagtttc taaatactgc tgccatgatt ctcatgtgat
19141
       gtgaaggata ttaagagate tagaactagg atagaactgt ageteaacea tgetagaaaa
19201
       aaattttttt aagttccact ctatttaaaa atatttgtgt agagacaaaa tctcactgtg
19261
       ttgcccaggc tggtctcaaa ctcctggcct caagtgatcc tcctgctttg gcctcccaaa
19321
       qtqccaqaaq ccactqtqcc aqqccaaaaa qttatactcc tqaqtaaaqa ctactttata
19381
       gtctcttgat aacctaccat aatttaaaga atattctgat tgctgtaaat ttctgtataa
       tttaaatctt aaggeegggg geagtggete atgeetgtaa teetggeatt ttggggagee
19441
19501
       aaggagggag gatcttctga ggtcaggagt tcgagaccag cctgaccaaa atggagaaac
19561
       tccatctcta ctaaaaatac aaaattagct ggatgtggtg gcaggtgcct gtaatcccag
19621
       ctactaagga ggctgaggca ggagaatcac ttgaacccgg gaggtggagg ttgtggtgag
19681
       ctgagattgt gccattgcac tccagcctgg gcaacaagag tgaaactcca tctcaaaaaa
19741
       aaaaccaaaa aatcttaaat ttgcagattt gtaaaatcct acatccctgt ccccacctct
       cttttcccat gatctttagt ttagttcctc aaggtcagga atgagtctca tcttctcttg
19801
19861
       cacctctcac agtgcctgac acaatattgg ggacattgat aagtattgtt ttgtgtatta
19921
       cttaggcata ttttctgtgg ttctctagca gtggagaata aatgattgat gatcttatga
19981
       agtcatagtc cacctaaaga ctgcgtagta cgcaaatgag tctaccatta tctcctcgat
```

```
qtaqattttc atcttttatq tatqqtqact ctaqqtaaqg agaagaggtg atctaqctca
20101
       cttqqtqqqa cgagcgtctt gacacaaatc cacaaaaaaa ggagagtaca cacagctgtt
20161
       catttgtctg tgtgttgagg tgggggtggg ggaggtgagg ttgtattcac acagagaaga
20221
       catcttagct ctgcacccaa accaaaacga gcgtcagtga tgttaaagat tgaaggtcag
20281
       gtgctaccca ggttttgtta gtctcttgac ataaacatgg tgtctagata tggtacatat
20341
       tatgaatcat tccatctttt caaatacatt ttgaagggtt tttttgtttg tttgttgtt
20401
       tttqaqacaq aqtttcqctq tqtcaccaaq qctqqaqcac cqtqqtqcaq tcttqqctca
20461
       ctgcaacctt tgactcccgg gttcaagcaa ttctcgtgcc tcagcctcct gagtagcttg
20521
       gattagaggt gtgcaccatc acacgcagca aatttttgta ttttagtaga gacggggttt
       caccatgttg gccaggatgt tgggattaca ggcgtgagct accatgccct gccagaaggt
20581
20641
       ttgtttttat tttaaacaca attgtgctgg gtgtgatagg atgggtatat gatagagcag
20701
       aaaatcacta gtatttcatc aatatggata gcatcctcca taggttactg atttaatatt
20761
       gacaacaata cccttcacac gagagagagg ttagagaatt tgcccaaagt caagcagctg
20821
       ggaagaggaa aactaggctg tgctctccct aaatctcatg ctctattata ttagggaaga
20881
       ctctggcaga tggcacatcc tgattatttg aaagtcacca atatttttag aaattgcata
20941
       gataattaag gttagtttat ccagaattct aataattgca gcacatgtcc ataaatctct
21001
       attacaacca cctgagatag gccatctcgg gcatgtggaa aaagcccagg tatagacact
21061
       ttggatcctg gatttccctt gtactggctg ctaccttggg taaggtaatt ctcctctctg
21121
       aageteetgt cagettetga ggattgegtg ggatageaca taccaaatta etgeacacag
21181
       agcctggtgt acacttaagc acacgagatg tgtcaaagtt ttcaaaaaaca ttgccaacga
21241
       ggcatcagtt acaaaacttg ctgcagagtg agctgatatt gtgccactgc actccagcct
21301
       21361
       aataagcaac ttgctgcaga aatgggtact cttgttctag aaatgtgact atagggaagt
21421
       tacaactacc aactegegtt aagggaaatg agtgacetge cacctacatg gtgttaggga
21481
       ggttttgctg agaaagtcac tcatgaagaa ggcaaaatat agttaagaca aaatgtaact
       atctatagag ataaggtaaa aattggaaat agaacttcat taaagatctt tcaaataggg
21541
21601
       agaatgtggt gaaaactgca gttaacattt gttaacagtg tgatcatcgg gttcagctta
21661
       tcagtaacct ggttcctgtc tcttaactga taaagaaaat gggaggtttt taaagagagg
21721
       ctqqctqttq tatttagtaa agctataaag ctgtaagaga aattggcttt ctgagttgtg
21781
       aaactgtggg cagaaagttg aggaagaaag aactcaagta caacccaatg agggtaagtg
21841
       gctttggggt atttttcaaa aatactttta tctcaaaggg aacaaaattt tcacatacga
21901
       atttaggtca ttataacaat aatattcatt gttatattat tataacatta tatgttataa
21961
       taatatataa tatataata atataattgt tataacatta taacaattat atattatata
       taatacaatt atataatat tattatataa ttgtaatata taatataatt atataatata
22021
22081
       tattatataa tataatatat aatatatcat atatgttata tattttatta tataatatat
22141
       attatatata atattatata taatatatat tatatataat attatatata atatatatta
22201
       tatataatat atttatatat attatatata atatatatta tatattaaat attatatata
22261
       22321
       tataatatta tatttatatt gaggaaaaca ttcactcaat gacagaggga agacttagct
22381
       ttccaaacaa tcttctgtta aaagaaaaac tttagctgaa ttaaatttaa aggagtttaa
22441
       ttgtgttttg gtgccttatg caaatgaagg gcctataata taatatata tgtaatatat
22501
       ataacattat aacaataata atcattgtta tcaattttgc ttgtgagaaa agtagtatta
22561
       ccaatgggaa aatcatggta cagattgatg aaatcccttg aaatcttgtt tgtgtaatat
22621
       gggactttgg tccaacgtgt tggaaaagac ctggttgagg gtaagaaagg atgggagctt
22681
       tatagtatac acacagagtg ctcatattct ggagttattc tttctgaaat ccacaaaatc
22741
       cagcagtgta atctactcag atggagatga aggaaaaaaa caaaacaaat gaagcaagaa
22801
       gctgctggcc aaaaggggtg aggaagcagg gtcataccca tctgcaagaa gaaaatagag
22861
       aacggacatg agaggggtgg atatgaggat atgagaaggc cctcttgagt gagggccatt
22921
       gtttctggag gggtggggag tgaatgactg atgtaagttc aagtttgttg cgggtgcctg
       ctttgatcct ctgcttggtg gctccattct tcttctctct gaaggctact tcaacagagg
22981
23041
       gtgcagaggt aaacatagca gaactggccc ccaaagcttt atccactgaa tgtcagtttt
23101
       teetttaatt tetteeagge ataaggaaat eggaggeatg tttetgtetg getataacag
23161
       ttctcctact agatcctgag gccaaactat tcacatgcca ttgttcctga ataccattgt
23221
       ggacataatc atctaatttt ggctattcca tgcagaggtt tggagacaga aagctctgga
23281
       ctaatgcaga tetgcattta atecetgete ttacaettae ttgttgcatg acettaagca
23341
       acticational geotteaaat oftigagettic tegettacaa agaggegaco agoatattag
23401
       taataatatc atgtttagag ggtggtttct gaaggttaaa tgggtgtaag tgtttagtgc
23461
       attactcage cagcagtcaa cacttaatga tgtcagacet catcattgte ettattttt
23521
       ttgttccaca ctctgggaac ttttatctct gcgccctgta gatatgctgc tgtgactgct
23581
       gacttgaatg ttacgggatc cccagcagga tctgggattc tctctgttct tgaatgcgtt
23641
       gtttaggtga ggagattgat tccaagttac ttttctactt gggtgcctct gtgtttggag
23701
       atgtggccgc atagctcaag tgtatctggg gtgtcttcag ttttcggata cctgtttttc
23761
       ctctaggaga attcttcctc tatgtccttt aaaatatttg tcctataatc acaagggttt
23821
       atatttetet geceaeteet ecataettee tggttttetg ggagtgeaga ggeagacaae
```

```
23881
       tgcaagggag agccccRcaa ggatttgggt gatgactttg agccaggctg acatccatgt
23941
       ccaagggcaa atgcagtctg gttgggagga tacaatgcag gagtcttaca acagtggcga
24001
       tgccaacatt gtgggagcat gcatgtgtgt gtgcacgagt gtgtgtgtgt gcgtgcRctt
24061
       gcatgcgcgt gcatgtgcag tagggtccta ctacaggttg gtggaaaaga tccctctggt
24121
       tctagaacaa aaatctaaca gcaaccaaga gtccagttcc aagaaagaga aaggaaaggg
24181
       ctagttaggc agggcgtctg gattctgaat aggggctgag gccttggctc agacacagag
24241
       gagaatggag aaatggggtg atgacagaga catctttaaa atatttgtgg gccaagggta
24301
       ggatggcttc aacttgtctg ggatgtattt gtccttagca cgcttggaat aacatagcag
24361
       cagcagcagc agtcgacact tactaagact gctatgtgcc agaccctgag acaagcactt
24421
       tocacacacc acatectgcc attettatga tgatagtgct gtgaagtagg gatgaaacca
24481
       agtccttgct gactccaatc ttgtacttcc aaccaatgct tgtgagggcc ccaggcccc
24541
       24601
       gagttgggtg tgtctgagcc ctgggtagaa agggatgcta tcctaatgaa aaaacacaga
24661
       ataaagatag tgtccaccaa gatgtaactg agatttaaag aagacttgat aacttatcag
24721
       ttgggtgggt agtcttgaat ttcaacaact gcagatccta tctaagacca cttttgtctt
24781
       gtgacaactt agttaaacat gactttagtg tggatatcag tggccaattt tgagcccatt
24841
       ctatgattgg aacaaaggct tccctgtcat aggtgagtga caactatctg gtctggtggt
24901
       acaggggtaa aagaatttac taagacagtt gtagataaag aaaggcagat ttattagaga
24961
       aagtatgaaa atacattaca aggttgcaat gggcagcaca gccagagagg agctgactgc
25021
       aaagaaacaa aggcttgctg gatatttata ggatagttcc tgggctccag ggggctatat
25081
       gcagtactga taatgccaag gttgtagtga gctaacttgc aggtgtctgg tgatagttgg
25141
       gtgcYggaag attctgagtt atttgctcag aagagctgtg tgtcctgggc catgaagaaa
25201
       ggcaaactta cagtttatct gctttctgtc tttgctttcc cttggtactg ctagtctgac
25261
       ttttttttcc ctaattagga ttccgcacat cctatgcctK tctctcagga gccttgcagt
25321
       caaagtgtga tggagtgtct gagggccgag tcattgagta atggggcctc tgccctctca
25381
       gccccactgc cacccaactg aggcttcctg gatgagggtg tatgccgtgg ggaaaaggag
25441
       tgggatgatg tggatccccc tggggttgca ggtcatcctc ccaaatcccc actctccctg
25501
       totacceaca ggaccagtca gcatttetet aatteetett cactetgggg cattatgate
25561
       agaaaatatc atgtaccttt tgctgcaatc ttggctgaaa gaacctcaca ctatacttga
25621
       atgtctttga tttaccatct gagcagagat ctcaaactat aaagaaaatc agagtctctt
25681
       gatgcccatg ttcaggacaa gttttcttgg atttcatgac tctttaggaa cagtgagctt
25741
       atgggtaaag accgagactt tagtgaggaa ttccctcatc atttttctct tcttgagcaa
       aaccttgtct gatttggggt tgtaatgtgc aaaattattt tcttttcatt tcaccttttt
25801
25861
       ttettaagta gtggtttetg gaaatetgag etgetttgtg caccetgaet agtetgtgee
25921
       agatcacgaa cgtacacacc aactetcaag acaccagaaa aatcaccett tagaggaatt
25981
       atctagaaaa ctagccccta tgcaaagtga ataacctttt caaaatataa tctacattaa
26041
       gacttttgaa actaacacat tcctcaatca tagcaaccaa atgaattact taaagtgggt
26101
       ttggttcctg ctttcattaa gggagtattt tcaggaaatt agaggtctaa tgtaaacatg
26161
       gateetttag gttttteeat tttetttgte tteaggetgt ttgetgeeae ttteateete
26221
       cccttatctc agtctgccag tgatggtttg tgcctctggt cKctgggtca gatccaggac
26281
       taattcactc acctctgatt tctagttcca cMcttatgac gaaagcattc ttaaatctga
26341
       tattcacata aaaaaggatt tgattgtctt tggcatgtct ctgcagttag ttaatcatgt
26401
       cttgagcaca gaaaaatggg aagaaaatgg aaataagatt tctttgatag gaggagtcct
26461
       gctaagactt gtctgtccct gagctgagat atgctaattt accttaaatc tttccataaa
26521
       ccacattctg aattccagtt tataaatctt tatcttattt cccgaacctc atctcatact
26581
       ttttatgtRa tctggtgggc atagatattc tcttggtgaa cctgaggaaa tctagaatga
26641
       caaaggatct aaaataagta acacagttgc tcaagttcct gggccctatg gtagaatttc
26701
       cttcttcaga gaaggggaga acaaggtaga taaagaaagg agcatttcat cttgcagcag
26761
       cttggaagat agctgggagt ttaattttta taaaaaagaa agcaatcaaa ccatccaata
26821
       cageccattt teaaccaaaa gtattteaga attacetgga gatttateta aaattatgta
26881
       aaaattacct agaattacct aaaaaaattg tacaatttca tttcccttct gttacatgat
26941
       atggataccc agctactatt tattagtagt agatacagac tatttaagag acaaaatatt
27001
       atattccaaa aaqtqtqacc cttaqaaacc aaaattccaa aqaaqaqttc ttgtaattgg
27061
       caaggctggc tctgattgag tgtcaaactt cctgtacaat gatggatcaa cttagaaatt
27121
       tgatagattc aaagcattct gaaattgtac aactcgctca gcacttgtat tatttacatg
27181
        tcttctgtta ttttccattq aatgatgttt tcatgtatac tcagctgctt ataagggcaa
27241
       tattttaaaa tcagagcaat ttctttaaaa ttaaaatgtt tcagcccagc atggtggctc
27301
       aYgcctgtaa ttccagcact ttgggaggcc gaggcaggtg gatcatgagg tcaggag\tc
27361
       gagaccatcc tggctaacac ggtgaaaccc catctctagt aaaaatacaa aaaattaact
27421
       gggcgtggtg gtgggcgcct gtagtcccag Stactcggga ggctgaggca ggagaatggc
27481
       gtgaacccag gaggtggagc ttgcagtaag cccagatggc gccaccgcac tccagcctgg
27541
       gtgacagagc cRgactctgt ctcaaaaaaaa aaagtttcag caaaatactt tttaaaatgc
27601
       actgtaatta ccctaacaat attacactgt tacagatagt ttaagaataa taaaaaatgt
27661
       aattcataaa ctcactcctt taatacaaca aacctgatta atqttctqtg ttccttatta
```

```
27721
       gactttctcc catgtatagc tttacaaagt ggtagttgat gagtaagtac aatttgtgat
27781
       attittitca cgigagaaca aactcaatat gicaattact icatagatit tiaattiata
27841
       aaaattttct catatctctt ataaaaataa aaaaagattt gatttcagta aacacacatt
27901
       tggagagaaa gttgatcctc atataaactc aaccatagac cattcagtgt gttgcaagtt
27961
       ataaatgcca tggaggaaaa aaatcaagca gagttaggaa gattgggagc acagggttag
28021
       ttggtcaaag ggtgggagag gatagacctc attaagaaag tggcagacct cattgagaaa
28081
       gtgagggagt gagtcacaca gacacctgga ggaagaaaat gcaagagagg gtaaacagtt
28141
       gctgcaacac cctgaggtgg gagcaggctc gggaggccaa gcaacaggaa ggacaaaagt
28201
       gcacgtgttg tgtagttaag attgggggtg tagcaggatg taaatcagag aggtgagagg
28261
       tgacagaccc ccagggggtc ttcttgttca ttgcaagcaa tttggctttt gttgggtaaa
28321
       gcagaggttt gaagagagga gtaatttgaa aggatcactc tagcttctgc tttgaaaatc
28381
       aattataggg gacaagggaa ggatcagtgc aaagactagt gcaataattc aggcaagaga
28441
       tggtggtggc ttggatcagc aaagatggcc atgagaagtg gtcacatggc tgagttctgg
28501
       ctgtatttgg aaagtaggtt cactgggact tattggtggc ttgagattga tgtgctggag
28561
       aaagagaggt ataaggatac ctccaaggat tttgatatgt gcagctgaaa aaagatgaag
28621
       ttgccatcaa ccaagatgag aatgatggtt ggaagagcag attttgggag aagaacaggg
28681
       attcattctt gacaggttat ttttgtgatg tctggaatat agacaagtga aggcattgag
28741
       tcagcaggtg gacatgagaa cctgaagttc aggggaaaag tgtggacggc aggtgtaact
28801
       gcaagagtgt aatcagtata gagatggtgt ttcaaaccag ttaactggat gacatcatgt
28861
       ggggagtgag tgtggacaga gaagaaaagt gcttcaagga ctgaaacttg aggtggtgga
28921
       gagaagagga gatgtgaagc agtagcctgt gggtcctgaa gaaagatgtt ggagaaattg
28981
       ttccactttg acagaggctg cttatagctc aaggtaaggg gaagacgcag gactgaggat
29041
       tcctaagcaa catggtggta attgacgata ttgataacag cataaagaca tcataattta
29101
       aactaattat taacaaaatt gaaatttaaa gaaacaagta atgcaatgtt tccaaagtgg
29161
       attcaaattc cataatatta ttgcaaaaca aacaaaccaa acaaacaagg aaaacaaaga
29221
       aagaagggat totaacataa aagaatttaa aaataagtaa atcagotogg aaaactcago
29281
       aggetatate treetggtgg tragaaatat etgettatta aaaggtttea aaggteettt
29341
       gggaatagaa cttattgcaa tgtgtttggc tgtacatttc ttacactttt ttttttaact
29401
       gaggagtcaa ttaaaatatc ctgaagtgac tgtccataga acacattttg aaaacacaaa
29461
       tgctatataa tcacaaatat atctccctcc cctttgctag tgattgttac caaactgcct
29521
       ctgctagcta ggattcaact ctttctctta acaactatgc tttgatggaa caatttttt
29581
       agatacaaat attgtcaatg agaacaatta ttatacagtg cagtcacaga agagaaacag
29641
       aaaagaaagg gaggtgcaac aaaatgaagg taaatattag ccttcacact taaatacatt
29701
       tgatgagggg ataaaatgta catgtaggtg tggatttatg ggctgtgttt ttcaaacaat
29761
       taccaaatat ttattaacca ccaactcagt gtaagctagt acactaggtg cctatagtga
29821
       tecageagge tataaggeag gatttetgee ettatagaat tttggggeat atteteagta
29881
       tettttgtat etetegeeet ettttettt ttttgtgaca eaagatgtea ttetgteate
29941
       caggetgtag tgeagtgaca tgateatgge teattgeage ettgacetee caggeteaag
       cgatcctctt gcctcagcct cccaagaagc tgggaccatg ggcatgtgcc atcatggcag
30001
30061
       actaattttt aaattttttg tagagatggg gacttactat gttgcccagg ctagtgttga
30121
       actectgggc tecageagte eteceacete ageeteecaa agtgetggga ttactggtgt
30181
       gaaccatcac teccagecet ttettttget ttgttatttt aaaaactttt etttattete
30241
       tttctgttac cttttaataa actataagaa gtctttcaaa acaaaaatgt cttaatgtca
30301
       gtgggttagg ttttgtcctg cattattttc ttttacccca tggtggggaa aaaagcaaaa
30361
       aactqtttgg tgcctataca atgccttatc gcacaqcttt ggtttgtgtt tactccagtg
30421
       tegtetttaa ggtggtgagg teagtgeage tggcetgggt cagaacteag ggteacttea
30481
       tgccagatgc tcaccacctc tgcagtgttg tatgcagaga tccaaattca tccttcaggg
30541
       aaggatgaac acttattctc tgggatgttt tacagtgaaa atcagaagtt gctctattgt
30601
       geocteatgg cetagetact tecactettt tgaagatett acetggtete atgggtttaa
30661
       tttcatctat acagtgacat atcccaaatt tgtatctcta gcccaatatg tctcctgaat
30721
       ttcagaagca Ratattcatc ctactgatgg gttcacatct ccagttttat gtctaatgag
30781
       gtgtcaagct gaatatggcc aaaactagtg taatcctact ctaaaaccta taccacctgc
30841
       catttectaa tteaqtaact aqeaacecta tttqaccaqt tactcaqqqc eccaaacete
30901
       tottgactgc totcatatcc tgtccatcaa gaagtcctac tagttctacc ctcaaaatct
30961
       atotgcatco caatcatgto toaccacotg cacttocoog cootggtoat agocacogto
31021
        totgtotcac otggatacot catecottgo cattottaag agtgttttca cattoagagt
31081
       gtttaatacc tacgtcaagt ttgaggtcat gtcactcctt agcttgaaac actccagtgg
31141
        tetteateae aettaataaa ageeaaggtg eteagggtga eteactegea atetagtete
31201
        tacccctctc caaccagctc tcactgctct ggagttcctc atgccctgtt ggccatgtgg
31261
       cttccctact ggccctcgga cacaccaggc ttgcctccat ggctggactt tcatgctgcc
31321
        tgccctttcg acctgggctg tttattaaat gtctccttcc tagttggcct tcccaggaca
31381
        totggcotga aattgctctc ctcctgctgt atttttctcc ttaagtggat cacactctaa
31441
       gatettttgt aacttttttt tatttettta tgttaettat tgteeteetg taggacatga
31501
       attccacaag ggcaggagat ttttgtttcc ctttgctctg tgtggtgcat tgctgtatcc
```

```
ccaqcatctq aaacaqtqat tqqcatqaaa qaataaccct cactqqttqt ctqqqttcaq
31621
       tattttgctt ttctaagctc agttgatgca gaacaactgg ttttctgggt tcagtatttg
31681
       cttttctaag cccagttgat gcagtgttgg tatgtgcggc catgttagat gttagaacac
31741
       tgacatcctt accactgatg agtcaatagc cactgcccc gaccctcctg atattttatg
31801
       gtccgagtaa ttactagatg catctacatt taaggctgca aacatgaatt tactggagga
31861
       agcacagtgg tggtggatga cccaatctct ctctcaaact taccaaaatc accaaggaat
31921
       cataaatcca ccctctttg catcttccaa gcaaaqgaaa ttgtagaaaa cacaagtgaa
31981
       actgtttcac agagatatca gcaatacata gcaaatttga gatgatggag attaagagga
32041
       aaattaaatc attottaagt ggtcaaggta aaaacattat tgtgYggttc gctatagtta
32101
       catttaaaac agtccgaagt caaacaactc tagtctgaYt agtcttgtca ttgatatgtg
32161
       ctgtcttaga attctccaga gagacagagc taataggaga ttatctatct atgtatctat
       gtatctacct aatctatcta tactctctct atctatttta aggaactgac tcatgagatt
32221
32281
       gtggggcctg gcagtctaaa agctacaagg caggctggca gactggaaat tcagataaga
32341
       gttggtgttg tagtettgae tetgaattee ataaggeage cagteggaat tteaggeggg
32401
       gtttctgtgt tgcagtcttg aggcagaatt ccttcttctt tgggaaacct cagtctttgc
32461
       totteagged tretgetget cacatteeac agtgetgted tretggeeat cetteatget
32521
       tttgtgttgc ctgtcccatt gcccttaatc ttatatggaa aaatggaaat aaattcagag
32581
       cggaaactta tctaaaagtt ttgagggaaa aatgatcgaa accagaatta ttcacaggac
32641
       aacttgttct atgaataatt tttgaaaagg aaccatttcc tctacagctg gcttattatg
32701
       aggagettga tgataaaata ttggcaagge etceagtgga catttategt gttgtgacag
32761
       atcccagaaa tgaatctggg tctcccgact tcaatacagt tctcttctta ttacataatg
32821
       agagaatatg attttctact cttttctttt actttacttg ttgagaaaca actaatattc
       aattaaaagc atcaatctta catattcagt tgaatgagat tcagcaatca tatcccccat
32881
32941
       gtatcatcat ccatatcacc atcccaacag cacagaaagc atttcttctt cccagagagt
33001
       tgccctcaca cctttccaat caaatttcca catactctct agtttccagc atgatagatt
33061
       ataatttgcc tattgttggg cttcctatag gtatgatatg taatatgtag tcttctgtgt
33121
       ttagtttett ttgettagea taatttttea ggageaagge tttattattt ttttgtttgg
33181
       aaaYgaactc agagttacat aacatgatgt ttttgacagt catcctatgg ttatatcagt
33241
       atttccttcc ttttcattgc tgagtagttt acattaaaaa aaggatccat tctcctgttg
33301
       atagacattt ggattatgtt taatttttgg tattatgagt atggcattta tgaacattct
33361
       tttacaagtg ttttgtggat atgtttgtta attctcttgg gtaagggggg agttattgag
33421
       ccatacagta gatgttcttt aactttttgt taggttttat atatatatat atatatata
33481
       atagtgtgtg tgtgtgtgt tgtgtatata tatatggtgt atataattta tatacactaa
33541
       aatacacaca tttttaagtg taaagttcaa taagttttga tacatgtata ctccattggc
33601
       tttgttttgt attgctataa aataatacct gaggctgggt aatttattaa gaaaaagttt
33661
       atttggctca tgattttaat ggctgcaaag ttcaagactg ggcaactgca tctggtgagg
33721
       gcctcaggct gcttccaatc ctggtggaag gttaagggga ggtggcttgt gcagagatca
33781
       catgggagag aagaagcaag gggtggggga gtgctaagtt ctttttaaca gccagctccc
33841
       atgggaagta atagagtgag aactcaccac caagggaggg gattgaagga gggaattaat
33901
       ctattcatga gggatccacc cccatgaccc aaatgctccc ccttaagccc caccttcaat
33961
       attggggatc aaatttcaac atgaggtgag gagggtacaa atatccatgc ataccacctc
34021
       tgaaaccatc atctcaattg acatattcca caaatctctc tcatactcct ttataagtaa
34081
       ttgctaccct tcatcttcca ttctaggtac ccactcactg gccttgcagt ttgggaacat
34141
        tatataaatg ggacgatata atatgcattc ttttaaaact ggcttctttc attcagtatg
34201
       tttctaagat tcatcaqtag ttctttttt tttttttgct gagtagtgtt ccattgtatg
34261
       aatatagaat aqtttaqcca tatatqtqct gacggtcatt ggtcttgttt ccagtatttq
34321
       gctatatgaa ttgaactggt atgaacattt gtgtacaagt ctttttgtgt gcataagttt
34381
        ttatttctct taggtaaatc cccaggaaga gaatttctgg agtgtaacac aagcacataa
34441
       tcagaaactg ccaaactatg ttcactgtat catgttacac tccatcaatg atgcatgaga
34501
       gttctaggtg tgccacatcc ttgccaaaat ttggtattgt cagtcttttt tattaqctat
34561
       gaaatagaat ctcgatattg tggagccaag atggccaaat aggaacagct acagtctaca
34621
       gctcccagcg tgagcgacac agaagacagg tgatttctgc atttccaact gaggtactgg
34681
       gttcatctca ctggggcgtg tcggatagtg ggtgcaggac agtgggtgca gagcactgag
34741
       cqtqaqccaa aqcaqqqcqa qqcatcqctt cacccaqqaa qtqcaaqqqq tcaqqqaatt
34801
       ccctttccta gtcaaagaaa ggggtgacag atggcacctg gaaaatcggg tcactcccac
34861
       cctaatactg tgcttttcca atggtcttag caaatggcac accaggagat tatatcctgc
34921
        acctggcttg gagggtccta cacccgtgga gcctcgctcg ttgctagcag agcagtctga
34981
       gatcaaactg caagttggca gcgaggctgt gggaggcgtg cccaccattg ctgaggcttg
35041
        agtgggtaaa caaagcagcc gggaagctcg aactgggtag agcccactgc agctcaagga
35101
       ggcctgcctg cctctgtaga ctgcacctct gggggcaggg catagctaaa caaaaggcag
       cagaaacctc tgcagactta aatgaacctg tctgacagct ttgaagagag tagtggttct
35161
35221
        cccaqcacqc aqctqqaqat ctqaqaacqq acaqactqcc tcctcaaqta qqtccctqac
35281
        ccccgagtag cctaactggg aggcacccc cagtaggggg cagtctgaca cctcacacag
35341
       ctgggtactc ctctgagaca aaacttccag aggaacgatc aggcagcaac atttgctgtt
```

35401				-	caggcaaaca	
35461					cctgacagtt	-
35521	ctaacaaaca	gaaacgacat	ccacaccaaa	accccatctg	tatgtcatca	tcatcaaaga
35581	ccaaaggtag	ataaaaccac	aaagatgggg	aaaaaacaga	gcagaaaaac	tggaaactct
35641	aaaaatcaga	gcacccctcc	tcctccagag	gaacacagct	cctcaccagc	aatggaacaa
35701	agctggatgg	agaatgactt	tgacgagttg	agagaagaag	gcttcagatg	atcaaactac
35761	accgagctaa	aggaggaagt	tcgaacccat	ggcaaagaag	ttaaaaacct	tgaaaaaaaa
35821	ttagatgaat	ggctaactag	aataaccaat	gcagagaagt	ccttaaagga	cctgatggag
35881	ctgaaaacca	tggcacgaga	actacgtgat	gaatgcgcaa	gcctcagtag	ccgatttgat
35941	caactggaag	aaagggtatc	agtgacggaa	gatcaaatga	atgaaacgaa	gtgagaagag
36001	aagtttagag	aaagtagaat	aaaaggaaac	gaacaaagtc	tccaagaaat	atgggactat
36061	gtgaaaagac	caaatctaca	tctgattggt	atacctgaaa	gtgacaggta	gaatggaacc
36121					tccccaatct	
36181					gatactcctt	
36241					tgaaggcaaa	
36301	_	_	_		ccatcagact	
36361					caatattcaa	
36421					taagcttcat	
36481					ttgtcaccac	
36541	_	_			acaaccggta	
36601					gaaaatgcat	
36661					cacacataac	_
36721					gactggcaaa	
36781					tcacatgccg	
36841					atggaaaaca	
36901	-			_	ccaacaaaga	
36961					-	
37021					caagaagagc	
37021		-		-	aagcaagtcc tttaacaccc	
37141		_				
37201					caggagttga	
37261					ccaaaatcaa	
		-		_	cacatagttg	-
37321					tctctcttag	
37381					cgctcaacta	
37441					atgaaggcag	
37501					aatctctggg	
37561	agcagtgtgt	acagggaaat	ttatageact	aaaagcccac	aagagaaagc	aggaaagatc
37621					aagcaagagc	
37681					gaactggagg	
37741					ttttttgaaa	
37801			_		gagaagaatc	
37861					gaaatacaaa	
37921	-				ctacaagaaa	
37981					cttgaatctc	
38041		-		_	ccaaccaaaa	
38101					gaggagctgt	
38161					cctaacgcat	
38221					aaaaaagaga	
38281			-		atactggcaa	
38341					ttcatctctg	
38401			_		ataaacagaa	_
38461	_				gacaaaattc	
38521					tatctcaaaa	
38581	_	_			caaaaactgg	_
38641	_				ctcctattca	-
38701					aagggtattc	
38761		_			gtatatctag	
38821					gataaaaaat	
38881					ccaaatcatg	
38941					caacttacaa	
39001					ataaaggagg	
39061			catacotaco	aagagtgaat	atcatcaaaa	tggccatact
	arggaagaac	accodigec	cataggtagg	aagacccaac	accacgaaaa	cggccacacc
39121 39181	gcccaagtta	atttatagat	tcaatgccat	ccccatcaag	ctaccaatga aaaagagccc	ctttcttcac

```
39241
       gtcaatccta agtcaaaaga acaaagctqq agqcatcacg ctacctgact tcaaactata
39301
       ctacaaggct acagtaacca aaacagcatg gtacttgtac caaaacagag atatagatca
       atggaacaga acagagccct cagaaataat gctgcatgtc tacaagtatc tgatctttga
39361
39421
       caaacctgac aaaaacaaga aatggggaaa ggattcccta tttaacaaag ggtgctggga
39481
       tatetggeta gecatatgta gaaagetgaa aetggateee tteettacae ettatacaaa
39541
       aattaattca aqatqqatta aaqactqaaa ttttaqacct aaaaccataa aaaccctaqa
39601
       aqaaaaccta qqcaatacca ttcaqqacat aqqcatqqqc aaqqacttca tqtctaaaac
39661
       accaaaagca atggcaacaa aagacaaagt tgacaaatgg gatctaatta aactaaagag
39721
       cttctgcaca gtgaaagaaa ctaccatcag agtgaacagg caacttacag aatgggagaa
39781
       aatttttgca atctacttat ctgacaaaqq ctaatatcca gaatctacaa ttaactcaaa
39841
       caaatttaca agaaaaaac aaacaacccc atcaaaaagt gggcaaagga tatgaacaga
39901
       cactteteaa aataagacat ttatgeagee aaaagacaeg tgaaaaaattg eteateatea
39961
       ctggccatca gagaaatgca aatcaaaacc acaatgagat accatctcac accagttaga
40021
       atggtgatta ttaaaaagcc aggaaacaac aggtgctgga gaggatgtgg agaaatagga
40081
       acacttttac accgttggtg ggactgtaaa ctagttcaac cattgtggaa gtcagtgtgg
40141
       cgattcctca gggatctaga actagaatta ccatttgact cagccatccc attactgggt
40201
       atatacccaa aggattataa atcatgctgc tataaagaca catgcatata tatgtttata
40261
       gcggcactat tcacaatagc aaagacttgg aaccaagcca aatgtccaac aatgatagac
40321
       tggattaaga aaatgtggta catgtacacc atgqaatact atgcaqccat aaaaaqatqa
40381
       tgagttcatg tcctttgtag ggacatggat gaaactggaa accatcattc tcagcaaact
40441
       ategeaagga caaaaaaceg aacaccacat gtteteaete atagatggga attggacaat
40501
       gagaacacat ggacacagga aggggaacat cacacaccac ggcctgttgt ggggtggggg
40561
       gaggggggag gcatagcatt aggagatata cctaatccta aatqacaaqt taatqqqtqc
40621
       aatacacaaa catggcacat gtatacatat gtaaaaaaacc tgcacgttgt gcacatgtac
       cctaaaactt aaagtataat aaaaagaaga agtagaatct cattgtgatt ttaattgttt
40681
40741
       acatttatct aataactaqt aactttqtqc actttqtcct qtqcttattq aacattqata
40801
       tattttcttt tgtgaaatat cagtacaagt cttttcctca ttttaaaaaat tgggctattt
40861
       tatettttte atatgaaett gtaagagtta tttatgaagt atagatacaa gtettteate
40921
       tgatatgcaa attatgagta ttttgtctgg actgtggctt agtctattca aattctaatg
40981
       atgtctttga agaagaaaag tttctaattt taataaagtc ccagctatga tttatttttc
41041
       tttcatgatt agtgattttt gtatccattt aaaaaatatg ttcctacttc aggtttaaga
41101
       tgttctgtta tgtaatacaa tccattttta tttcattttt ctgtataata tgaggtagag
41161
       ctttttcaca tagagagtca atttttctat cattatgtgt cgaaggcagt gtttttcaca
41221
       ataaacttat qttqqcactt taatcaaaca tcaatttact atatattttt qaqtctactt
41281
       ctggactctc tcttctttga tcaacatgtc tgtctttatg cccataccac actgttttca
41341
       ctatggtagc ttttatatta agtcttaagg tcaggttgtg taagttctct gactttgttc
41401
       tttttcttta accatgtctt gggtattccc aggactttaa tttqccatqc agactttaga
41461
       tttccataga agctttagaa gcagattctg ctgctgggag aataagttac actgacctca
41521
       ggacaaaaga tgtacatgaa ctaaataaag gacgacaaat gcagaccaca gcaataaagt
41581
       ctgtggaatt agtatgtgag aaattacaca atggtctaaa atattagtct tgaagggatc
       ttaatgatga tttagtcatt tcacatgtgc aaactgttaa gactgctggc tgtttttcac
41641
41701
       atggccacca gcccatttct ttttcctgag tgtctgcagt ccacagagct tcttctggct
41761
       ctctatgctc catctgttct cggcccagga gttcccatgg gacctcccaa gggtgactca
41821
       ggtctcatct tccgataaga taaaagaaat aagacagata ttcqqqqtaq aqatqtaaqa
41881
       gcatggtgaa agacagactg ctcctgttta gatccatata taaaaqcttt attttttqaa
41941
       teceaaatte catecettgg teatetgtae tggtteatte aataaatatt tetgaggetg
42001
       ggcatggtga ctcaagcctg taatcccagc actttgggag gtcaaggttg gtggatctct
42061
       tgagcccaga tgtttgagac cagctttggc aatatggtga aaccccatct ctacgaaaaa
42121
       tacaaaaatt agctgggcat ggtgatgtgt gcatgtagtc ccagctattc tggaggctga
42181
       ggtgggagga tcacctgagc ctgggaggtg gagactgcag tgagccatga tcacaccact
42241
       gcactccatc ctgggtgaaa gagtgagacc ttgcctcaga aataataata ataatgataa
42301
       taagtaaaca tttctgagca cctacttggt gctcttccag gcacatgagc tatatcaata
42361
       gacaaaactc acaaaaataa tMttgccttc tggaggtcat attccaatgg ataaaacaga
42421
       aaataataaa catgatgcat aagtagaatg ttagaagtga tgaatgcctt agaaaaagat
42481
       tggagcaggt taaggagatg gggattggga gaggagacat tgcaattcaa aattgattgg
42541
       tcaggaactc aaacaaatac ttggacacYg atagcagcat tattcaaaat agtcaatagc
42601
       tagaagcaac ccagatgccc attaagggat gaatggataa acaaagtgtg gtaggcacac
42661
       acagtggaat attattcagt cattagaaac aataaagtac agacccatgc tacaatatga
42721
       acaaaacttg gaaagattat gctaagtgaa ataaggcagt cacaaagggc acttattgtg
42781
       tgactccatt tataggaaat attcagaata gataaaccca tagaatcaga aagcagattg
42841
       gtggtggcta tgggtttgtg gtggaagagg aagaatgggg acttcatgtt aatgggtatg
42901
       gggttatact cggggatggt gaaaatgttt tgaaactaga tagagttagt ggttgtacca
42961
       cattgtgaat gtaccaaatg tcactaaatt gtgcacttcg aaatagttaa ttttatgtta
43021
       tgtaaatttc agcgcaattt ttaaaaatca gaagcaaaaa aagaacagaa aattgagtag
```

```
43081
       ttaggctaag tctcacagag aatgtacagt ttgagtgctc acttggagga ggtaagttag
43141
       ctcttggagg aggtgggtaa caggaagaag ctgtccaggc agaaggaaca atgtctgcct
43201
       agtgeettea aggaacagea atgaggatae ggggttggag gggaggggte agagaggtga
43261
       ttggtggtgt tccatcgtgt ggtacctccc agcttctgta aagacgttgg cttttcctct
43321
       gggtgaggtg gaagactttg gaggccccct tgatttgacc taggttctaa cagaatctcc
43381
       agagtgtctg ttttaggaat atacttaaga gggtacaagt ggaagcaggg atagcaattt
43441
       agaagcagtt attaaaatcc aagtgagaga ccatggtggc ttggacaagg atggtggcat
43501
       tgagaatgga gagatgcaat tgcaggtagg atgtcctttg aagatgaaac cataagttgt
43561
       43621
       ggtctgagcc aacattcagc atggagttgt catcattaca attagaagac aatgggagat
43681
       gctgaatttt ctttggggca agggaggcaa ggaggctgga ggagatcaga agttctgttc
43741
       tggagacatg aagcttttgt gaattccact tgttccttga atatatgatg gtttcttctg
43801
       tttttgtttg tttgttttt taaattttag attctaaggt acatgtactt gtttgttaca
43861
       aaggtatatt gcattactgg tggggactgg tcttacccaa attgtgaata ttgtacccag
43921
       taggtaattt ttcattcatc accccctacc ctccctactt ttggagtccc cagtgtctat
43981
       tatttccatc tttatgtcca tgagtaccca tggtttagct cccacatatg agtaagaaca
44041
       tgtggtattt ggttttctgt ttcttaactg gttcacttaa gataatggtc tccagctcca
44101
       tocatgtago aaaggacata atotoattot tittatggot gaatagtaat cogtggtgca
44161
       taggtacaac attttcttta tctagtcaac cgttgatgga cacttgagtt ggttccatga
44221
       ctttcctatt gtgaaccgtg ctgcagtgaa catactagta gaggtggaga ggtgtctttt
44281
       ttatacacta atttctttc cttagaaaag tagtgagatt tctgggttga atggtagttt
44341
       tacttttggt tcattgagaa atcttcatac tgttttcctt agagattgaa ctaattgaca
44401
       ttcccagcaa caatgtataa gcattccctt ttctccatat ccatgccaac atctgctgtt
44461
       ttttgagttt ttgataatag ccattcagac gggtcttctg gtttttggag gggaggattg
       ggtgaagcaa gaaggagttt ggagggaagg agtggaggct tgagtgggcc tagagtttgg
44521
44581
       agtatgggca agaaggatcc cagagacaag cactttgccc acagctacac agctaatgga
44641
       gctggggggc ccagcatatt ctccagaggc ccaggctagg ccgtctaggg gcatgcttgt
44701
       totgocccat coactgoagg tootgaaaac attttoatca ataaaaaaaa ataaaaaaac
44761
       agaataaaaa tgatacccag atgtcctcta gtgaaatgag gggaaaaaaa acatccatcc
44821
       ccRgcttatt gtgagagtca ctgaatgaga gcctgactca ctgaaattca tataagatta
44881
       atgtaaccaa gttccctgtc ttttgacact ggtttacagt aagagcaggc cacacatggc
44941
       cagctctgga gtgtgttagg acatttacat tttacatttc agtgtgatat ctgctaagtc
45001
       aaatgaagaa gtcttgaaag ataccttcta agttcggaag tatttgagtg tcacattcca
45061
       tatqccaqca qttaqttqtt qccctcaaaa cataaqqttt tqtttqttct tttqtttttt
45121
       gtttttttaa tggttatgta agtgaagtag attataaata ggcacataca ggatttcaaa
45181
       actgtaacaa aattaagaaa aagctggttc aatgagStta gattctatga gattaatctg
45241
       aaaagggaga gtagttatga gaagtettaa aaaagtgggt gtttgecaga gaaacaageg
45301
       gacactggaa gatttctgat taagttggca gaaatttatg ggtgcttaaa atgccttgct
       ttattcatgt attagaagcc cttgcctttt tgcagtttgt cttaagttgc tataaactac
45361
45421
       tctctgttat gttgaatggt gcctgaatag tgagaagcca aaataattta gttcttttca
45481
       aagaagaaat tatagactag cttattttaa taacccagtc caaattataa aaagaaaagc
45541
       tttaccaScc taatctctgg tatagagaat gttctctttt tttagttgac attgggggga
45601
       Raaaagcttg actttgaagt tcagcaagtc tggtttcaac cccagccaca actgggtgaa
45661
       ctttgtcaaa ttacttaaac agtcttatcc ttgggttttt atttgtaact ataaaatttg
45721
       aStgatgtta atgataagta ccacagagta ttgtactttg aatgaaaggt cttaattgtc
45781
       aaatgaacaa ctgaattatt ttataggaag agttcaactt taataaatac tcctgcagaa
45841
       tttcatcatt atgatgttaa aattgatgga agctttattt tttaaactga gaaactaatg
45901
       cttacgtggc ctagtttaag aatgaatgct ccaacttaat acatgtttta aaaagatgtt
45961
       ttgggaYacc aaatgtaaaa gaattccgag tttgtgttat tttattgaac aaacctattc
46021
       ageacttggg gaategeagg cageattgea gactegetgg etectggetg tgateteaca
46081
       cccaaactgc aataatatcc tgataaaacg gacttccaca agtaggtcaa gaataataat
46141
       gtgcagacct aattcctcta atttacggcg aacatctcat ggtcaatctc cccgcggtca
46201
       aaggactgtg tatctcttcc tctcaqaqcc actccacagg ctqaaaggct gactaaaccc
46261
       ctgacagata agagattcaa gagtagtccc agccccatg gggagcaaag acgctggctc
46321
       aaaaaacgta ctgataactc tcccttttgc agctactcct ttctgaacct tccacaagca
46381
       gtggcagttg gattctgtga ttagtctagt cttcagagcc agccttcttg agttcaaatt
46441
       ctgacttcac cccttaatgt gaagtgacat gggcaagttg cttaatctct ctgtgcctca
46501
       gtgtccttgt ctgtaaaatg ggcatcataa taatagcgcc tgccacattg ggtgagtgtg
46561
       agaatgaagg aattaataca tgtaaatcac ttagactgtg tctggcatag aggacattct
46621
       aaagaaaagt tagctattat cattatatta ttatatgggt ctggaattag ttcctgaatc
46681
       cttctgagat gtgatgactt ataaacgtag gttgagttta ctcatgatag ggtcatcgca
46741
       actatgcata gctaaaatca aattttgctt ttcaagtttg ttttacctgg agccctagag
46801
       ttcagggtta tggttttctt tgtcactccc cttgagggaa gcttcttagt cacactctcc
46861
       ttctctttct ctgcactcta tgcactctag aaaagctcct ttttttttt cttcatccag
```

```
qcaqaqaqqc ctactqggac ttaaatccaa qqaqctqaaa tctqttttqq qatqqggtgg
46981
       agtcacattc tggaacctag acagagaatt tctaagttcc agaaagtgct gcttacttcg
47041
       cattteetet ceceaeett tgettttgaa aeteetggea ceaatgetge caaggetgge
47101
       ggagctttcc tgagtggtgt ctgccaaatg aggagtcaag gaatatctgg aaaggcagcc
47161
       tecaggteee egatgteaag accatttaga actgaaagtg teceaatate ggggtacagg
47221
       caataagcat tagttattaa tcagcctgag aagttgattc taaaatagga ggaaatgatt
47281
       caattatttc ctctcaaggg attactcaat gttgttttta tgtttaaata tttatttgtc
47341
       aacatcaaga attettagta catgatKeac cagcattttt gaacaagtea tagatttgge
47401
       cacaaatcaa atttcaggat gggaggagtg tctccccttt aaaatagaag agagtgagta
47461
       qtctatqaqq aqqqacctac aaaqactqqa aactattctt aqctccqtca ctqactccaa
47521
       gttcatcccc tctgtctttc agtttgggta agcatcaatt acttatctaa aatttgtaat
       aagaaaagtc ttcataattc atgattgtgt ttatctttat gttagtaaat ttctatgttg
47581
47641
       gKtctattct tccccttata ttttaagaaa agaagtaaga aagtaaatga ttattttct
47701
       gaaagaaaaa attaatgtat ttattattat tcatagacct tcaactactc ttagaaagcc
47761
       tttggtgact cctgggaaca agctgagtca ggaacaaaag tttgagggtt tggggttgag
47821
       aataggacaa ttacttggct aatgctttag tctagttctg attttgctac tccttagttg
47881
       ctttgtgatg actacttgca catgtgtctg tttgtgagtt atttttgtga gcatgctgct
47941
       accageetgt gtggatgtet gtggttteae atgatacatt ttatttteta aagtetaetg
48001
       aattttatgg tttttttttc aaaaatagaa actcatattc tttccttctt aatttagttt
48061
       ctttcaggac attctcatga gactggtgaa atataactgc gaagtagcat gataactgtg
48121
       aaataRcatg atcattgtga aagctatatg acccaaaact tccataggaa taagtgcaat
48181
       gettgeatat etgaaacetg etgttacate tetttgeact taataaatat gtgttgaetg
48241
       attcattctc ctgggtggtg ctgagaaagt aaaaatcatg gctgataaaa attctgttta
48301
       tggttttgtc taacttattt ttcagttgag atataggcta ctcttcccaa ctcagtcttg
48361
       aagagtatca ccaactgcct catgtgtggt gaccttcact gtcgtatgcc agtgactcat
48421
       ctggagtaat ctcaacaacg agttaccaat acttgctctt gattgataaa cagaatgggg
48481
       ttttggatct tagcaattct cacaattctc atgtattcca cagcagcaaa gtttagtaag
48541
       tattgccttc taagtataat ttaaattttt ataaattaaa taatttcaag aacatttacc
48601
       ttgttttaaq caqtttqctt ccaqttgttc caqttgaatt gtaaactgaa aaatatattg
48661
       cttttgtaca atcatttttc attacaaatc tatcagaaga aattctttgt aatgaatata
48721
       tcatcagttt ggctttgcat attcaaatgc agcagaattt aatgtggtgc ctgcacatgg
48781
       gaagtgttgt ggatgtttat tgactgacac acagtttgtt tagatagata aggctgcttt
48841
       tccattggat aactgtgggg gatttagctc caaattaatt cattaacatt tcataagaga
48901
       tcttaacagt tacttattct ttcccaaggt atcacagtat atgaccggct tctaaattta
48961
       atcctagaag aaaatattga gagtataaga attatgatct tttcatttga tcatttcagg
49021
       attgtcttta tatcttttat ttgcaggtaa acaatcatgg ggcctggaaa atgaggcttt
49081
       aattgtaaga tgtcctagac aaggaaaacc tagttacacc gtggattggt attactcaca
49141
       aacaaacaaa agtattccca ctcaggaaag aaatcgtgtg tttgcctcag gccaacttct
49201
       gaagtttcta ccagctgMag ttgctgattc tggtatttat acctgtattg tcagaaggta
49261
       ttatgcagaa ggcNcccatc ttctttcacc ctgctcccct ttcttcagtg gttgattgcc
49321
       tgagctgccc ttgctttcat tccttcccta gtcctttctg gaacagttaa atttataaaa
49381
       tgatttgaat aaaagtgatt tggataaact tctaggaata ctatcaggtt gaKgtctagc
49441
       tcattctgag ctatttggat ttacagttgc agggattgat ttgtagctga cttagagaaa
49501
       aacctagctt tcctagtgac caagataact gagagcaatt gcttactttt ggctggaatt
49561
       aagaacaaac tagcacagaa aatagtaatc tggatgtttt ccatctcagK gggcctctag
49621
       taggtgaaaa ggggcttcta accttcaagt taagccacag aaggctgatg agattgtgtg
49681
       cctaaaaatt aattactttt gttcacaaat tgttaaatgt tttgattttg tggctgtatt
49741
       tggcacacac aaaatgtcaa atgaattaaa tcaaaaagca ggatgtatta gttaagggcc
49801
       taggtctgag actagacaag tgtcaaattc atgctctgcc actagttaac tgtgcgatgc
49861
       cagcaagtta cttctctqcc cttcaqtttc cttctctqta aaatqcatat aatqtaataa
49921
       tatatttcat ctcaaagcat tattgtgaaa ataaactaag gtaatgtgtg aaaagtgctc
49981
       atcatcatca ctggtacaga gtaaactctg aataaatagt aattatcttt attacactgg
50041
       gattacagca ttacagcaga tgtagtgtat gaataaatgg tgaagaagtc attgttaggg
50101
       ctactatggg ggctatgttt ttagctcaag tgtaacaaaa aagtatttaa actctagatt
50161
       ttaatgttta tttttaaata ataaaataac cattatgtat attgatggtt tttaagaaga
50221
       aaagcaatat taagtaaagg ctgaatttag attaagttat ttcacaatgc taagtgactc
50281
       ttttaattgt ctgacttatt ttaacagtcc cacattcaat aggactggat atgcgaatgt
50341
       caccatatat aaaaaacaat cagattgcaa tgttccagat tatttgatgt attcaacagt
50401
       atctggatca gaaaaaaatt ccaaaattta ttgtcctacc attgacctct acaactggac
50461
       agcacctctt gagtggttta aggtaagaag aaatttggaa ggaaatagat gaaaattaca
       caattaaaat agacacaagt ggccgggcac agtggctcat gcctgtaatc ctagcacttt
50521
50581
       gggaggccaa ggcaggcaga tcacttgagg ccaggagttt gagaccagcc tggccaacat
50641
       gaggaaaccc catctctact aaaaatataa aaatcagctg ggtgtggtgg cacacacctg
50701
        taatcccagc agcttgggag gctgaggtat gaaaatcact tgaacctaga aggcagaggt
```

50761	tacagcgagc	ccagattgca	ccactcactc	cagcctaggc	aacaaacatt	ctgtcaacag
50821	ataaataaat	aaaagtgaag	aattactgag	aaggaaatgg	aatttcttat	ttcagaattg
50881	tcaggctctt	caaggatcaa	gatacaggac	qcacaaqtca	tttttqqtca	ttgataatgt
50941		gacgcaggtg				
51001						
		acggcgacca				
51061		acgtgaaaga				
51121	ttgcacttct	ccctcctccc	tttacttcct	cctgctccat	cttatcttat	acattctgaa
51181	ctatgacgca	aagaggtttt	ctgaacacac	tatcaagatt	taagaaattt	cagggggaaa
51241	ttacattact	aattcaaagc	cacatctgtt	ctttattctt	tYtttgtgac	ttaattttcc
51301		caatctgaat				
51361		aaaccagctt				
51421		gcttaatctt				
51481		actatggtgt				
51541	tactttcaac	atcatcaatg	gccttgagta	agtcacttca	ttctaaaaat	gtgttttcca
51601	agttatttta	aattttataa	aagcttattt	aagggaaaga	tttcacaatc	atagcttatc
51661	aatctacaaa	ggattggggt	ctccttagca	caagtcgatc	tacagacgta	gatgtaatag
51721		tatgacagtt				
51781		tccaccgctg				
51841		atatcaaccc				
51901						
		tggaggattg				
51961		agtcagaatt				
52021		gagcaaatct				
52081	tcttctctgt	agtcttattc	ttattctgta	gaagagtagt	tttctaacaa	ctactaggtc
52141	atgtaattag	ttttatgggt	tggatctgca	tttgtttaag	tgatatcaga	gaataatgat
52201	attaaaqaqc	atcataggta	ataaaaqaaa	qttttattta	agtqcctttc	tqtttcqtqt
52261		gtgtgtgtgt				
52321		aaaaacattg				
52381		ttcaatgctc				
52441						
		tgtttccagt				
52501		aaatttatca				
52561		gaattccctt				
52621		acatccctga		_	_	
52681	acacagcaat	tgcagtgttt	acatttgttt	attgtacttt	gtaattcatg	atgctttcat
52741	gtatgcatct	aatttcatct	tcatctctat	cccagagett	gggatggaga	cctgcagggt
52801	gttcattctg	ggcaatggta	gccagatccg	gtaaaacatg	tttatcttca	aagtagctta
52861	tqqaqaqatq	aagagagttc	tqtaqaaaqa	tgtggaagag	ggcagttgga	aaqaaactct
52921		agagggcaat				
52981		tggccttgtt				
53041		tgggtgaatg				
53101						
		tgctagagat				
53161		ttttttacat				
53221		cttactcccc		_	_	
53281	ggaaaaggca	ctcagttctt	ggctgccgtc	ctgtggcagc	ttaatggaac	aaaaattaca
53341	gactttggtg	aaccaagaat	tcaacaagag	gaagggcaaa	atcaaaggta	tttttatatt
53401	gaagagaacc	atcctcttcc	ccttgcacat	ggtttgcacc	tgcaaagtag	gcattaaaag
53461	taacaggttg	ctttcttagt	ttcagcaatg	ggctggcttg	tctagacatg	gttttaagaa
53521		gaaggaagag				
53581		aaggcacacc				
53641		tcacctgaac				
53701		aagacaatgg	_			~ ~ ~
				-		
53761		tgtctgatct				
53821	_	gtcctcccc				
53881	ttctttggtc	atccttgttt	tctaacttta	tgaactccct	ctgtgtcact	gtatgtgaaa
53941	ggaaatgcac	caacaaccgW	aaactgaacg	tgttcttttg	tgctctttta	taacttgcat
54001	tacatgttgt	aagcatggtc	cgttctatac	Stttttctgg	tcataatgaa	cactcatttt
54061		gtggtaaagt	_		_	
54121		aggtgctact				
54181	_	ctgagactgg	_	_		
54241		tgggaagcat				
54301						
		gcaaagcaaa		_		
54361		gaaacagcca				
54421		tgctaaacca			_	
54481		ctcgaatact				
54541	acccaaacca	taccacacac	attatcattg	ttaRactttg	taaagtattt	aaggtacatg

```
54601.
       gaacacacgg gaagtetggt ageteageec atttetttat tgeatetgtt atteaceatg
54661
       taattcaggt accacgtatt ccagggagcc tttcttggcc ctcagtttgc agtatacaca
54721
       ctttccaagt actcttgtag catcctgttt gtatcatagc actggtcaca ttgccttacc
54781
       taaatctgtt tgacagtctg ctcaacacga ctgcaagctc catgagggca gggacatcat
       ctcttccatc tttgggtcct tagtgcaata cctggcagct agccagtgct cagctaaata
54841
54901
       tttqttqact qaataaatqa atqcacaacc aaattattqa taccaaatqt tttttttqtq
54961
       tacatttcta cttctctagc tataagtctt aattatacaa caaaatacta tttttatatt
55021
       tatgtttggt aaattcaata actttcctca tcatttggaa agtcaaattg tttattgctt
55081
       ccctacagtt ttttctgaat ctagcaggat tttaatgata tcattataat ttgacacaat
55141
       aaaaggacaa catgaaactg atgaatcttt attgggttaa tttcagacac tatataatct
       tttaaaaatg taacattctt ttttatatat aaataattgg tggcatcaca aatagccaaa
55201
55261
       gcagggtgga gagagtgatc cttcctgggt gcaggcaaga aggggatatg ttttctacag
55321
       agttttcaaa acagtgataa agctgtctac aagtcattgt gctttttatc atcactatgc
       ccagacaatg tgaaacatca gagatgaagt gctcttccca cagaggtgga ctgatccttc
55381
55441
       tececactee ettggtgtgt etetgaatge aatgttgtet tggaaaacag etttecaage
55501
       atttcactcc tgagcacttg ccagtttcct cacttgttct tcacatatcc aggcaaagac
55561
       atcctgtttg ctatatgaag cattgtatcc cgtataaaag gaaggaaaga gagaaatata
55621
       ttttttacact catcactcct caggggctgt acaatcatgt agaaattgtt taatgtgcYt
55681
       gtcaaatagc caaagagtgt taaaccctga gttcccaccc atgtgtgtgg tatggttagg
55741
       attcatccag atacacagag agaggcacaa caggaggaga aaggataggg gtgtggggac
55801
       agegggeece caatatggtg taategtgge aggtetetge etgaagtget atgtggggtt
       tttcttgttt taattttgac tttaacccct gatttgtaag tttttcataa aataaacaga
55861
55921
       atcataactc atgtagatgg ctataagtgc cgtagtgttc tgtgggtctc tggtgtctgc
55981
       cagtgataag tgtggcaccc caggaaggct gtggacccca tcaaggtgct atgtgagggc
56041
       catgcttggg gtggtggtgg gcccagtaga ccctgcagcc atccatccag cctgcccacY
56101
       cacactgccc ttgtgtactc ctgctttgct acgttatcat tgatcaatgt ccctggttac
56161
       ctatgtgttt gaattatctt cRtgttacag gtgtttaatg attttgctcc ttctagctta
56221
       tttqtatttc acctgttttt ctttaaatca acatqgttac actctgtttc aqcaactqta
56281
       taaattaaac acaaattatt actactgcta ttgagttgtc atgatgaatt cttttttatt
56341
       tctgaaatta tagcatttct tgaatttaag agaacaaaaa cttgaaaggc ctatgtctcc
56401
       attittattaa ctattactaa atacatatti gatgotoata ataataatac acattitatto
56461
       attatttcct atgtacaagg actcatctgc ttattttaca tttaactttc tcatttattt
56521
       tctcaatact tcaaagtaaa agacaaagaa agttgaataa cttgcgaaat actacagtta
56581
       tggagcaagg attcaaacac agccagcatt ttcctagcta tatgtgtata cagaaagtaa
56641
       tgttttgtca tcacatacct gaattactta tacttttata aaataattca cacttacgaa
56701
       gactteetet gatgtettgg ateaacttet tteetetaet tggaagegte cagecaatgg
56761
       catggttact tcccagcaat ccccacagga agtacacatt cctctgtgca tccagctggg
56821
       gattttagag agagagtgac ctggaaagga atcctgttga aatgatttac ttacggagtt
56881
       ttcattattt aacctgatga cagtaagctc tttgtcaatt ttcacttttt ccccccaatt
56941
       ttgtgtcaca tcaccttgat aattcttgat tccatactgc tggtcattga gagaactgat
57001
       aaactattag aggttgtgga ggaattcgtg aatatgggcc agtgattttt ctaccttaaa
57061
       actgggagcc catgcatgga gacttaagac ggaaaagaac cgtatcagca aatctcattt
57121
       gagattette teacatteat cagtgeatat geatgttgea tetacatttt gtgaageaag
57181
       gagtactagg aaaaatttct gggttggttg agtaatgttc accatgagag aagtatgggc
57241
       tatgagtagg acctagaact tagtaatttg acttttgaat tctttatcaa agacgcttta
57301
       gccacttagg tgactettgc ttcttctcct acctcccaca ctgctaagcc cttagaggca
57361
       acaagtgtct cagacccgtt gatgccttct ccttaatgcc ctgacagcac cttgtccgtt
57421
       cettteteat tattgeatga gttacaacce tetgeatett gcaacceaag taccacacte
57481
       cattettete tgagtgeeca tagtttettt ceacagaaac aaactttetg etaccateag
57541
       attaatetet tgggaaaace actcaagtet ceagtgetee tgtgacette agaataagge
57601
       aatctgcctt agttggggtt tcaagtctgg acagaacctg gccctgacca aactttctaa
57661
       teatatette caccaattee ecaaactggg ttatteeatt teeteagaaa acacaettee
        tctctgcact taggcttatg gtattctctc tgcccaggtg ggcatttcct tttttgcatc
57721
       tecetgaate etacteatet tteagaggee teeteeteet eeaggaagge teeettteee
57781
57841
       agtectgeat ggageaaget etecttetaa aetectetge eageaaaaat eatgettttg
57901
       cottoccatet tagttaagtt gacaagtgto tototgcoto tittigcotti gagaatgcoo
57961
        tataatgctt agcctgatgt gttgtagagg ctcaataaat gtgactgtcc tggtacttaa
58021
       agcagccatg aaaattccca gtgggtatat cttatgtgaa attcttgtac tcccaaggga
58081
        agccaacatc catgtatcag tttattatat cttgtttcag tagtaataat aatctttttc
58141
        tttcttttga atagttgatc atcatagcat ctactgcata attgcagtat gtagtgtatt
58201
        tttaatqcta atcaatqtcc tqqttatcat cctaaaaatq ttctqqattq aqqccactct
58261
       gctctggaga gacatagcta aaccttacaa gactaggaat ggtaagtggc aaataccaag
58321
        tttttctccc aaagaaaaag tcccataata actgttggtt acctgtctat taatctttca
58381
       gtagctaggc tgctaagccc agattccatt ttgcttgcta atctgttatc agtgagttgt
```

```
58441
       gtttttggat tttcttaaag gccattttcc atcctgctat gtaaatcctc acggtcctga
58501
       gatecatete aacageteae tittetteee egataggatt getatteeta etgagiteee
58561
       aataacagaa tetgeeccaa geecagaaag gtgtaaattt cataatgtat eggtaagaca
58621
       ttatgaagtt aaacacagta gcaaaattgt tcctgttttc cagatgtgag caggttagaa
58681
       aggtccatcg gaatgcatgt gtgttctacc tgaaaccctg atctgtaaag tctaacccag
58741
       gcacacaagc atqtqcatqa aaqqaqtcaq ttctqcctqc taqatqcaaa cttcatqctt
58801
       acatttggag gcagatacat gttgcatttt tcaaacagtt gactgaagta ggagctcatt
58861
       gtttgatctg tggaaaattc agatccctta aaaaattcta tgcccttgct actattagct
58921
       ttaaattttt tcacttcccc aggggagtat ggggaaacca aagatgaaac caaagtctat
58981
       tcagcacaga aggccccctt agtcatagat tattaatcaa aattgatgag atggacaggt
       tettgtecae ttttttgtta tttagtetgt gacagtaaaa aggagaaaca etttgggatg
59041
59101
       aagactgtta tttcctgata gtgttttgtg gcagtggttt gacgtcaaca tctcttgagt
59161
       ctagaatatt ttggaggatg acatacattc accaacagcc ataaaacttg gagaggaatg
59221
       tetteaaaga gteeagtgag aatttttaga ateaagtaag aagttgtaet tettgtttte
59281
       attttcagat ggaaagctct atgatgctta tgttgtctac ccacggaact acaaatccag
59341
       tacagatggg gccagtcgtg tagagcactt tgttcaccag attctgcctg atgttcttga
59401
       aaataaatgt ggctatacct tatgcattta tgggagagat atgctacctg gagaaggtaa
59461
       agctattgac atacattagg gacagaaatt catgcttatt aaaggctgtg aactaggtgg
59521
       ccttatccct gcattggata atgaattgca tttactacca caggccttaa gaccagaact
59581
       ttaaatattt atccagaagc agacacttat ccttcaatcq cccctctccc atcattqtcc
59641
       tggtgatgag atcttcacag taatgttggt ggtgtgaatt cagagttaga agtcctttgc
59701
       ttcaaqqact taqcaaacca tcatccctac tttattccct tqqtccccca ccaqqataac
59761
       tctgccactt cttaattctg tccataagat ttgaaagagg acttaaaaaat tgatgaRttt
59821
       tgttctggta gccataggca ctagctgaaa taccttaaaa gtactcagag agtcttcatg
59881
       actttcttta tgttggtgaa acatttgcaa atttatattc tcaccagaac aaaaagaact
59941
       atagettetg tteettaata tteetaeeca attttatata ettttgacag ttttateaea
60001
       actcctgtgt ttgcagtagt tatttttccc ttcccttcat acatttatgt ccagtacctg
60061
       tggaccetee ttgtgaacte ttetetaatt teteacagae caggaagggt ttggaatata
60121
       actggggctc agctatcaag ggcctaagca taataaagta aatgttcagt tttacatttt
60181
       aaactcattt tactaagaag aggaattcac atgcctatag atgtaaaggt atgaacaaca
60241
       ggtgactttg gtttaccctg gaatatctgg gaatgctaat agcctcaata acggctcaag
60301
       agacttgtga aagatacaat ttaggagaat ctaatgcatt tcttctctca aggctctctt
60361
       cttcccccat tttcccactg gccacccatg caggtagaag taatgagtaa tgctctcaaa
60421
       cactetttte atattacaaa tgtaeteaag aaeetgeagg gaeteetgtt gtttgtgage
60481
       teaacaatgt gccacatctg gccacaagct cttcacctct tctttttcag tcYatccacc
60541
       taagttctag ttacacttct cctcagccaa acccagatgc tagctctctc acactcaagc
60601
       ttgtgctgcc ttcaatgtgt gacaacatac taatctggaa tggttttcta cttccYtctg
60661
       tcaatgcatg tcctctgacc tttcctcaac caaaataaat gaaacccaac aacaccaaac
60721
       ttcactcttc acttaggaaa gttcatgact gactaaggta ctgtctatag tctcatgcaa
60781
       cttaccttgc atttgcccca tcaaaatcct ctctgcacca ccattccatc cagtggccct
       gggtggctga tcatcatgga atccattgca cccttgtcct ctgctccagg ctgggtttgg
60841
60901
       ccaatgggag gcactggcag gaggtcagag gqtqqtcatq tacctqcctc tccacqtqqq
60961
       gettetggag cagetgtget teteettetg ggageacete tteteagetg geetgeetea
61021
       tgcatctgca atccctcaga cacccgtctg tgagaatatc ttggattgtt ttattcacgg
61081
       ttgtgctttc tcctgttgca gtgaggactg ggtatttcat gttctttctg agtccaccca
61141
       gagctaattg gaatcaccat ggtgtctacg ctgtataaat gccaaatggg aaaggggatc
61201
       tattgtccct tgagattttt ctagtttgga tcaaatcagt ttctacaaca gagattaatg
61261
       ggggtgaagt aaagtcagga cttactttcc tagagacttt ggacaatttt gaggggcaaa
61321
       gcttcctgag tttgtgattt ttaatatgtg agttaaaatt ttaagccatt gaaaaacata
61381
       taagcaaata ttctaagaag ataatatgta ttctaagaag aatgaaccag aattctaagt
61441
       ctctaatgca acaatagact aaatatcttt acattcccat ttacttggaa ggactctgtt
61501
       ttagttaaga ctaagctacc tgggaaataa aaagaggaca ttggggctga tattctcaca
61561
       aatgaaagga acacaaagaa caaaacgggt tctctatcca cacatacttc cacttctctg
61621
       agtaagtgac ttgatgtcag agaatctcac accagattat aacaataaga cttttaaatg
61681
       ttcaggatgt ttatgtttaa agcattagac tgataagaaa tctgatctat ttcttgtatg
61741
       actagatgta gtcactgcag tggaaaccaa catacgaaag agcaggcggc acattttcat
61801
       cctgacccct cagatcactc acaataagga gtttgcctac gagcaggagg ttgccctgca
61861
       ctgtgccctc atccagaacg acgccaaggt gatacttatt gagatggagg ctctgagcga
61921
       gctggacatg ctgcaggctg aggcgcttca ggactccctc cagcatctta tgaaagtaca
61981
       ggggaccatc aagtggaggg aggaccacat tgccaataaa aggtccctga attctaaatt
62041
       ctggaagcac gtgaggtacc aaatgeetgt gecaagcaaa atteecagaa aggeetetag
62101
       tttgactccc ttggctgccc agaagcaata gtgcctgctg tgatgtgcaa aggcatctga
62161
       gtttgaagct ttcctgactt ctcctagctg gcttatgccc ctgcactgaa gtgtgaggag
62221
       caggaatatt aaagggattc aggcctcagg tttcatctgg taactttcct tccatttccc
```

```
tgcttctqqq tgqatqcaaa atgqccaqaa atttttctcc tcaatcctcc accatcctcc
62341
       agecaccatt ttetteeett tteeettttg ettgeetgte tteaetgtgg tgtgggatgt
62401
       tttcactact ctttttcct ctctctctgt cttgctttgt ttctcctcat tcttgtcaat
       aatctgaata gcaaactaat cattcgagag tagattttca cgtcacttga agaacattct
62521
       gattccctca ggcagaatgt cagttgtaaa tcttgttccc aacatgccag attttcttta
       62581
62641
       62701
       tataattata tataattata taaaatataa tatagaatat ctaataatgt ataatatata
62761
       62821
       tataatatat attttatata taattattaa ttatataatt aatatataat atatattta
62881
       tacataatta ttaattatat ataattaata tataatatat cttatacata attatcaatt
       62941
       tataatatat cttatacata atatatata atatattata tataatatat attatatata
63061
       atatattta tatacaatat gatatataat ataatttata tattatatat atttatatat
63121
63181
       aattattata taaattatat aaatataaat tatatattta tatataatta ttatataaat
63241
       63301
       qtattctata taaataatat aacatatatt ttatataqaa tattatatat aatataatat
63361
       atattttata tagaatatta tatataatat aatatatt ttatatagaa tattttatat
63421
       atataatatt atatatatgt atttgagaca gagteteeta tgaacataet tgtggaetta
63481
       tgattttatt ttgacactta agagtggtca tagggtaggt gcatatttaa ctttcagaga
63541
       aacttataca ttgtttcgca aactaattgt actactttac attcccacca gttgtgtatg
63601
       agagetetge tgetteattg ceteaceaat aattggtgtt gteaagttte tttatatttt
63661
       aacaattttc acaggtgtta gatgaagatc accatggtct taacttgaag ttccccgact
63721
       catgatgatg agcccttttc atgttttcat tggccattcg tatatatttt ttagtgaagg
63781
       caaaatattt tgccttattt aaattgtatt tctttttata ttgttgatgt gtaggaatta
63841
       tttatgtatt gtggatgcaa gttctttgcc agatagatgt aaagtgagaa tatttccagg
63901
       gttatctatt catttttaag tgatgtcttt gtttagtaga catttttaat tttggtgatg
63961
       tecaattaat caatqtqtcc caaatqqcta tttttttttq tqtcctaqct aaqaattttt
64021
       tttttttgcc tgtccaagag catgatttat ttatttattt tttgcctgtt caagggtctt
       ctctgtattt ttatgaaata tttacggttc tagcttttat gtttaggtcc atggtctatt
64081
64141
       gtgaattgat acttgtatat agaacaaggc aaggacaaag ttcttttttc tgctgctgct
64201
       tettetgtag aggtaceeag etgtteeage aatgtttgtg aaagaetgtt tttteeeeae
64261
       tqaattqaca tqqtqctttq ttaaaaaaaat caattqacta qatatqtqtq qatctattaq
64321
       totactitot attitiquatq titiquatqtt totqtqaata toacattqcc ataattacta
64381
       ttgccttatt ataaatcctc aagtcagtat aagtctctca actttgttca tctttgtcaa
64441
       gattgcccag acaggccggg caggtggctc acccctgtaa tttcaacact ttgggaggct
64501
       gaaatgggtg gatcacctga ggtcgagagt tcaaaaccag cctgaccaac acggagaaac
64561
       cctgtctcta ctaaaaatac aaaagtagcc aggcgtgata gtgcatgcct gtaatcccag
64621
       ctactcgggc ggctgaggca ggagaatcgc ttgaacccgg gaggcggagg ttgcagtgag
64681
       ccgagatcgc gtcattgcac tcactccagc ctgggcaaga agagcaaaac tccgtctctg
64741
       tccggacaat cctagattat ttgcattttc atatgaattt tagaaacagc ttcttaattt
64801
       ctttgaaaaa ttttcctggg atttggattg aattgtatta aatatataga ttctgtaata
64861
       tatagagtet aaaaatatac agtgtggaga atagatgtet taagtettee aateegtaaa
64921
       cgtggtatat ccctttatta cttagatqtt tcatctctgt tagcaatatt ttgtagtttt
64981
       cacaggagat atctcatata ttatccaatg aatttatact ttggtatttt gatttttaaa
65041
       tgctatggta aatggcattt ttaacaaact tcacttttca agtttccatc tctaatatat
65101
       agaattgcta ttgattttta tattccatga tcttgataaa cttaatattt ccagtagttt
65161
       ttttttttgt agatttcaaa agttggtcta catataatta tgctctctgt taaaaccagt
65221
       agttttaatt ctttcttctc aatctttatg ctttctaatt ttatttttat ttcttcactt
65281
       gttgcaaagc agttgtccca ggtgggagcc aaaatcctta tctcattact aatcttagga
65341
       aaaagtaaaa tgtttcacta ttatgaggtc acctgtaggt ttcttataga tgttctttat
65401
       cagattaaga aagatcattt ctactctcgg ttttctgatc gttattaatt cagaaagggt
65461
       gttgaatttt gccagatgct tttatttgca tttattgaga taattacata tttttattta
65521
       ttatgtggtg aattatctaa attgggtaaa aagaattaaa atcattgtcc aattgaacat
65581
       tttgcctgtc ggctatgggt ttcccttttc ctttggttaa ataacagttc tgccacaaaa
65641
       taaaaatcta gaaacacaca ttcccctttg gcctccttaa aaaaaattaa aacttcaaca
65701
       attgcacccc tctctattac ttccctatag tgaactccca tccttctggg aaagcaaaaa
65761
       gcctggggcg catttgtgta cttgagacgc ggagcccctc tcaccttcgg catcccccgg
65821
       ggaccectet tecacectea geetteeet cetetteete tgaagageeg ggeggeeget
65881
       geggecageg gteeteecte etggeateet etgetgtgaa eegeggeetg eagggtgett
65941
       qcqaqcqqcc cqqqcqcqcc cccttccqqc cccqcaqqtc acccqqctac aqcccgqqtt
66001
       tteceggae eegegeege eggetgggea ggaagegeea gaegeetggg geecaegeeg
66061
       tccgcgggaa aagggcaagg cgctgggttt tccagcagca acctttggac cccgcgatcc
```

```
66121
       agtageteeg gtaacteeae geggggegte tgggtggagg ageeggteet ggageaegge
       tgcgaggagc acccggaccg gagggtcccc agaccgggac ctccgagtca gggaggattc
66241
       tacgccaggg agcgcccag actgagagcg ccccagaccg ctacactccc ggacccggag
66301
       cttcgcccga ccgcgggcag tgcccacctg cagcctccac cggccggggt tagcagccag
66361
       gagetgeeag aegeetgaca ttettettte tgtteetact tttttteett ettettett
66421
       tttttttttt ttgtageect etetgggtge ettatetett taateacaec tetettteae
66481
       tttccacggt agtcaggagg cggagatcgc tgcttctcac ctactttctg aacttggcct
66541
       cegcagtege gacetggegt gaaggaggag etgeegeece egecceagee teggggaege
66601
       ctctctgaag gtaagggtgg gctccgctgc cacccgcatc ccctccccac cccccagagg
66661
       aagggaagac caccatccgg ggagggattc tcaaaacaag atcaattaag agagagaga
       agatagtgtg tgtgtgaata cgtgtgaata attgatgaga cagcatcctg tctttctctg
66721
66781
       aaagtatagt tttagaaatt aagacatccc caattatttt ctggagaggt gagaacgact
66841
       gaaataaatg ataacaatag ttgtcatttt ttaagtgctt actgtgtgcc tgacactgta
66901
       gaaaattttt agatatctta ttttcactta atcctcctaa gggccacatg aaataggaac
66961
       tattattatg ctttaataaa agggaggagg aattccgtca tttacccaag agtatccagc
67021
       tactaagtag tagatcccag atccacacct acagtttttc tgattccaga gcctttgtct
67081
       tttttctgcc cagccttggg aagttaaaat gcccagcacc ttgcatcgca gttatcagac
67141
       aaattaatga ggtgtcaatt gcagccagat acagagatcc aagattattg tattcatttc
67201
       actgacttcc cttgcttcat tctcaagacc ctttgcaaat actgattttg tgtcatacac
67261
       agtgetgeta ttgggageae aceteteaae eteagggeet tggatggget aggaggggta
67321
       aatttgggga attgggtttg tgcaggttct ggggacagtg ttttccctct ccctgcacaa
67381
       atatacatcc aaaggctgag aactaaggct aaggcctagg aatgtggatt gaaaaggcag
67441
       ggtctaaatc ccagcacttg ggaggctgag gtgggcggat catgaggtca ggagatcgag
67501
       accateetgg ctaacatggt gaaacceegt etetactaaa aatacagaaa attageeagg
67561
       catggtggca tgcatctgta atcccagcta ctcgggaggc tgaggcagga gaatcccttg
67621
       aacctgggag gcagaggttg cagtgagcag agattgtgcc actgcactcc agcctgggcg
67681
       acagagtgag actctatctc aaaaaaaaa aagaaaaaaa agaaaaggca gggtccctgc
67741
       cttcagggac ctcaccagag agtgggagct agtccaggag tgatggaaac acctqcaqaq
67801
       tgatggagac gggcttggac ataggcatgt gccatttggg gctgttgcag ttggagagct
67861
       teettgagaa ggcaaataet gagecaagtt gaaaggaete tagggaeetg geatetggga
       taagtgggga gggtggccca ggtagcacag gcataggcag atttgggggg aagcataagg
67921
67981
       ctaacaacag acacttgtgt gacaaggact atgaggtggc attgattttc ttggccttac
       agatgagaag ccagattcca cacaaattat tttgtttgtt agagtttaag cgtgtgtgt
68041
68101
       tgtgtgtatg ttatgcccca agatcatttt ctcctctttg catttggttg ttttatcatc
68161
       tcacaggctg agtaaaacga ctatgcatgg aacactgtag aagaaaaatt atattaaaaa
       aaaactggtg ttatgatttt catccaggca ttttgtctca ggagttgctc gattgtatgt
68221
68281
       gttgctaatg agtgatttat attgaggcat ctacattgat tcatgtgtgt ccttgtgaat
68341
       agctgtaatc tgaggctggc catttcactt cctcaaatct caggttcctc attagaaaga
68401
       gctttgaaat ggctctttcc ctgtacatta tgtgtttctg tgaattactt ttagagaaag
68461
       gtcagtttct cagggtggga cccataaaca agggcaactg ggcagagcca ggggttcagg
68521
       tgactgtttg gaccctctgg aagggctttt tactgtgctg ccttgggaga cagagtggtc
68581
       agettggaac tgacettaag cetaggteac ecaatttttg aaggaaactt tttttttea
68641
       tttagagaaa acttcatgcc tttgggtgat taggtaagtt tgtgactatt cgttatgtgt
68701
       gtgtacacac acgtacgtac atatgtacat gcacaaatat ttaatatgtc ttagacacac
68761
       caaatatatg catttaagat gtattaaatg tactttcctg agaatacaaa taagtgccct
68821
       acgtcggcgg ctctcttcct gactgtatgt tagagtcacc tgcggagttc tgaggactca
68881
       agatgcccag tctgtcctcc agaccactgg gtggagcccg ggcatcatta ttttttcca
68941
       aacttgggtg agagctggtg gcctagactt cacattagtt gctaaaacat actttccaag
69001
       atatcatgcg ttaaaaagca tttcaaaagt agattgtgca gatgtgccaa tggatcagat
69061
       gatcacctag attcaaaatt atgtatattt tatggatgta attcattcaa taattcagca
69121
       aataacaaag aaacaagcag acataaagtg aaaatatgac agttagtgat aagtgctatg
69181
       aatagaaaaa ctaaacatgg gagggtagag gaacattggt gtgtgtgagc atgtgtgtgt
69241
       gtgtgcgtgt gtgtgcatgt gcatgtttgc atttggaaaa gcatgctgtg ctagacttgc
69301
       agtagagtgc tcagggagga cctgtacatt ataagctagt gtcagcaaac cgtaggggaa
69361
       69421
       ttagttactc cctatcttca tgtcaatcaa gaaaattaat tgttggtgga tgaaaaaagt
69481
       taaatataaa aacaaaacaa aacaatactg agccataaaa atgaagaaac tatgtaagca
69541
       tctatttgac caccaacagg aaaggatatt ctacacttaa aagcaatgaa aacatctcaa
69601
       agcaaaagaa tgatcggttt gactacatca atataaaatt tgtcatatat tagcataaca
69661
       aaagagagaa agtaaacaac aagttagaca aacatctgag aactggacaa aagacataaa
69721
       gagatggttc ataaacatga aaataaaatg gctgataata agtgattaat atttgcctca
69781
       tttataccca aagaaatgca aattcagaaa acaatgagat accatttttc cattttggct
69841
       gtctaattac attttaaaag taaagctgct cagtgtgctt ctgggtgtgg tcgatagatg
69901
       ctgtcttgtg ctattgagtg agagaatatg aatattgcaa cacttctgga atgcagtttg
```

```
69961
       tcagcatgca tctgtcaggt ttcaatgtag tcttcaactc aatcactgta ctgccatcga
70021
       gtaacatcca ggaaataata agaaatgtag acaaataata atagctaaca tttaagcggt
70081
       attaccattc accagacact tgcaatagta actcatttaa ctttcacatc agttccagga
70141
       gatggtatta ttatcccctt ctatagataa gaaaattgag gcacagaaca ggcaagtcag
70201
       ttgcccacag tcggttagct ggttgaaggt agaatggaga ttcaaatcca cccaggctgc
70261
       cacctgagec catectgeaa aaaaactgag ceatectgge ttacctgtga agatggteat
70321
       gtgattatct cataatacat ctggtaaatg ataatatatc cacatgaagt aatattaggt
70381
       ggcaattaga atggttttaa caaagagttt tgtgacacag gaaaaatgct cattatgtaa
70441
       tattaggtgg aaatgcagga tatgcaatct tttaaacatc ttttattttt tgagaaggag
70501
       tettgetetg teacetagge tggagtgeag tgatgagate teggeteaet geaageteeg
70561
       cctcccggat tcacaccatt ctcctgcctc atcctcctta gtagctggga ctacaggtgc
70621
       ctgccaacgt gcctggctaa ttttttgtat ttttagtaga gacggggttt caccatgtta
70681
       gtcaggatgg tctcagtctc ctgacctcat gatccgccca tctcggcctc ccaaagtgct
70741
       gggattacag gcatgagcca ctgtgcccgg ccaaacatct taattatgtg acaacattta
70801
       taaaaagtac aagaaaacat gttaaaatat tatacatgtt gagattattt tcttctttca
70861
       gtcttcattt tgtaaatctc ctacattgag tgtaaattgc tttggaaatc agaaaaactt
       taaaaagaaa gaaacttaag agttacggaa aagcaaaaat attgatgtta aacataacaa
70921
       ttaaaaaccc ttaaaattct tacatgtgag acatttaata tttatattct tctatgccta
70981
71041
       ttttattaat acttgctaga gtatctgtta catttatcag aatatctctt tttctggaaa
71101
       tgaattcatt atttaatata accttaatct gttaaattca cacataaaac ttaagtacat
71161
       taaaaatttt acatacacac atccacattt gcagaatggt gtgcaaatac cagctaattt
71221
       ttgtatactg agatcttaat atgtttcata ttgaaactca gagttaactt gtcagtttgc
71281
       tgagccatct cctgataact ttgtctcttg gtagatattg aatatttttt ttaaagtggt
71341
       gctttttata aggttacatc agtcttaatc agagtgacat aaaggaaata acatttttta
71401
       ttctatccaa gataaaagaa taggcagagt taataataaa atgtgtgtgg aggtagtggg
71461
       ggaggtttca aaccatagac ccagtctaat ttgattatgc acagttggta ctttccattc
71521
       atacttgttt ctttggcctt tacattattg attattatct cttggaaatt ccaccaaata
71581
       tttggtgatt tttccataag gatgtatgtg aatggtttgg cagcttagcc attaagcttt
71641
       atcctggatt caggcattga caactctgtt tttaggatgg aagatgagta gttgttggtg
71701
       agaggggcat tattctgagc tgagaagctg ccatgcactc aggaagtctc cataagcctt
71761
       agatggttta atcactcaat aattccagaa acaggccttc ttatcatctc cattttggta
71821
       atgaccaaat aggctctaag acttgctcaa gttttgcaca gcttgtcagg actcaaactg
71881
       gagtggcagc ccaggtttgt gtgtttctga aaccagactc tgccatcagg ctgtgctccc
71941
       tgcactcagg atggtaaaca agggatatgg tgttgtttat qcagtattta qactctqaat
72001
       ggtctccact gtgggggatc cttcctccac tgggctcaaa gttgtaatca gaactccgaa
72061
       tgctaggttg cagtaggatc aaagttcaat gcagatctta gttccactgg gacacagtca
72121
       atgtaaagga ggggtctgga gtgtaggaga ctgtcacttc atctctccat ttccctaaag
72181
       taaactcata tgggcagcca cagacacaYg cgaacacatg catacacacc tatgcctcag
72241
       tgacacatgt tagtgagtgc tggccatgtg ctaggcatcg tgctggattt tagtgataca
72301
       gtgaagagca aaatagactt agtgcctaga attatgaaag aattatgaaa cctacgttct
72361
       gaaggaggaa gatggagttt taaaaattca accctaaatg aagatctcat tacaattcag
72421
       gaatgttttg atgtgaaaac acacacacac actctgttac atgggcagaa ctgccataca
72481
       tagctcagca tgtcacagat tgtttattca tagaatggta atatttgtag cagtactatt
72541
       gttctgaact ttggccttgg aagcaggatt tagtagacca ggcctgaaat tcttcagact
72601
       ctgttgtatc agagaaataa attattaggg atttcttcca gaagaaaaac ttcattgcct
72661
       tgaaatttta ttaattttat atggctgaaa gttctgagaa ttggttacta tattaaggat
72721
       tataaaaatc ttctaggttg tttttttaaa aatctgtgtg ccagaagatt tttaaaacYtt
72781
       cataagatag gcacactttt gtttgaaagc tttggctgaa tctgttttat tctgttttcc
72841
       agagaagcca tttgaagcag aatccaaacc atgaattgta gagaattacc cttgaccctt
72901
       tgggtgctta tatctgtaag cactgcaggt aagtgattat acatactctc aaacatattt
72961
       catgagtaat tgaggaagaa tgtcaaagtt ttttttttta tgtggtggct actgaagatg
73021
       ctgaatttat tcagaaaaga aaagattctg cagtctgagt tcatgtagtg agagcagagt
73081
       gaaaaacttg catcttgaaa taaattacat ttgtctgagg tgaaaattta tttcatgtta
73141
       tttgaaactg aggaggtggc agccatcttt attattgggt tttaggaagt ttgaaacaca
73201
       ctttctcctt cctgagaggg aggcaaagat tcatctcatt atttaggtaa taatgagaaa
73261
       agtaaacgac aaatatatta gacagcttaa aaaaattaga gtgagacaga atatacaaaa
73321
       aagagactgg Ctagaccaag gagctcaagt aggcttcata gtagatgtca aatgccaatt
73381
       ttaatttatt agcagtgatt gcctagagtt ttttttttt atttgagaag gattctgaga
73441
       cttcattctt tagcgttcca tgattgaata gcaaggcatt ccttaggctt tgggggagaa
73501
       agtgggagca caaatattaa tttgccttga tctattatct aggagaggga aacacatggg
73561
       ctcatttccc tactttgata attactgctq aagagataqc tqaattaaqc tttttqqtact
73621
       ttcaaataac acaggtccaa tctcagggta tatttactta cgtgaatcca tttattattg
73681
       agatccctgc aatggaaccc tacccagatg ataaaagaat gaggtagatc ttcattttt
73741
       tatatgtaat gatctgtcaa acaagtgagt ggaaaagcaa ggtacaggaa aatgtgtcat
```

```
73801
       gagttccatg tcctgtggac atacactaat agccttactc ctcattataa tgagagcatg
73861
       tttcaggcac agaaacagcc atgcagccag ttgtgcaaag ttgccaagct cccagctgcc
73921
       tgccctgaag agcacttttc atgaatgttt cactctaggg agacatgaca atgttctaca
73981
       tgtaggattc actgaaccta gttgagttta aatgacaaaa tgctatgatt gtccaaactc
74041
       aagacagttt caaacacaga ctttattcca aagggcattg gttgttggca ctgctgggtt
74101
       cttatactgt gtttatggaa caataggagt gcctattttg gcacttggaa ctccacaaac
74161
       cttgaattcc acttatttat cccatcaagg accatttcca acaagcatta tggtaatcaa
74221
       caaagcctct aatggtaaac catgattaga tatccaaagc ttctagctgc Ctgaggaggg
74281
       gggtggtgtg gaStttcaat tatcaacctc ccacctctgg agagtgtgta attcacatta
74341
       agtttcttca aatggtaata cttccagaga gcacatgtcc atccaaacac agtcatatgt
74401
       gtttatagaa gctttttaag acaagcgaaa acctgaaaac ctagaaataa gccaatatcc
74461
       aacagacatt tgtttaataa accaaatccc caacatgcat ccttttataa aatggataaa
74521
       ttgcgaaatg gtaacacgat tagatactac attacaatga gaaaagaaca catcactgct
74581
       acagcatgaa tgcatttata acaatcatgt caaaataacc cagacataaa ggagcacata
74641
       ttgtatagcc tatgcatttt attatattaa ctctaaaaca ggaaaataaa tgaactgtta
74701
       atgactcacc agggttttgc atgtcaattg aagctgacac ccacatatac acatgtatca
       caccttctat gttgaagtat tagtataaaa tagagattgt attgcaacca gaaaacatga
74761
74821
       agttggtgct ccttaactct tgtgttttgt ggtagattcg acatccatgt tttcaaagaa
74881
       tgtgtattag tccattctca tgctgctaat aaaggcatac ttgagactgg gtaatttata
74941
       aaggaaagag actttattga ttcacagttc tgtatggctg gggaggcctc aggaagctta
75001
       caatcatagt ggaaggggaa gaaaacatat ccttcttcac atggtggcat caaggagaag
75061
       gctgagcaaa aaggggaaaa gccccttata aaaacatcag atcttatgag aactcactca
75121
       ctatcaggag cacagcatga gggtaatcac ccccatgatt aaattacctt ccaccaggtc
75181
       cctcctatga cacatggggc ttatgggaac tacaattcaa gatgatttgg gttggggtat
75241
       agccaaatca tatcattcca cccctggcct ctcccaaatc tcatgctctc acatttcaaa
75301
       acacaatcat geeettetaa cagteeecca aagtetteac teatteeage attaacteta
75361
       aaatccaagt ccaaagtctc atctgagaca aggaaggtcc cttcctatga gtcagtaaat
75421
       tcaagagcaa tttagttact tcctagatac aatgggagta caggcattga gtaaatacac
75481
       cccttccaaa tgggagaaat tgactaaaac aaaggggcta ctggccccat gcaagttcta
75541
       aatttaatag ggcagtcatt aaaccttaaa gttccaaaat agtctccttc aactccatgt
75601
       ctcatatcca ggtcacgctg ctgcaagagg taggatccca tggccttggg cagctatgct
75661
       cctgtggctt tgcagggtac atctttcttc ctggtttctt tcatgggctg gtgttgagtt
75721
       tctgggcttt ttcaagtgca tggtgcaagc tgtcagtgaa tctatcatcc tggggcctgg
75781
       aggatggtgg ctttcttctc ttaactccac tagctgtact ccagtgggga ctctgtatgg
75841
       gageeecaae accaeattte cetteeacae taccetagea aagattetae atgagggtte
75901
       cacccctgca gcaaactttt gcctggacat ccagcatttc catacatcct ctgaaaccta
75961
       ggcgtaggtt cccaaacctc aattettgac ttttgtgcac ctgcagcctc aacaccatgt
76021
       ggaagetgee aaggtttggg gettgeacee tetgaageea eageetgace tggtatettg
       geetgtttta geaatggetg gaggggtagg atgeagggea ccaaateeet aggetgeaca
76081
76141
       cagaaaggga ccctgggccc atcccaggaa atcatttttt cctcttaggc ctctgggcct
76201
       gtgatgggag ggtctctcat gaagatetet gacataceet agaggcattt tteccettgt
76261
       cttqqcaatt aacatttqaq tcctcatttc ttatqcaaat ttctqcaacc aqctttaatt
76321
       ccatgcccc cattcccacg aacatggatt tttctctcct accacattct caagctgtga
76381
       attttccaaa ctcttatgct ctgtcacctc ttgaatgatt tgctctttag aaatttcttc
76441
       ttccagataa tcgtctctca agttcaaagt tccatagatc tctaggacaa gggcaaaatg
76501
       caactaatct ctttgctaaa cataacaaga ttcacttttg ctccagctcc ccaaaagttc
       ctcatctcca tctgagacca cctcagcctg gactttattg tccataacat tatcagcatt
76561
76621
        ttggttaaag ccatttaaca agtctttagg aagttccaaa cttttccata ttttcttatc
76681
       ttcttcaaac tgttccaact gttccaaact gccctccaaa ctgttccaac ctctgcctgt
76741
       atgcagttcc aaagtcactt ccacattttg tgtatcttta cagtagcacc ccacttctgg
76801
       taccaattta ctttattagt ccattctcac actgctaata aaggcatacc tgagaccagg
76861
       taatatata aggagaaaga tttaatggac tcagttctgc agggctgggg aggcctcagg
76921
       aaacttacaa aagggggaaa agccccttac aaaaccatca gatctcatga gaactcactc
76981
       actatcatga gaacagcatg agggtgactg cccccatgat taaattcaag atgaggtttg
77041
       gatggggaca cagccaaacc atatcagaat acttctatca catcacttcc tctttatgtt
77101
       gtggttaaca attttgagtt aaaaccagac tgctgctttg agaaacaaac cctgactata
77161
       gagaaggcat totgtocatg tootaaggga aatgtttogt toagtttaaa tactotoatt
77221
       ctcaaaatgt ggtaaaagtg cacccgagag tcactggagg gtagattatg agaaagaccc
77281
        attaatgctg aacttgaagt accacaagtg ccgaaagagg ggaacgagca cctaYcctgg
77341
       gtagatgatg gtcagatata tttccaaagc cagagaatct gggtcaattg agaaagagg
        aaaqtqacaq atatqcaaat ttqqattqat attatattaa tqaattcttq attatccaca
77461
        tggtgaccta cagtaataat aaatgtacac acactgtaag ttacataggc acacttttaa
77521
        ccatccccac acttaccagg gaatgttaga taggcagggt gaatgcaagg tgagtgagat
77581
        ttctgatggt gcagaggaag aacgccaagg ttgctagatt tcttgttttc ttttctactg
```

```
77641
       agactectea geagggtata ttttgaggta ggtatggeag caeegtgaet tetgtggett
77701
       tocctatgaa gaagttgtaa tgtgtaatat tttcttctaa gaggtgacac aattatcagt
77761
       cttctagcta tggttggtta agttctgata ccttcctgtt ggggagaaac tgtcagaagt
77821
       tgtgttttca tctatgttct gaaagtgcct tatcctttct agtgttggaa agattgacgc
77881
       aaaaataaag aacccacagg aaattgacac cgagatttct caggactttc ttgctaacct
77941
       tgcttcttca cctaatgcat taggagacca aaaaaagtta ccttgtcatt ttggtttttg
78001
       tttttattta ttttacttta ctaatcttYt gaagaatctt gtacttcacg tccccacatt
78061
       actgtggttg aaggggaacc tttctatctg aaacattgct cgtgttcact tgcacatgag
78121
       attgaaacaa ccaccaaaag ctggtacaaa agcagtggat cacaggaaca tgtggagctg
78181
       aacccaagga gttcctcgag aattgctttg catgattgtg ttttggagtt ttggccagtt
78241
       gagttgaatg acacaggatc ttactttttc caaatgaagt gagtaaccct ttctttcaa
78301
       aatgtattte acageeetet tgteetttgt teageaacte aaatatgeag taatetgaaa
78361
       ggtcagataa tcaScacatc cttttctttc ctactcttcc tatgacatga aatacattct
78421
       Wtgttatgga acaRaataag tttatctctc tctgcttatt ttcttagtga ccttaatgaa
78481
       aggagattac titgatagga titcettita ggggateett accaatgaca agtgacgatt
78541
       gcatgaaagt cagttttaaa attcagctgc tcctgttgtc tacatgcccc cagtcaaatg
78601
       aaatcagcag ctcggacttc acgataatat taatgtgttg cgaatggggc agtgtcagct
78661
       tcctqqqtqq catccctqqt qqqqaqcaqq qatqqqqttq atqcaqaaac accctaqtcc
78721
       agtotgcagg ggotcactgg gagocaggtg gcagtgctag tggaggaagt gtgtgtctgg
78781
       cagccaatcg caacccacag ctggcggcag tgagagaggg caacacggaa gggctttgca
78841
       ttttactcat gaggtggccc tagattgtgc cagaatactc gaaattgggt tgcataactt
       ccatgaaatg agcacagcca actgaaagca tcacccacac ttaccgaaca tgtttctttt
78901
78961
       tagatgtacc tacctacata tgtaggtagt tttatagaga cttgagactg acctgtaaga
79021
       aatagctcag ggccagctga aggtgatggt gtgtagattt catatgtagc agatttcttt
79081
       tagaagcact tctagagtga gtcctgatca ttatcaacca tgctgttgag agcagagtga
79141
       gctaccactg tgaaaccttg gtagcacttc tgtRgttttg caccaaatca ggtcattttt
79201
       gtttgttttt atacatagga actcccaggc atcagaaaaa aattaaataa ctcttttgat
79261
       aaaaatcagt acatctttag atgacctttt aagagtgata attacagaaa tgggtcacaa
79321
       ctgggatttt tcataacaag ctaaggaaga atgaatactc acaaatgggg taaaggaaaa
79381
       agaacaacaa acaccgNccc ccccaccaca aacttccaac tcctgattga atgaaggcac
79441
       aatttaaagg aataaagtat totggacagg cotgagagag cacgttgaga ggaacagcca
79501
       ggcatctatg atcatcactt ggagataaca gaagcaaatg gcattggcca tctttctgat
79561
       aYgggtagta gagaatacag cctggcttag gaagcaacaa tggaagcaag agattgaggc
79621
       aggatgtcag agagaggtgt ggagagacag ggtagcagat aggagatgct, aagaaaaatg
79681
       aggtgtccca gagtggataY tggagactga tcactatgag agtaaagtgW tgaggagaga
79741
       agcagetttg acaatggeet tgaaaataaa tgggatteta catggaagea ggttgtttta
79801
       gacacccagg agcatagaag atacagacat attcaaatat agcataatta gtaagcagta
79861
       tttactgcag atgtgtgtt atacacatat gtgttctaac ttaRgtgtgt aaccetttgY
79921
       agtttgatgt gggatctRtc agtgaacaga ctttacacct gaaaatttcc cRttagaaca
79981
       gtaattettt aatttttagg agtacattge taataateaa atgeeactga ggaetgataa
80041
       tttaatcctg cagtgaaaat cactttatcc caccaaattt taagaacgaa atcaactaag
80101
       aaggaaggg atacaaaagg gataRttttc aatgatagag atgaagaaat gaaggtagag
80161
       ggaaatgtac acacagaaat ctaataaacc aatcctaggc tgacaggcag caagttcagc
80221
       ttggtgagca gcctcagaag tgggggctgg tgtttatccc ctggtgtctg tggacaaaaa
80281
       tgagtggttt tcataaccag gatgggctgg gcctgcagca ggaaagtgtg gtcacagctt
80341
       tgggtcagtt ggcctcagtg ttaacccYag ccttggagct cctggatggc aggtgcagtg
80401
       tgttgttgat tgtttcagtg ttcccagagc agtgtcctgc acatggagtg tcatcagtcc
80461
       tctagcattt gttgagtgtt aggaattgag gattacagag gtagagaaca atgtagtctg
80521
       aaaaaatggg gatattcatg taccctattc cttttaagca acaagtcata gagatccaga
80581
       aatctctatc tctccaattc attgaggtca aagttgggga gacattacga aaacacactt
80641
       gagaagaagg tgctgccaat gggatcctga atccatttca agagccaggt agctctatca
80701
       tgtcccatgg ccatggaccc actgcctata actgtgacaa cagcctgcca cattctccag
80761
        tcacggccaa tcttttataa attcatgccc aaggtgaagg ggccagattc attggattgt
80821
       gaaagtgtag aattgtgcag agcacttctg gcctctgggg ccaaatagat gtgcatttgt
80881
       atcctggccc catcttgtag caagtgtgtg gcctgagtga gttgtttaac ctgtctgaga
80941
        tttagcattc tcatctgagc ctaaggaaaa tagtcatctc ccttgcatgt acatagtcta
81001
       atacctggaa ctcagtaagt gttagtcttt tcttccttct atcctattgt agacctaaag
81061
       gggaactgga agagcctaag gtgggcacaa atgtgtcagg catagaggat aacttaaaaa
81121
        cacgcatctg ctgtatttag caaataaatc tatacatcac aggtgcagtg gaagcaaaca
81181
       tgaaatttga aacattttct ctttaaaatc ttcatttaaa aaatactgta ttgttttcca
81241
       attocagate acctititt gttgttgttg ttgttgtttt tttgagatgg aatttegete
81301
       ttgttaccca ggctggagtg caatggcgcg atctccactc actgcaacct ccgcctcctg
81361
       cctcagcttc ccgagtagct gggattaccg tacccagcta atttttgtag ttttagtaga
81421
       gacggagttt caccacattg gccaggctgg tctcaaactc ctgacctcag gtgatccacc
```

```
tgccttggcc tcccaaagtg ctgggattac aggtgtgagc cactgcacct ggcctaqtqa
81541
       tagtectgat cactaattte tgateaggag taacaacaac tataaaatet atteacaete
81601
       acttttggtc agtaaggcat ttaaagagct aattcaatta cactgagggc agcaactcca
81661
       gactttggtt tattcaacag taaaagaagg ggattggaag aagttgcctt gagaaacaaa
81721
       acctaatata tatatacaca catttagaaa acaacatcag gagtttatag accttttgaa
81781
       accaaagcca ccctcccatc ctgggccaag actctctgaa cttatccatc agtgaagatc
       actqacattg taaagatttg cttttctgac atccataata tattgtctgc agtaggatat
81841
81901
       gtgcgtgtgt gcgtgtgcat gtgtgtgtct ctgtgtgtag actgcttagg ttttcattta
81961
       ttaccccctg ctcccatgac agtttttaaa gttcttgctc ttctgaatca agcggaggca
82021
       gagcagggtt tgcacatcaa tgaatcaatc tctttaataa aaataaagga aaaatattgt
82081
       aactggttac taaagggaag atgggtgata tttgcaaagt ttctgtagca ttataaaata
82141
       cctttacact tttattccat agaaattata ctcagaaatg gaaattaaat gtcatcagaa
82201
       gaaataaaca cagctqtttc actgaaagac aagtaactag taaaattgtg gaagttaaaa
82261
       aattttttca gataacctgt gaaaacagtt actatcaaac actggtcaac agcacatcat
       tgtataaggt aatgctttta taatatttta actttaagac ttgatggaaa agataaaatt
82321
82381
       cccaaatgag gggctgagag ctaccattct ggtctcataa caaataagct gcctgatccc
82441
       acatggctca cttgccctgt ttgagccaag tatcctcact ggcttgttgt caagagctga
       tacatagtat taaaatatgt tccacagtag tttccctgga tttggtgttc aataaatgat
82501
82561
       cacagittet teteccacae tetteettea titteettet etitegataa gitagittit
82621
       aggaatgcta gtttaaggat ttccattttc tcctgtcaca tgtacttaac atgtaaaact
82681
       caattgtata tgtttttctg aatatattag ttcattttat aaaattggaa atggtaatgc
82741
       ttttataata ttttaacttt aaaactgaaa aatgaagaaa gcaaaaatca ctcataaatc
       82801
82861
       tttatgtttt ttcctctaag aattacgttg tgactatccc ttcacatctt taaatgatct
82921
       tctaaacatg ggctttcagt ttttatgtgc ctggtttcat ctaactaacc cccttgtttc
82981
       gtctaactaa tcccattgtt attggacatt tgtgaattgc ctatttttca ctattattta
83041
       aaggctaaaa tgaataccat ttccacaaat ccttgggtgt ttatatgatt tgtccttcaa
83101
       ccaaatccct aaatgtagaa ttactggatc agaggctgtg agcatgtttt tatgtctttt
83161
       tecteacact ateaaacttg tetecaggga getetgecaa ttttacecca caettgaaaa
83221
       aagaatccca attttttgtc acctccacaa cacagggaaa taatggtccc agaatgttta
83281
       aaatttgata tgaaaaattt ctaaaactcc tagtggtttt ttcacttttc ttagtaataa
83341
       attaaacttt tgtatatatt tattgcatct tactgcaatc aggcaatttt gtgggttgcc
83401
       tgtttatttt tttctgtttt cttcagttgg tatacctttt acctgttgaa ttttaaaagc
83461
       tcttatacac catggaatac tatRcagcca taaaaaaagca tgagttcatg tcttttgtag
83521
       ggacatggat gaagctggaa accatcattc tcaqaaaact aacacaggaa taqaaaacca
83581
       aacactgcat gttctcactc ataagtggga ggtgaacaat gagaacacat ggacacaggg
83641
       83701
       ttaggagaaa tacctaatgt agatgacaaa ttgatgggtt cagcaaacca ccatggcacg
83761
       tgtataccta tgtaacaaac ctgcacattc tccacatata ccccagaact taaagtataa
83821
       taaaaaaaaa agetetttta gaccaegeta taggaattat atatacacat acatagegtg
83881
       tgtgagtgta gtgtgtgtat gcataaaaaa catacgcact cccatccaca cacatacgca
83941
       acacaaattg tgtagttcag gggtccccaa tccccatgcc actccccctc actaacatta
84001
       ccacctcctg tgagatccgc agctgcatta gactctcaca ggagtaagaa ccctattgtg
       aactgtgcat gcgaggaatc tatgttgcat gctccttatg agaatctaat gcctgatgat
84061
84121
       ctgtcactgt ctcccatcac ccccaqataq qaccacctat ttgcaqqaaa acaaqctcaq
84181
       ggctccactg attctacatt atggtgagtt gtataattat ctcattatat attacagtgt
84241
       aataacaata gaaataaagt gcacaataaa tgtaatgcgc ctgaatcatc cccaaaccaa
84301
       ccattccccg ctcaacccct gcccaccctg gtccatggaa aagttgtcat tcatgaaatc
84361
       agtetetgtg teaaaaeggt tggggaeage tgatgtagtt tatatggaat aacaaatetg
       aataatatga ctctgtcgtg gaaactttaa agacctataa gagagtggtc tgaacagaac
84421
84481
       ttgggaaatc tgaagcgagt gatgataata tccttgaaat ttttgtttgt gccttatgct
84541
       tgaacactgt ttacttgtga tttccaaaga cagctcttat ctgaaatgtt ttctaaacac
       togaatcagt cotgagttga ctttattota tttcccatca atgtacaaat gtgttacagt
84601
84661
       atatgtaaat tagtgagget tggtgtgetg agttttaaga aatteegtte gggtgeagtg
84721
       gctcacacct gtaatcccag cactttgtga ggctgaggtg ggcggattat taggtcaggt
84781
       gatcgagacc atgctggcta atatgatgaa accccgtctc tactaaaaat acaaaaaatt
84841
       agctgggtgt ggtggcacat gcctgtagtc ccaactgctc gggaggctga ggaaggagaa
84901
       tcacttgaac ccgggaggca gaggttgcag cgagccgaga ctgggccact gcactccagc
84961
       85021
       atttaaatta tgagcattgg tgcaccaagg atatttagct aaacaattga ccacagggct
85081
       aagtatteta ageetgttga etgateatga etteaeeega gageaaggag getaatteag
85141
       tagataattg attggaaagg aaaatgaaga gtctctgatc ttttctatag cttgaaaaaa
85201
       ttagtttggc tttataaaat tttgacttat acataaataa tttactgacg tgaggtacac
85261
       atgtgggtgc ttcacgggca cattctatgc acaatgggtc tagccccagt ggctqccagc
```

```
85321
       tetgagggag aagtetgtgg geagtaggtt eteettgett ggtgtttgea gaaacagtgt
85381
       ccaatggtca aggatggcca ttcagccccc agacatccca tgtggtccct gcatggctct
85441
       gggacctacc tctgggtgtc agatgtgctg gattgaagaa agtggcttta ttacaattag
85501
       tattaattcc tgtgacatgt atggaaatgt gtgtgWgaga gagagagatg tttgagtgta
85561
       cctatggatt tgaatattgc tgatgcattt ctaatcagct cgctggcttc tgatggaggg
85621
       tgtcaaaggt gtgggtggtc tgctagctac tcattctctg agtaattctc cacccatccc
85681
       actatccacc ccttggatgc atagtccaga atacacaaaa ttggactgaa catgtatgaa
85741
       atagagttgt gactcaagca agggtttttg aagtagtgtt ctctctctct ctctgtctct
85801
       cgctgtttct ctctctgc atgtgtgtRa aagagagaga tgaaagaata ttgatgaaag
85861
       cagctacagt geteatattg ceceatgete tactteecag catatattaa taatgataat
85921
       catgccaaca gcaatgggat cgcatttaat tgtttcagta aaagtagtac caagtaactg
85981
       ggacatatgt ccaggaattg ctactttcca ttctttttc ctttttctct aaagatcttt
86041
       aactcaaaaa tttggatcac tgtaatatca atttggcttt tacttaaagt attttaactt
86101
       gttttattaa aaacagaact gtaaaaagct actactggag aacaataaaa acccaacgat
86161
       aaagaagaac gccgagtttg aagatcaggg gtattactcc tgcgtgcatt tccttcatca
86221
       taatggaaaa ctatttaata tcaccaaaac cttcaatata acaatagtgg aaggtaaggg
86281
       aaatcttaga attgggaaga aacagacgta ttttagtaaa atggaatttt ttcattcttc
86341
       aaaactgtgt aggaattgag agtccactta gttggggttg tgaccacacc actagttcaa
86401
       atatgatggt caaaatcttc attcYggtgc tcctctactt ggctaaataa acctttcttt
       ttggcaggaa atgaaaatct gtatgaataa acctcacatg agtcagctct ctcttatata
86461
86521
       tgaaatttga gtaaattgtg gaatcttttg ttgttgaatt tacattttca ctttacaagt
86581
       atttagtgaa gatatctata ctcaataaac tgtacttagc acaaggaggc agggaagcaa
86641
       ggaagtatga gatgaagtga gcactgtact attctcaagg gatttgattc taatcaggag
86701
       aggcaataag ctacataaag gtgtatgaaa cagcatcatc tgtgaggctg caagagtgca
86761
       gggggaggtt ggaggaagga cagatgcagg aggctccatt tgacctgggt cttggagggg
86821
       gagceteatg aaacagtgae etgetggggg aagggatete caetgggaga gtgtaaggaa
86881
       cactgatgtg ctggtgggag gggagggatg atggacccag gctgtgtgtg atggaggaac
86941
       agaatcctgg gacattaaag gtatttaatt aaacctcacc cagtgaagat attcatcata
87001
       aggcacagec aatgattgat catecagtec ettettggga actateaagg aaatgaceaa
87061
       atcacacage agtgtagtee agattttaat atctgattgt tgaagtettt tttttccatt
87121
       gagtccaaat atcccttctc gtagcttcca accactgatt ttgatttgcc ttcaagaatt
87181
       tcacagatta aatctggctc cttcttaaaa tgacagctcc tttcgtgatt tgaagacagg
87241
       ttgcttatcc agtctggatt tttttcccca tcatcctgag aaactggaaa ctgtaacact
87301
       tgtgtgggtg gcttgttcaa caaaatctgt gggctcacat ttgctacagc ctttgcttta
87361
       gcaacatctt tettetteac tacaaccata tagetatata cacccacatg ttgccacagt
87421
       aacaaattat cccaagttta gtgtcttcgg taaaatcaag gttttgccag aattgcttta
87481
       cttctggagg ctgtagggga aaatgcgttt cttttatttt tttatttatt ttatatttt
87541
       87601
       tgcaagetee geeteetggg tteacaceat teteetgeet eageeteeeg agtagetggg
87661
       actacaggca cccgccacca ctcccggcta atttttatat ttttttagt agagacgggg
87721
       tttcaccgtg ttagccagga tggtctcgat ctcctgacct cgtgatcccc ctgccttggt
87781
       ctcccaaagt actgggatta caggcataag ccactgcacc aggctggaaa atgtgtttct
87841
       ttgcctttgc tttctccagc ttctagagac cacttgcatt tcttggcttg tgccacttca
87901
       tttcagagcc aaacacattt gcttccctca cctcatctcc tttttgactc tgacctttct
87961
       gtotocotot gatgaggaco ttgagacago attgtgocca cotagataat coaqqqtqat
88021
       ccccctatct caagatcctt aatttaatca cacctagagg ttccttttgc catgtaacat
88081
       atYgtattca caggaactca ggagtaggat gtgggaaact tcagggtcta ttattctgcc
88141
       tgccacagcc cagaacctgg gaaacccata actgaatttt aaactcaaaa acttaaattc
88201
       88261
       catataaatc tctgttactt tgatactttc agaacctaca ccctctttcc caccatcttt
88321
       gctttctgcc catcagcagc tctttccttc cttctttgat gaagctgtag gaatatccat
88381
       tattaaaagg gtgcttgtta tgggccaggg atagtgccaa gtactggagt tgaatcatgt
88441
       catataatcc tcacaacagc tctgtgagac tgtggccttg ggtgagttaa tttccttagt
88501
       tcctctctgc ctcagattcc tcatcatgaa atgggacagt aatggttcat ccctcgtagg
88561
       gttggctgac aatgactgag tgagtgtgtg tacagctcta ggatggctct agggacagta
88621
       agagttctgc tgtaatgacc atctcccact gctaggttcc caaacctcac ctgcagcaca
88681
       agctattctt ctcccataat tgtccatctt atttttcttg tcttcttgga gaatgtcaac
88741
       tgagtetete aggaetecae teaaatgteR etecteette atgeececat ggeetettgt
88801
       ttgtatttct gattgctgtc agaattgcaa tacatgtgtg tctctgtcaa taacttgaaa
88861
       ggtttctgga aaggtttgca ctgacttctg ttcctcatct ttgcatcttg gatacctaca
88921
       attggcactt ggcacataaa ggtgctcatt acatgcctgt tgtgtgcatc actgagacag
88981
       aaacaactcc tttgatgcca catacaacaa gaagtctaga cacttcacca tcctggctat
89041
       ttttctttgg atttggcaaa gtttgccacg agtctacaga aattagagct tggagttggg
89101
       gaagacagcc aggaggtgtt ctgacctgtg ttgaggacag agaggttctg agctcgtgcc
```

```
89161
       ttctccatgt tgtgtcttta ttacagtccc cttgccttca cagatggaga tctctcagct
89221
       gctacatagt gactgctgta gtggtggaaa ggtgtgatac ctctcctccc catcctaagg
       gtcatggcaa tgttcccata acaaaagaca gattaacaag gggcaagcaa gttttatgtg
89281
89341
       acatgggagc cttcagaaat gaagaccctg gggccgggtg tggtggctca tgctggtaat
89401
       cctagcactt tgggaggccg aggcggatgg atcacctgag gccaggagtt cgggacaggc
89461
       ctqqccaaca tqqcaaaacc ccatctctac taaaaacaga aaaattagct gggcttggtg
89521
       gtgggcacct gtaatcccag ctacttggga gactgaggca gggagaattg cttgaaccca
89581
       ggagatggag gttgcagtga gccgaggttg tgccattgca ctccagccca ggtgacagag
89641
       caagactcgg tctcaaaaaa aaaaaaaaaa aaaaaaggaa agaaatgaat accctaagac
89701
       tcagggaaac ccgttttttt ctttttaaat gcttaggttc tagaagcatg gacagccatg
89761
       tagaaatgtg actggacaaa agggtatgac ctaatgctaa tggactgagt ggggaaaccc
89821
       agcaagacct gtccagattc ttcttggcct ctctgttata tttccttttc ctgggtagag
89881
       ggcagagggt cttaagactt actgtcagac caaagaaggc caaagaattt atagccaact
89941
       cctagacaga aaggcaggag aaggttagag tcatgcttct aggttctatg gcttactttg
90001
       gggaaaagga ttctagtttt tatgatctgc ctttgggaag agggagtctg gtttctgtga
90061
       cttgctttgg ggagaatgga ggcatgaagc aggagggtca gagagatctt gctttttgag
90121
       gctgctttga ggccttccaa tgtcgttgag tttaaagtac tcagcctgcc aaaacaccat
90181
       actitigggac tgtttttgga gcccaggagt tccattgcct tcagcttgtg ctcgtgaatc
90241
       acagtcacct gaattcctac tgggcatctg ccactttcct ggtcttatgc tctggctgat
       cgggtggcgt ctcctcctac tgctgacaac ctctatgcca ttttgactct gatagcctcc
90301
90361
       taggtaggct ggccatgtca agggcatgca ctaagatgtt gtcttccttc catgacctac
90421
       atgggactgc accettcact ttacaccagt tgtgcaaact cagggctgat ctggatttcc
90481
       aatgatataa tttgagtgtt agtagacagc atcctttgct gaagatgtac ttcacactgc
90541
       cccttctcaa gtctcccatg taagtttctg ggggaaaaaa tggatccaga gaaaggtagt
       cagtagcagt tttcccctc aYaccgaaat tgcctcctgc cctttgtctt tgaattcaag
90601
90661
       coggtgttgc ttcttactgt cagacctgag agttagttct ttccctggat acatctcagc
90721
       tectectgee tgtttgecaa catteeagge ttgeecagag acteeaaate ecaggataae
90781
       ccagacetga etetgaacte eggagteeac tgtgetggtt teatteteet gttettgete
90841
       acatatetge ceaetgeatt eteaactget caetetggea etgeaettte tgagaggget
90901
       cagggaggtg tgggagctgc aggWgcactc ctgagtctca ctgtactcca ctcaaccagg
90961
       cattctgctt gctgagcacc tgcaagtccc agaagggatg ttctggacat tgcttgggaa
91021
       tgcctccaaa gcagtggggc tactagcagg gaacagtcct ggtcatgagg aaagttacca
91081
       atgaggatat cagggctaac aataacttga ctatgcacca gttcaaatct atgaccgcag
91141
       ctgctcttag ggtttgcagt tgccagctgc cttaagcaag agcctgttgg gtgatgactt
91201
       actteceaag cateaceaga tggggaetga gggggeagaa tttatgtgea gtaetteeag
91261
       tcggggtcag aagccactca ggagcagaag tagaacttgg cctacaggca tgaacttaac
91321
       agcagggcca gggccaaggt gaggttcctg gggcacaata ttgaagggac actgctctca
91381
       gcatcctgca agggaacggt gggcacatag agtgagggat ttgcttttct caacctgtct
91441
       tggccctgca acctgtaaat ccatcactgg ccagctgagg aggacagatg gagagaatcg
91501
       agacagggca gaaaggctct tcccagtgat tttgcatggt ttgactcttg cacgtttgat
91561
       catctgagag tcaggctgca gtggatgggc atcttcctct gccttttagt tgaattgagc
91621
       agaagctgca gatMtgagtg agagaaaacc agatacaagg tgctttcaac atctgtgacc
91681
       tttagttaat gcaaatagga ctcaagtgaa cgtaaaccag taactattgg aatctttgtt
91741
       acatgaaatg agcatattat catagatgct tgtcataaag aagctcaagg caacactaat
91801
       acatttgtet tttettttgt caettetaga tegeagtaat ataqtteegg ttettettgg
91861
       accaaagctt aaccatgttg cagtggaatt aggtatattt caatatacat atattctgca
       tttataagta ggggacataa tagagccttc caagcgtaaa tatgattcat tattgaatga
91921
91981
       cactggatac acatttcata tttactgaat aacattacaa aactaagaRa agagtgccat
92041
       tttttgaatg ttgccacttt taacctgact taaaagtggc aacattttaa agttcaagag
92101
       acagtggaac tgcctgtggt ctgtagacta aggtttttgt tgtgctgtgt gtggcttgtc
92161
       tttaactctc tctatggtca tttctaaata tgaaaattta ggccgggcat ggtggctcac
92221
       gcctgtaatc ccagcacttt gggaggccga ggcaggtgaa tcacttgagg tcaggagttt
92281
       gagaccaacc tgaccagcac ggtgaaaccc catctctact aaaaatacaa aaattagcca
       ggcatggtgg catgcacctg tagtcccagc tactcaggag actgaggcag gagaatctct
92341
92401
       tgaacccagg aggcaaggtt gtaatgagct gaaatcacac cactgcactc cagcctgggc
92461
       aacggagcaa gactctatct caaaaataat aataataata acaataataa atttaaaaat
92521
       tccgtgagag cactctggag agatgagtca aactaatttt gtggtgtgaa tgtttgatta
92581
       attgattcac tttgaatcaa ttgactgaat tgattgtata tttggagttt gtctagataa
92641
       gtcttatgaa tcaacatgtc caattgagta atttctttca aaagagaaag gaaagaaaga
92701
       aaggaagaaa ttcacatatg ctaaatatgc taagccaggt atcatgctag gtcattgaaa
       tatattgttg aattaaatcc tccaagcaac actgaggcat gtattctcat tttatgaatt
92761
92821
       aaataattga agtgcaaaaa ggctaagtaa cttgtctaaa gttaaatcac taaaaagtgg
92881
       caagtetetg tttggggaet ttgetteeaa agteeceace ttataaceae agttggggte
92941
       acagtggttt atatgaccac atgggcccag tgcagctggg tcctatgcca tttgtacatg
```

```
93001
       aggittacct getttatatt titattiatt tetgietitg tettiettat tittitette
93061
       tactcagtgg ggcttaactt gtatcttttg ttctagtttg ttgacctgaa gagttagata
93121
       attaatttta attatcctct ggtagtatat accttcaaac tccaagtttc cctgtaaaca
93181
       ctactttagc tgtttctcat attttcattc tattcagttt aaaccattac ttaaaatttc
93241
       ccttatgatt tctttttcca acaaaaaaaa ttggtatttc ttaatttcca aatatttggt
93301
       gaactttcta gttcttgtct cttattcatg tccagtttag attcattata gtccaagatt
93361
       atacacagtg tcatttcaac cttttggaat ttgtagtgac ttacttaagg cccacagtat
93421
       ggtatatctt gctgaatgtt ccaagtgcat ttgaaacaaa ggtatattct ataggttttt
93481
       gttttattat tetttaaatg tetatttggt taagtteatt aatggtgttg tteaaaaaat
93541
       ttatatectg actaatettt tatetgettg tteeatgtgt tgetgaaaga gatgtgaata
93601
       totocagota ttatggtcaa tgtgtctcta totoctttag ttotgtcaat ttttgtctta
       catagtitgg agctatctta tatgcataca cggatttatg ctcgttgaat taacactttt
93661
93721
       atcattatga aatatetete titatetate ataacaette eteteteaat qtqtaqetet
93781
       gtcatatgaa ataatcacac aagttttctt ttgcttggaa ttggcatatt tgtcttttt
93841
       taaacatccc tttactacta acatttttt ttgttctata cataagtgtc tctcataaac
93901
       tgcatgtatt tgggtctttt ttgaaatatg gtcagtatgg tgtttagtcc atctgtatat
93961
       ttaaaattac tgttacagtt gctatatcat atatgcatga tatacaaata ttcaatttat
94021
       totttttaaa ataaattaca otttgttgaa aatgaccata tttocatoca ottgtttato
94081
       ttttcctcta ttttctttca aacagttaca gtttttaatc tgctaaattc agcatctgta
94141
       ggtctgcttc tattagctgt cttttctatt gattatggtg cacattttct gtgtctttgt
94201
       atgtctctta actttttatt gttgttgatc tttatgtatc aaacaattaa agtgaataca
94261
       gtttgtgtta ttttttctta caagtaggta gcctgagagt tgagtatctc agttgactca
       gaattgagtt ggggtggctg ggatgtaggt atgttcactc caatctgtct tcatctctga
94321
94381
       tagetggtga gggaggggt getgeagtea tgtgtttgat geaggeetet tetetaatga
94441
       gacttgctct cctgattacc atgagactgc aagtgatatc atacagcctt tacatccttt
94501
       tacgctgtcc gagcgcttag cagtaattct gcaagcctgt gcagctgccg agagcttcac
94561
       tggattttct tgtttctatg agactctttc tgttaattac aaaccaaagc aaattcttcc
94621
       atggaatgaa atggccacct atcttagctc accctagaaa ggatccttct taagtacgct
94681
       tgtagttcaa aaatttctct ttattttctc tqtactttqa tqtctttaat qtttqcttaa
94741
       tttaattttt ttgtgattta tccatttggt ttatctagaa gtctcagtga gagtattgtc
94801
       ctgatgctac gaaaagaaag tacactgatc actctaagtt ttaatagttg attaaaaaat
94861
       aaatgtttaa aaagaagcca aatcatgttt tgatgatgac ttgtaattgc tacctatttc
94921
       ttctgcttta tatattttgt tttatttaaa gcaaggcata tttattctct ttctgggatg
94981
       gagcaccacg ttttgcttta ggttaacata cattgattat attgttttga aacttttaag
95041
       aaacctgagt ttcaaaaatc agaactttag ttgccaactt ttaccaggaa aaaacgtaag
95101
       gctcaactgc tctgctttgc tgaatgaaga ggatgtaatt tattggatgt tYggggaaga
95161
       aaatggatcg gatcctaata tacatgaaga gaaagaaatg agaattatgt atgtatgtgt
95221
       aatatatatg tcatgatata taccatataa caatcatata tattatatat aacatcatat
95281
       taaaaacact gggtggctga tgtgtatttt gggctagttt aaattataat tataatggac
95341
       atattttRta attatagtgg ccatacccta ggtggacgca ttatgcagaa atgtattgcc
95401
       aagatctaag ctggaaaaga aggtccatga gaaggccatt gagggtattt agcatcacct
95461
       tataaagtag cacttggtca gtcaggatcc gatggggatg gggattttac ttagcttctt
95521
       agaaaagggc agttgcctag gccaagcttt tttttatttt gaaaaatttg ttaagttgtt
95581
       tatttttcta atgggccaaa aaggctccaa atggcatgca tattgatagt taaataagcc
95641
       tatattette cattaaggaa tggteaaaet tataaaaaga ateagtetaa acatatagte
95701
       acctaagett ttetgaatga etttgggtea taetgtaagt atteattatt ataetgatae
95761
       actactcagg gtctagtgag tgctgagcct aagttgtcta ccacctttgc tgcacttgtt
95821
       gagtttttte ttetgetttt tacacatttt ttggtacett etttgttace atgettgeag
95881
       ctctacattc ttgctgacac ttgtctacaa gctctttaca ccacaaagtc tggcttcttg
       cccgggagca ggcatgtggt tcatgcccac tacatgttga attcatttga aatttgtgga
95941
96001
       totootgoca toatatotoa tootoagoca otggootoca tttgagtttt gtatttotoo
96061
       ctttggataa tctctttcta atttttagag gggtagaaaa ccacagtgaa tttcttagaa
96121
       atgcagctta agctacaaaa gcagRgaggt ctttctttat agttgggtgc acacaccttt
96181
       gatateatet gttttgteet eteaatggee teetgtgttt taettetgga ttteattggt
96241
       tttagatggt tgttatttct ggatgggtga cttaggaaaa gaattacatt ccaaaatgta
96301
       cttgaggcag ttggaggcta cagtggggct ctgtgtcccg ggtcctagat atccctgttg
96361
       ctaggaatgt gggcacatgg ggcccatgtc tggctgtgtt gactggtgta gctatcagag
96421
       cctgtcattg gtgggggaag tggatgtcat aagtgtttac ctggcatcaa ccacaggaca
96481
        acttgaatcc tttaaggaag agaaaaagaa cgtgctgcca gtggctttct caagtcattt
96541
       tctgacactc atggaacaac ttaactggca cttctcatgt gcctcggggt gcttgagaac
96601
       agtgttatgt tttttaatgc tgaacacatt aaaccagttt gttggattcc atggacaaat
96661
       gtaatacgaa tctgcatcta tqctaaacaa cttacaqcct tqtcataatt agtqqqaacc
96721
       actctgtgga aagaacccca agttgttagg aagagagaag tagagaaaaa taattgagag
96781
       agaatatttt ctYatatttt caacttagca tcatgatctt ccagccattt aaacttcagg
```

```
96841
       taattgacag caccattccc aaatggcctc aggagggctg tttctgtcag ttctgatgct
96901
       ccacggagga ggggacgttt attgtagtcc atgctttaag agtactggcc agaaagggaa
96961
       gaaataatgc aggcaacatc acagctgcct tcaagcattt taaacatgtg aattMccctc
97021
       tcaagggtaa cgaacagagc ctactgctaa attattttgc cctYttacag gactccagaa
97081
       ggcaaatggc atgcttcaaa agtattgaga attgaaaata ttggtgaaag caatctaaat
97141
       gttttatata attgcactgt ggccagcacg ggaggcacag acaccaaaag cttcatcttg
97201
       gtgagaaaag gtgagaaaga tttatttttg gaagttttga aacttagctc attgctactg
97261
       agagacattt caaagatggc cactttctca ttttccacac cagaacccag ctcctgagag
97321
       gggaaatcag tcacctcatg tcacaggcat tcagtggggc ccctctgtct catgatgcca
97381
       agcagagaat gaaaacagaa ctcattcacc tgctctcaca gtcatattcc atcccatgtg
97441
       gaaatcgggg ggcctgaggc ttggagatac agatgctgag agtgaggctc atggggaggc
97501
       catcaggaca acccctgcac ccagcactgc acccagagca gccagagggg agtttgcaga
97561
       tctgcgtggg gtgaggccag tgcaaacttg gtgtgactgc tgctccagcc tgcctcagca
97621
       gggtaccagg ctcagggtcc ttgtgtgtgc tcattcattc attgcctcct gaacttggca
97681
       agcaggccat tgatagccct gatctgtggg Mtcttaaagt ttttcagtgc tcacctgatg
97741
       gagggcccca gggtcgccat ccctttcctc caccaaagta gcacagccca gggccttcag
97801
       cccagtagct gctcagaaat ggctgtgact ggagccagaa agctgtgatt tccaKttgtc
97861
       actactcatt cactgttcat tcccaggctc caccatcaac tccctggata tgtagagaaa
97921
       ggtcaagatc tgcaagagag agagaaaggg agagagacag agactaacac acccagagac
       agatggagac ggagagaca agacacagag agaggagaaa gagagataca tgagatggag
97981
98041
       ggacagagac aaacagacaa ctctagagag atgaagacag aaagagaaag agaaagagag
98101
       acagagacac agagagacag agaaatgaga gagacaggga gagacacaga gacagaggaa
98161
       caqaqaaaqa qqqaqataqa qaqqcaqaqa catqaaqqqa tqcaaataca qaqaqatqaq
98221
       agtgaaacag acagagagac agagtaaggg agagatacac acagacacac acaaacatgc
98281
       agagacagcc acacacaga ctgggcctt ggctttgtga gaaacggaag gagtgggaca
98341
       ggaagagcac aaaggctgga aaccccctcc agctgggcta tcctggtgcc caggggcagc
98401
       teccecagea cacatgitti agageegeet etetaceata igeacteata acaacagete
98461
       acatcccata catgttaatt acatgacagg atctgggtga agttcctgcc atgatctcat
98521
       ccagtcttaa cgacaagctt tatagaggtg aagaaacttg cccagggtca cacggcaggt
98581
       gaaggtattg aaactcagac ccgattaaat ctggaatctg tgttcttacc ttctgcaggg
98641
       cttgtcgcct gtggtcttcc acccctgtgt gattgcctgt gctgtggtcc actcacttgt
98701
       gagggagYgg ggctcaaatc tttgttggac catcacctgg ccaggctttc ctggggcctg
98761
       ctcaccaccg gatcacctct cctttcttag aacagggcag gactgtccta ttcctgccct
98821
       agacgettet ggecagteaa tteteetget getgetetea caaatgtagg tttagaaatt
98881
       ggggctcaca acccaccatc accccgtgcc aatgtgtgcg tgcgtgtttt tgcaaatgtg
98941
       tgacacagtt
```

[0251] Three alternatively spliced transcript variants encoding distinct isoforms are depicted hereafter. cDNA sequences 1-3 show the human cDNA structure for transcript variants 1-3 of IL1RL1, respectively. cDNA sequence 1 encodes the longest isoform (1), which is depicted hereafter as amino acid sequence 1. cDNA Sequence 2 differs in the 5' and 3' UTR, compared to variant 1. The resulting isoform (2), depicted hereafter as amino acid sequence 2, has a distinct and shorter C-terminus, as compared to isoform 1. cDNA sequence 3 differs in the 5' and 3' UTR, and contains an additional internal segment, compared to variant 1. The resulting isoform (3), depicted hereafter as amino acid sequence 3, has a distinct and shorter C-terminus, as compared to isoform 1.

#### IL1RL1 cDNA Sequence 1 (SEQ ID NO: 2)

NM\_016232 Homo sapiens interleukin 1 receptor-like 1 (IL1RL1), transcript variant 1, mRNA

```
1 aaagagagge tggetgttgt atttagtaaa getataaage tgtaagagaa attggettte 61 tgagttgtga aactgtggge agaaagttga ggaagaaaga acteaagtae aaceeaatga 121 ggttgagata taggetaete tteecaaete agtettgaag agtateaeea actgeeteat 181 gtgtggtgae etteaetgte gtatgeeagt gaeteatetg gagtaatete aacaaegagt
```

```
241 taccaatact tqctcttgat tgataaacaq aatqqqqttt tqqatcttaq caattctcac
 301 aatteteatg tatteeacag cageaaagtt tagtaaacaa teatggggee tggaaaatga
361 ggctttaatt gtaagatgtc ctagacaagg aaaacctagt tacaccgtgg attggtatta
421 ctcacaaaca aacaaaagta ttcccactca ggaaagaaat cgtgtgtttg cctcaggcca
481 acttetgaag tttetaceag etgeagttge tgattetggt atttatacet gtattgteag
541 aagtcccaca ttcaatagga ctggatatgc gaatgtcacc atatataaaa aacaatcaga
 601 ttgcaatgtt ccagattatt tgatgtattc aacagtatct ggatcagaaa aaaattccaa
 661 aatttattgt cctaccattg acctctacaa ctggacagca cctcttgagt ggtttaagaa
721 ttgtcaggct cttcaaggat caaggtacag ggcgcacaag tcatttttgg tcattgataa
781 tgtgatgact gaggacgcag gtgattacac ctgtaaattt atacacaatg aaaatggagc
841 caattatagt gtgacggcga ccaggtcctt cacggtcaag gatgagcaag gcttttctct
901 gtttccagta atcggagccc ctgcacaaaa tgaaataaag gaagtggaaa ttggaaaaaa
961 cgcaaaccta acttgctctg cttgttttgg aaaaggcact cagttcttgg ctgccgtcct
1021 gtggcagctt aatggaacaa aaattacaga ctttggtgaa ccaagaattc aacaagagga
1081 agggcaaaat caaagtttca gcaatgggct ggcttgtcta gacatggttt taagaatagc
1141 tgacgtgaag gaagaggatt tattgctgca gtacgactgt ctggccctga atttgcatgg
1201 cttgagaagg cacaccgtaa gactaagtag gaaaaatcca attgatcatc atagcatcta
1261 ctgcataatt gcagtatgta gtgtattttt aatgctaatc aatgtcctgg ttatcatcct
1321 aaaaatgttc tggattgagg ccactctgct ctggagagac atagctaaac cttacaagac
1381 taggaatgat ggaaagctct atgatgctta tgttgtctac ccacggaact acaaatccag
1441 tacagatggg gccagtcgtg tagagcactt tgttcaccag attctgcctg atgttcttga
1501 aaataaatgt ggctatacct tatgcattta tgggagagat atgctacctg gagaagatgt
1561 agtcactgca gtggaaacca acatacgaaa gagcaggcgg cacattttca tcctgacccc
1621 tcagatcact cacaataagg agtttgccta cgagcaggag gttgccctgc actgtgccct
1681 catccagaac gacgccaagg tgatacttat tgagatggag gctctgagcg agctggacat
1741 gctgcaggct gaggcgcttc aggactccct ccagcatctt atgaaagtac aggggaccat
1801 caagtggagg gaggaccaca ttgccaataa aaggtccctg aattctaaat tctggaagca
1861 cgtgaggtac caaatgcctg tgccaagcaa aattcccaga aaggcctcta gtttgactcc
1921 cttggctgcc cagaagcaat agtgcctgct gtgatgtgca aaggcatctg agtttgaagc
1981 tttcctgact tctcctagct ggcttatgcc cctgcactga agtgtgagga gcaggaatat
2041 taaagggatt caggcctc
```

#### IL1RL1 cDNA Sequence 2 (SEQ ID NO: 3)

#### NM\_003856 Homo sapiens interleukin 1 receptor-like 1 (IL1RL1), transcript variant 2, mRNA

```
1 gaggagggac ctacaaagac tggaaactat tcttagctcc gtcactgact ccaagttcat
  61 cccctctgtc tttcagtttg gttgagatat aggctactct tcccaactca qtcttqaaqa
 121 gtatcaccaa ctgcctcatg tgtggtgacc ttcactgtcg tatgccagtg actcatctgg
181 agtaatetea acaacgagtt accaatactt getettgatt gataaacaga atggggtttt
 241 ggatcttagc aattctcaca attctcatgt attccacagc agcaaagttt agtaaacaat
 301 catggggcct ggaaaatgag gctttaattg taagatgtcc tagacaagga aaacctagtt
 361 acaccgtgga ttggtattac tcacaaacaa acaaaagtat tcccactcag gaaagaaatc
 421 gtgtgtttgc ctcaggccaa cttctgaagt ttctaccagc tgcagttgct gattctggta
 481 tttatacctg tattgtcaga agtcccacat tcaataggac tggatatgcg aatgtcacca
 541 tatataaaaa acaatcaqat tqcaatqttc caqattattt qatqtattca acaqtatctq
 601 gatcagaaaa aaattccaaa atttattgtc ctaccattga cctctacaac tggacagcac
 661 ctcttgagtg gtttaagaat tgtcaggctc ttcaaggatc aaggtacagg gcgcacaagt
 721 catttttggt cattgataat gtgatgactg aggacgcagg tgattacacc tgtaaattta
 781 tacacaatga aaatggagcc aattatagtg tgacggcgac caggtccttc acggtcaagg
841 atgagcaagg cttttctctg tttccagtaa tcggagcccc tgcacaaaat gaaataaagg
901 aagtqqaaat tqqaaaaaac qcaaacctaa cttqctctqc ttqttttqqa aaaqqcactc
961 agttettgge tgeegteetg tggeagetta atggaacaaa aattacagae tttggtgaac
1021 caagaattca acaagaggaa gggcaaaatc aaagtttcag caatgggctg gcttgtctag
1081 acatggtttt aagaatagct gacgtgaagg aagaggattt attgctgcag tacgactgtc
1141 tggccctgaa tttgcatggc ttgagaaggc acaccgtaag actaagtagg aaaaatccaa
1201 gtaaggagtg tttctgagac tttgatcacc tgaactttct ctagcaagtg taagcagaat
1261 ggagtgtggt tccaagagat ccatcaagac aatgggaatg gcctgtgcca taaaatgtgc
1321 ttetettett egggatgttg tttgetgtet gatetttgta gaetgtteet gtttgetggg
1381 agettetetg etgettaaat tgttegteet eeeccaetee eteetategt tggtttgtet
1441 agaacactca gctgcttctt tggtcatcct tgttttctaa ctttatgaac tccctctgtg
```

```
1501 tcactgtatg tgaaaggaaa tgcaccaaca accgtaaact gaacgtgttc ttttgtgctc
1561 ttttataact tgcattacat gttgtaagca tggtccgttc tatacctttt tctggtcata
1621 atgaacactc attttgttag cgagggtggt aaagtgaaca aaaaggggaa gtatcaaact
1681 actgccattt cagtgagaaa atcctaggtg ctactttata ataagacatt tgttaggcca
1741 ttcttgcatt gatataaaga aatacctgag actgggtgat ttatatgaaa agaggtttaa
1801 ttggctcaca gttctgcagg ctgtatggga agcatggcgg catctgcttc tggggacacc
1861 tcaggagctt tactcatggc agaaggcaaa gcaaaggcag gcacttcaca cagtaaaagc
1921 aggagcgaga gagaggtgcc acactgaaac agccagatct catgagaagt cactcactat
1981 tgcaaggaca gcatcaaaga gatggtgcta aaccattcat gatgaactca cccccatgat
2041 ccaatcacct cccaccaggc tccacctcga atactgggga ttaccattca gcatgagatt
2101 tgggcaggaa cacagacca aaccatacca cacacattat cattgttaaa ctttgtaaag
2161 tatttaaggt acatggaaca cacgggaagt ctggtagctc agcccatttc tttattgcat
2221 ctgttattca ccatgtaatt caggtaccac gtattccagg gagcctttct tggccctcag
2281 tttgcagtat acacactttc caagtactct tgtagcatcc tgtttgtatc atagcactgg
2341 tcacattgcc ttacctaaat ctgtttgaca gtctgctcaa cacgactgca agctccatga
2401 gggcagggac atcatctett ccatctttgg gtccttagtg caatacetgg cagctageca
2461 gtgctcagct aaatatttgt tqactgaata aatgaatgca caaccaaaaa aaaaaaaaa
2521 aaaaaaaaaa aaaaaaaaaa aa
```

## IL1RL1 cDNA Sequence 3 (SEQ ID NO: 4)

## NM 173459 Homo sapiens interleukin 1 receptor-like 1 (IL1RL1), transcript variant 3, mRNA

```
1 gaggagggac ctacaaagac tggaaactat tcttagctcc gtcactgact ccaagttcat
  61 cccctctgtc tttcagtttg gttgagatat aggctactct tcccaactca gtcttgaaga
 121 gtatcaccaa ctgcctcatg tgtggtgacc ttcactgtcg tatgccagtg actcatctgg
 181 agtaatctca acaacgagtt accaatactt gctcttgatt gataaacaga atggggtttt
 241 ggatettage aatteteaca atteteatgt attecaeage ageaaagttt agtaaacaat
 301 catggggcct ggaaaatgag gctttaattg taagatgtcc tagacaagga aaacctagtt
 361 acaccgtgga ttggtattac tcacaaacaa acaaaagtat tcccactcag gaaagaaatc
 421 gtgtgtttgc ctcaggccaa cttctgaagt ttctaccagc tgcagttgct gattctggta
 481 tttatacctg tattgtcaga agtcccacat tcaataggac tggatatgcg aatgtcacca
 541 tatataaaaa acaatcagat tgcaatgttc cagattattt gatgtattca acagtatctg
 601 gatcagaaaa aaattccaaa atttattgtc ctaccattga cctctacaac tggacagcac
 661 ctcttgagtg gtttaagaat tgtcaggctc ttcaaggatc aaggtacagg gcgcacaagt
721 catttttggt cattgataat gtgatgactg aggacgcagg tgattacacc tgtaaattta
781 tacacaatga aaatggagcc aattatagtg tgacggcgac caggtccttc acggtcaagg
 841 tttggtgtca gagtttctgc aaattaaaaa agagcttaat ctttagtaat actcattgga
 901 ttcaaagtct aatgagaggc tttgtgatgg tatactatgg tgtacataaa tgttgtcgag
 961 tggtttttaa tctttgtttg caatactttc aacatcatca atggccttga atgagcaagg
1021 cttttctctg tttccagtaa tcggagcccc tgcacaaaat gaaataaagg aagtggaaat
1081 tggaaaaaac gcaaacctaa cttgctctgc ttgttttgqa aaaqqcactc agttcttgqc
1141 tgccgtcctg tggcagctta atggaacaaa aattacagac tttggtgaac caagaattca
1201 acaagaggaa gggcaaaatc aaagtttcag caatgggctg gcttgtctag acatggtttt
1261 aagaataget gacgtgaagg aagaggattt attgetgeag tacgactgte tggeeetgaa
1321 tttgcatggc ttgagaaggc acaccgtaag actaagtagg aaaaatccaa gtaaggagtg
1381 tttctgagac tttgatcacc tgaactttct ctagcaagtg taagcagaat ggagtgtggt
1441 tccaagagat ccatcaagac aatgggaatg gcctgtgcca taaaatgtgc ttctcttctt
1501 cgggatgttg tttgctgtct gatctttgta gactgttcct gtttgctggg agcttctctg
1561 etgettaaat tgttegteet eecceactee eteetategt tggtttgtet agaacactea
1621 gctgcttctt tggtcatcct tgttttctaa ctttatgaac tccctctgtg tcactgtatg
1681 tgaaaggaaa tgcaccaaca accgtaaact gaacgtgttc ttttgtgctc ttttataact
1741 tgcattacat gttgtaagca tggtccgttc tatacctttt tctggtcata atgaacactc
1801 attttgttag cgagggtggt aaagtgaaca aaaaggggaa gtatcaaact actgccattt
1861 cagtgagaaa atcctaggtg ctactttata ataagacatt tgttaggcca ttcttgcatt
1921 gatataaaga aatacctgag actgggtgat ttatatgaaa agaggtttaa ttggctcaca
1981 gttctgcagg ctgtatggga agcatggcgg catctgcttc tggggacacc tcaggagctt
2041 tactcatggc agaaggcaaa gcaaaggcag gcacttcaca cagtaaaagc aggagcgaga
2101 gagaggtgcc acactgaaac agccagatct catgagaagt cactcactat tgcaaggaca
2161 gcatcaaaga gatggtgcta aaccattcat gatgaactca cccccatgat ccaatcacct
2221 cccaccaggc tccacctcga atactgggga ttaccattca gcatgagatt tgggcaggaa
```

```
2281 cacagaccca aaccatacca cacacattat cattgttaaa ctttgtaaag tatttaaggt 2341 acatggaaca cacgggaagt ctggtagctc agcccatttc tttattgcat ctgttattca 2401 ccatgtaatt caggtaccac gtattccagg gagcctttct tggccctcag tttgcagtat 2461 acacactttc caagtactct tgtagcatcc tgtttgtatc atagcactgg tcacattgcc 2521 ttacctaaat ctgtttgaca gtctgctcaa cacgactgca agctccatga gggcagggac 2581 acatctctt ccatctttgg gtccttagtg caatacctgg cagctagcca gtgctcagct 2641 aaatatttgt tgactgaata aatgaatgca caaccaaaaa aaaaaaaaa aa
```

[0252] Following are show human amino acid sequences for isoform 1, isoform 2 and isoform 3 of IL1RL1.

#### IL1RL1 Amino Acid Sequence 1 (SEQ ID NO: 5)

NP\_057316 interleukin 1 receptor-like 1 isoform 1 precursor; interleukin 1 receptor-related protein; homolog of mouse growth stimulation-expressed gene; ST2 protein [Homo sapiens]

MGFWILAILTILMYSTAAKFSKQSWGLENEALIVRCPRQGKPSYTVDWYYSQTNKSIPT QERNRVFASGQLLKFLPAAVADSGIYTCIVRSPTFNRTGYANVTIYKKQSDCNVPDYLM YSTVSGSEKNSKIYCPTIDLYNWTAPLEWFKNCQALQGSRYRAHKSFLVIDNVMTEDA GDYTCKFIHNENGANYSVTATRSFTVKDEQGFSLFPVIGAPAQNEIKEVEIGKNANLTCS ACFGKGTQFLAAVLWQLNGTKITDFGEPRIQQEEGQNQSFSNGLACLDMVLRIADVKE EDLLLQYDCLALNLHGLRRHTVRLSRKNPIDHHSIYCIIAVCSVFLMLINVLVIILKMFWI EATLLWRDIAKPYKTRNDGKLYDAYVVYPRNYKSSTDGASRVEHFVHQILPDVLENKC GYTLCIYGRDMLPGEDVVTAVETNIRKSRRHIFILTPQITHNKEFAYEQEVALHCALIQN DAKVILIEMEALSELDMLQAEALQDSLQHLMKVQGTIKWREDHIANKRSLNSKFWKHV RYQMPVPSKIPRKASSLTPLAAQKQ

#### IL1RL1 Amino Acid Sequence 2 (SEO ID NO: 6)

NP\_003847 interleukin 1 receptor-like 1 isoform 2 precursor; interleukin 1 receptor-related protein; homolog of mouse growth stimulation-expressed gene; ST2 protein [Homo sapiens]

MGFWILAILTILMYSTAAKFSKQSWGLENEALIVRCPRQGKPSYTVDWYYSQTNKSIPT QERNRVFASGQLLKFLPAAVADSGIYTCIVRSPTFNRTGYANVTIYKKQSDCNVPDYLM YSTVSGSEKNSKIYCPTIDLYNWTAPLEWFKNCQALQGSRYRAHKSFLVIDNVMTEDA GDYTCKFIHNENGANYSVTATRSFTVKDEQGFSLFPVIGAPAQNEIKEVEIGKNANLTCS ACFGKGTQFLAAVLWQLNGTKITDFGEPRIQQEEGQNQSFSNGLACLDMVLRIADVKE EDLLLQYDCLALNLHGLRRHTVRLSRKNPSKECF

## IL1RL1 Amino Acid Sequence 3 (SEQ ID NO: 7)

NP\_775661 interleukin 1 receptor-like 1 isoform 3 precursor; interleukin 1 receptor-related protein; homolog of mouse growth stimulation-expressed gene; ST2 protein [Homo sapiens]

MGFWILAILTILMYSTAAKFSKQSWGLENEALIVRCPRQGKPSYTVDWYYSQTNKSIPT QERNRVFASGQLLKFLPAAVADSGIYTCIVRSPTFNRTGYANVTIYKKQSDCNVPDYLM YSTVSGSEKNSKIYCPTIDLYNWTAPLEWFKNCQALQGSRYRAHKSFLVIDNVMTEDA

## GDYTCKFIHNENGANYSVTATRSFTVKVWCQSFCKLKKSLIFSNTHWIQSLMRGFVMV YYGVHKCCRVVFNLCLQYFQHHQWP

[0253] Modifications may be made to the foregoing without departing from the basic aspects of the invention. Although the invention has been described in substantial detail with reference to one or more specific embodiments, those of skill in the art will recognize that changes may be made to the embodiments specifically disclosed in this application, yet these modifications and improvements are within the scope and spirit of the invention, as set forth in the aspects which follow. All publications or patent documents cited in this specification are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference.

[0254] Citation of the above publications or documents is not intended as an admission that any of the foregoing is pertinent prior art, nor does it constitute any admission as to the contents or date of these publications or documents. U.S. patents and other publications referenced herein are hereby incorporated by reference.

PTO/SB/16 (08-03)
Approved for use through 7/31/2006. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

# PROVISIONAL APPLICATION FOR PATENT COVER SHEET This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EL 961008095 US

	IN	VENTOR(S)			
Given Name (first and middle [if any] )	Family Name or Surname		· •	idence te or Foreign Country)	
Steven	MAH		San Diego, Californ		
Andreas	BRAUN		San Diego, Californ		
Stefan M.	KAMMERER		San Diego, Californ		
Additional inventors are being	r named on the	1			
Additional inventors are being	<del></del>	<del></del>	ely numbered sheets attach	ea nereto	
METHODS FOR IDENTIFYING	ITLE OF THE INVE			HEREOF	
Direct all correspondence to: COF	RRESPONDENCE	ADDRESS			
X Customer Number:	25225				
Firm or Individual Name			e		
Address					
City		State	Zip	1	
Country		Telephone	Fax		
ENCI	LOSED APPLICATI	ON PARTS (c	heck all that apply)		
X Specification Number of Pages 121 CD(s), Number   X Drawing(s) Number of Sheets 3 X Other   X Application Data Sheet. See 37 CFR 1.76 (4 pages) (specify): Return Receipt Postcard					
METHOD OF PAYMENT OF FILING	FEES FOR THIS P	ROVISIONAL A	APPLICATION FOR PATE	NT	
X Applicant claims small entity status. See 37 CFR 1.27.  A check or money order is enclosed to cover the filing fees.  The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 03-1952 80.00  Payment by credit card. Form PTO-2038 is attached.					
The invention was made by an agency of the United States Government or under a contract with an agency of the					
United States Government.  X No Yes, the name of the U.S. Government agency and the Government contract number are:					
[Page 1 of 2 ]  Respectfully submitted, Data A 114 0004					
SIGNATURE TYPED OR PRINTED NAME Bruce D. C.	Grant		REGISTRATION NO. (if appropriate)	April 1, 2004 47,608	
TELEPHONE (858) 720-	7962		Docket Number:	524593009000	
USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT					
I hereby certify that this correspondence is being deposited with the U.S. Postal Service as Express Mail, Airbill No. EL 961008095 US, in an envelope addressed to: Mail Stop Provisional Patent Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450, on the date shown below.					
Dated: 41104 Signature: Durrel Wy (Deborah Wykes)					

## PROVISIONAL APPLICATION COVER SHEET Additional Page

PTO/SB/16 (08-03)
Approved for use through 07/31/06. OMB 0651-0032
U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

	Docket Number 5	24593009000			
INVENTOR(S)/APPLICANT(S)					
Given Name (first and middle [if any])  Matthew Roberts Rikard Henry Maria L.	Family or Surname  NELSON RENELAND LANGDOWN	Residence (City and either State or Foreign Country)  San Marcos, California San Diego, California San Diego, California			

[Page 2 of 2]

1- 1